

TABLE E-1 Patient Demographics

	Entire Cohort (N = 71)	Skeletally Immature (N = 35)	Skeletally Mature (N = 36)
Age (yr)*			
At time of injury	21 (4 to 64)	11 (4 to 17)	31 (16 to 64)
At time of final follow-up	42 (19 to 81)	31 (19 to 47)	52 (34 to 81)
Sex			
Female	27 (38%)	17 (49%)	10 (28%)
Male	44 (62%)	18 (51%)	26 (72%)
Fracture side			
Left	39 (55%)	18 (51%)	21 (58%)
Right	32 (45%)	17 (49%)	15 (42%)
Injury to dominant side			
Yes	32 (45%)	15 (43%)	16 (44%)
No	39 (55%)	20 (57%)	20 (56%)
AO/OTA classification†			
22-A3	46 (65%)	31 (89%)	15 (42%)
22-B3	18 (25%)	4 (11%)	14 (39%)
22-C	7 (10%)	0 (0%)	7 (19%)
Associated ipsilateral fracture of upper limb			
Yes	11 (15%)	1 (3%)	10 (28%)
No	60 (85%)	34 (97%)	26 (72%)
Fracture severity			
Closed	47 (66%)	25 (71%)	22 (61%)
Open, grade I	17 (24%)	7 (20%)	10 (28%)
Open, grade II	2 (3%)	1 (3%)	1 (3%)
Open, grade III A	2 (3%)	1 (3%)	1 (3%)
Open, grade III B	2 (3%)	0 (0%)	2 (6%)
Open, grade III C	1 (1%)	1 (3%)	0 (0%)

\*Values are given as the mean, with the range in parentheses. †AO/OTA = Arbeitsgemeinschaft für Osteosynthesefragen/Orthopaedic Trauma Association.