

Appendix

References

1. Eisele R, Kinzl L, Koelsch T. Rapid-inflation intermittent pneumatic compression for prevention of deep venous thrombosis. *J Bone Joint Surg Am.* 2007;89:1050-6.
2. Kocher MS, Kasser JR, Waters PM, Bae D, Snyder BD, Hresko MT, Hedequist D, Karlin L, Kim YJ, Murray MM, Millis MB, Emans JB, Dichtel L, Matheney T, Lee BM. Lateral entry compared with medial and lateral entry pin fixation for completely displaced supracondylar humeral fractures in children. A randomized clinical trial. *J Bone Joint Surg Am.* 2007;89:706-12.
3. Parratte S, Argenson JN. Validation and usefulness of a computer-assisted cup-positioning system in total hip arthroplasty. A prospective, randomized, controlled study. *J Bone Joint Surg Am.* 2007;89:494-9.
4. Gerber JP, Marcus RL, Dibble LE, Greis PE, Burks RT, LaStayo PC. Effects of early progressive eccentric exercise on muscle structure after anterior cruciate ligament reconstruction. *J Bone Joint Surg Am.* 2007;89:559-70.
5. Matziolis G, Krocker D, Weiss U, Tohtz S, Perka C. A prospective, randomized study of computer-assisted and conventional total knee arthroplasty. Three-dimensional evaluation of implant alignment and rotation. *J Bone Joint Surg Am.* 2007;89:236-43.
6. Placzek R, Drescher W, Deuretzbacher G, Hempfing A, Meiss AL. Treatment of chronic radial epicondylitis with botulinum toxin A. A double-blind, placebo-controlled, randomized multicenter study. *J Bone Joint Surg Am.* 2007;89:255-60.
7. Poggie RA, Turgeon TR, Coutts RD. Failure analysis of a ceramic bearing acetabular component. *J Bone Joint Surg Am.* 2007;89:367-75.
8. Canadian Orthopaedic Trauma Society. Nonoperative treatment compared with plate fixation of displaced midshaft clavicular fractures. A multicenter, randomized clinical trial. *J Bone Joint Surg Am.* 2007;89:1-10.
9. Moroni A, Faldini C, Hoang-Kim A, Pegreffi F, Giannini S. Alendronate improves screw fixation in osteoporotic bone. *J Bone Joint Surg Am.* 2007;89:96-101.
10. Neumayr LD, Aguilar C, Earles AN, Jergesen HE, Haberkern CM, Kammen BF, Nancarrow PA, Padua E, Milet M, Stulberg BN, Williams RA, Orringer EP, Gruber N, Robertson SM, Vichinsky EP; National Osteonecrosis Trial in Sickle Cell Anemia Study Group. Physical therapy alone compared with core decompression and physical therapy for femoral head osteonecrosis in sickle cell disease. Results of a multicenter study at a mean of three years after treatment. *J Bone Joint Surg Am.* 2006;88:2573-82.
11. Baker RP, Squires B, Gargan MF, Bannister GC. Total hip arthroplasty and hemiarthroplasty in mobile, independent patients with a displaced intracapsular fracture of the femoral neck. A randomized, controlled trial. *J Bone Joint Surg Am.* 2006;88:2583-9.
12. Morgan SJ, Jeray KJ, Saliman LH, Miller HJ, Williams AE, Tanner SL, Smith WR, Broderick JS. Continuous infusion of local anesthetic at iliac crest bone-graft sites for postoperative pain relief. A randomized, double-blind study. *J Bone Joint Surg Am.* 2006;88:2606-12.
13. Canadian Orthopaedic Trauma Society. Open reduction and internal fixation compared with circular fixator application for bicondylar tibial plateau fractures. Results of a multicenter, prospective, randomized clinical trial. *J Bone Joint Surg Am.* 2006;88:2613-23.
14. Draganich L, Reider B, Rimington T, Piotrowski G, Mallik K, Nasson S. The effectiveness of self-adjustable custom and off-the-shelf bracing in the treatment of varus gonarthrosis. *J Bone Joint Surg Am.* 2006;88:2645-52.
15. Forward DP, Singh AK, Lawrence TM, Sithole JS, Davis TR, Oni JA. Preservation of the ulnar bursa within the carpal tunnel: does it improve the outcome of carpal tunnel surgery? A randomized, controlled trial. *J Bone Joint Surg Am.* 2006;88:2432-8.
- 16.

- Bahr R, Fossan B, Løken S, Engebretsen L. Surgical treatment compared with eccentric training for patellar tendinopathy (jumper's knee). A randomized, controlled trial. *J Bone Joint Surg Am.* 2006;88:1689-98.
- 17.** Digiovanni BF, Nawoczenski DA, Malay DP, Graci PA, Williams TT, Wilding GE, Baumhauer JF. Plantar fascia-specific stretching exercise improves outcomes in patients with chronic plantar fasciitis. A prospective clinical trial with two-year follow-up. *J Bone Joint Surg Am.* 2006;88:1775-81.
- 18.** Riew KD, Park JB, Cho YS, Gilula L, Patel A, Lenke LG, Bridwell KH. Nerve root blocks in the treatment of lumbar radicular pain. A minimum five-year follow-up. *J Bone Joint Surg Am.* 2006;88:1722-5.
- 19.** Jones AL, Bucholz RW, Bosse MJ, Mirza SK, Lyon TR, Webb LX, Pollak AN, Golden JD, Valentin-Opran A; BMP-2 Evaluation in Surgery for Tibial Trauma-Allograft (BESTT-ALL) Study Group. Recombinant human BMP-2 and allograft compared with autogenous bone graft for reconstruction of diaphyseal tibial fractures with cortical defects. A randomized, controlled trial. *J Bone Joint Surg Am.* 2006;88:1431-41.
- 20.** Iannotti JP, Codsi MJ, Kwon YW, Derwin K, Ciccone J, Brems JJ. Porcine small intestine submucosa augmentation of surgical repair of chronic two-tendon rotator cuff tears. A randomized, controlled trial. *J Bone Joint Surg Am.* 2006;88:1238-44.
- 21.** Drogset JO, Grøntvedt T, Robak OR, Mølster A, Viset AT, Engebretsen L. A sixteen-year follow-up of three operative techniques for the treatment of acute ruptures of the anterior cruciate ligament. *J Bone Joint Surg Am.* 2006;88:944-52.
- 22.** Busch CA, Shore BJ, Bhandari R, Ganapathy S, MacDonald SJ, Bourne RB, Rorabeck CH, McCalden RW. Efficacy of periarticular multimodal drug injection in total knee arthroplasty. A randomized trial. *J Bone Joint Surg Am.* 2006;88:959-63.
- 23.** Kelly MJ, Rumi MN, Kothari M, Parentis MA, Bailey KJ, Parrish WM, Pellegrini VD Jr. Comparison of the vastus-splitting and median parapatellar approaches for primary total knee arthroplasty: a prospective, randomized study. *J Bone Joint Surg Am.* 2006;88:715-20.
- 24.** Ly TV, Coetzee JC. Treatment of primarily ligamentous Lisfranc joint injuries: primary arthrodesis compared with open reduction and internal fixation. A prospective, randomized study. *J Bone Joint Surg Am.* 2006;88:514-20.
- 25.** Keating JF, Grant A, Masson M, Scott NW, Forbes JF. Randomized comparison of reduction and fixation, bipolar hemiarthroplasty, and total hip arthroplasty. Treatment of displaced intracapsular hip fractures in healthy older patients. *J Bone Joint Surg Am.* 2006;88:249-60.
- 26.** Stannard JP, Lopez-Ben RR, Volgas DA, Anderson ER, Busbee M, Karr DK, McGwin GR Jr, Alonso JE. Prophylaxis against deep-vein thrombosis following trauma: a prospective, randomized comparison of mechanical and pharmacologic prophylaxis. *J Bone Joint Surg Am.* 2006;88:261-6.
- 27.** Wang CJ, Wang JW, Ko JY, Weng LH, Huang CC. Three-year changes in bone mineral density around the knee after a six-month course of oral alendronate following total knee arthroplasty. A prospective, randomized study. *J Bone Joint Surg Am.* 2006;88:267-72.
- 28.** Hartrick CT, Martin G, Kantor G, Koncelik J, Manvelian G. Evaluation of a single-dose, extended-release epidural morphine formulation for pain after knee arthroplasty. *J Bone Joint Surg Am.* 2006;88:273-81.
- 29.** Vendittoli PA, Makinen P, Drolet P, Lavigne M, Fallaha M, Guertin MC, Varin F. A multimodal analgesia protocol for total knee arthroplasty. A randomized, controlled study. *J Bone Joint Surg Am.* 2006;88:282-9.
- 30.** Salk RS, Chang TJ, D'Costa WF, Soomekh DJ, Grogan KA. Sodium hyaluronate in the treatment of osteoarthritis of the ankle: a controlled, randomized, double-blind pilot study. *J Bone Joint Surg Am.* 2006;88:295-302.
- 31.** Egol KA, Paksimma N, Puopolo S, Klugman J, Hiebert R, Koval KJ. Treatment of external fixation pins about the wrist: a prospective, randomized trial. *J Bone Joint Surg Am.* 2006;88:349-54.
- 32.** Bohm ER, Bubbar V, Yong Hing K, Dzus A. Above and below-the-elbow plaster casts for distal forearm fractures in children. A randomized controlled trial. *J Bone Joint Surg Am.* 2006;88:1-8.
- 33.**

- Webb GR, Galpin RD, Armstrong DG. Comparison of short and long arm plaster casts for displaced fractures in the distal third of the forearm in children. *J Bone Joint Surg Am.* 2006;88:9-17.
- 34.** Ha YC, Jung WH, Kim JR, Seong NH, Kim SY, Koo KH. Prediction of collapse in femoral head osteonecrosis: a modified Kerboul method with use of magnetic resonance images. *J Bone Joint Surg Am.* 2006;88:35-40.
- 35.** Kim YH, Kim JS, Kim DY. Clinical outcome and rate of complications after primary total knee replacement performed with quadriceps-sparing or standard arthrotomy. *J Bone Joint Surg Br.* 2007;89:467-70.
- 36.** Kim YH, Kim JS, Yoon SH. Alignment and orientation of the components in total knee replacement with and without navigation support: a prospective, randomised study. *J Bone Joint Surg Br.* 2007;89:471-6.
- 37.** Spencer JM, Chauhan SK, Sloan K, Taylor A, Beaver RJ. Computer navigation versus conventional total knee replacement: no difference in functional results at two years. *J Bone Joint Surg Br.* 2007;89:477-80.
- 38.** Church JS, Scadden JE, Gupta RR, Cokis C, Williams KA, Janes GC. Embolic phenomena during computer-assisted and conventional total knee replacement. *J Bone Joint Surg Br.* 2007;89:481-5.
- 39.** Molloy DO, Archbold HA, Ogonda L, McConway J, Wilson RK, Beverland DE. Comparison of topical fibrin spray and tranexamic acid on blood loss after total knee replacement: a prospective, randomised controlled trial. *J Bone Joint Surg Br.* 2007;89:306-9.
- 40.** Albert JD, Meadeb J, Guggenbuhl P, Marin F, Benkalfate T, Thomazeau H, Chalès G. High-energy extracorporeal shock-wave therapy for calcifying tendinitis of the rotator cuff: a randomised trial. *J Bone Joint Surg Br.* 2007;89:335-41.
- 41.** Saudan M, Saudan P, Perneger T, Riand N, Keller A, Hoffmeyer P. Celecoxib versus ibuprofen in the prevention of heterotopic ossification following total hip replacement: a prospective randomised trial. *J Bone Joint Surg Br.* 2007;89:155-9.
- 42.** Blomfeldt R, Törnkvist H, Eriksson K, Söderqvist A, Ponzer S, Tidermark J. A randomised controlled trial comparing bipolar hemiarthroplasty with total hip replacement for displaced intracapsular fractures of the femoral neck in elderly patients. *J Bone Joint Surg Br.* 2007;89:160-5.
- 43.** Kim YH, Yoon SH, Kim JS. Changes in the bone mineral density in the acetabulum and proximal femur after cementless total hip replacement: alumina-on-alumina versus alumina-on-polyethylene articulation. *J Bone Joint Surg Br.* 2007;89:174-9.
- 44.** Uvehammer J, Kärrholm J, Carlsson L. Cemented versus hydroxyapatite fixation of the femoral component of the Freeman-Samuelson total knee replacement: a radiostereometric analysis. *J Bone Joint Surg Br.* 2007;89:39-44.
- 45.** Muller SD, Deehan DJ, Holland JP, Outterside SE, Kirk LM, Gregg PJ, McCaskie AW. Should we reconsider all-polyethylene tibial implants in total knee replacement? *J Bone Joint Surg Br.* 2006;88:1596-602.
- 46.** Middleton F, Coakes J, Umarji S, Palmer S, Venn R, Panayiotou S. The efficacy of intra-articular bupivacaine for relief of pain following arthroscopy of the ankle. *J Bone Joint Surg Br.* 2006;88:1603-5.
- 47.** Karunakar MA, Sen A, Bosse MJ, Sims SH, Goulet JA, Kellam JF. Indometacin as prophylaxis for heterotopic ossification after the operative treatment of fractures of the acetabulum. *J Bone Joint Surg Br.* 2006;88:1613-7.
- 48.** Brouwer RW, Bierma-Zeinstra SM, van Raaij TM, Verhaar JA. Osteotomy for medial compartment arthritis of the knee using a closing wedge or an opening wedge controlled by a Puddu plate. A one-year randomised, controlled study. *J Bone Joint Surg Br.* 2006;88:1454-9.
- 49.** Shetty N, Hamer AJ, Stockley I, Eastell R, Wilkinson JM. Clinical and radiological outcome of total hip replacement five years after pamidronate therapy. A trial extension. *J Bone Joint Surg Br.* 2006;88:1309-15.
- 50.** Karn NK, Singh GK, Kumar P, Shrestha B, Singh MP, Gowda MJ. Comparison between external fixation and sliding hip screw in the management of trochanteric fracture of the femur in Nepal. *J Bone Joint Surg Br.* 2006;88:1347-50. Erratum in: *J Bone Joint Surg Br.* 2007;89:421.
- 51.**

- Hallan G, Aamodt A, Furnes O, Skredderstuen A, Haugan K, Havelin LI, Palamed G compared with Palacos R with gentamicin in Charnley total hip replacement. A randomised, radiostereometric study of 60 hips. *J Bone Joint Surg Br.* 2006;88:1143-8.
- 52.** Kean J, Wigderowitz CA, Coventry DM. Continuous interscalene infusion and single injection using levobupivacaine for analgesia after surgery of the shoulder. A double-blind, randomised controlled trial. *J Bone Joint Surg Br.* 2006;88:1173-7.
- 53.** Kesteris U, Aspenberg P. Rinsing morcellised bone grafts with bisphosphonate solution prevents their resorption. A prospective randomised double-blinded study. *J Bone Joint Surg Br.* 2006;88:993-6. Erratum in: *J Bone Joint Surg Br.* 2007;89:421.
- 54.** Vendittoli PA, Lavigne M, Girard J, Roy AG. A randomised study comparing resection of acetabular bone at resurfacing and total hip replacement. *J Bone Joint Surg Br.* 2006;88:997-1002.
- 55.** Grübl A, Weissinger M, Brodner W, Gleiss A, Giurea A, Gruber M, Pöll G, Meisinger V, Gottsauer-Wolf F, Kotz R. Serum aluminium and cobalt levels after ceramic-on-ceramic and metal-on-metal total hip replacement. *J Bone Joint Surg Br.* 2006;88:1003-5.
- 56.** Grondal L, Broström E, Wretenberg P, Stark A. Arthrodesis versus Mayo resection: the management of the first metatarsophalangeal joint in reconstruction of the rheumatoid forefoot. *J Bone Joint Surg Br.* 2006;88:914-9.
- 57.** Girard J, Lavigne M, Vendittoli PA, Roy AG. Biomechanical reconstruction of the hip: a randomised study comparing total hip resurfacing and total hip arthroplasty. *J Bone Joint Surg Br.* 2006;88:721-6.
- 58.** Campbell DG, Duncan WW, Ashworth M, Mintz A, Stirling J, Wakefield L, Stevenson TM. Patellar resurfacing in total knee replacement: a ten-year randomised prospective trial. *J Bone Joint Surg Br.* 2006;88:734-9.
- 59.** Bhan S, Pankaj A, Malhotra R. One- or two-stage bilateral total hip arthroplasty: a prospective, randomised, controlled study in an Asian population. *J Bone Joint Surg Br.* 2006;88:298-303.
- 60.** Tanaka N, Sakahashi H, Ishii S, Kudo H. Comparison of two types of ulnar component in type-5 Kudo total elbow arthroplasty in patients with rheumatoid arthritis: a long-term follow-up. *J Bone Joint Surg Br.* 2006;88:341-4.
- 61.** Kalteis T, Handel M, Bäthis H, Perlick L, Tingart M, Grifka J. Imageless navigation for insertion of the acetabular component in total hip arthroplasty: is it as accurate as CT-based navigation? *J Bone Joint Surg Br.* 2006;88:163-7.
- 62.** Cobb J, Henckel J, Gomes P, Harris S, Jakopec M, Rodriguez F, Barrett A, Davies B. Hands-on robotic unicompartmental knee replacement: a prospective, randomised controlled study of the Acrobot system. *J Bone Joint Surg Br.* 2006;88:188-97.
- 63.** Kalairajah Y, Cossey AJ, Verrall GM, Ludbrook G, Spriggins AJ. Are systemic emboli reduced in computer-assisted knee surgery?: A prospective, randomised, clinical trial. *J Bone Joint Surg Br.* 2006;88:198-202. Erratum in: *J Bone Joint Surg Br.* 2006;88:1407.
- 64.** Khan RJ, Fick D, Yao F, Tang K, Hurworth M, Nivbrant B, Wood D. A comparison of three methods of wound closure following arthroplasty: a prospective, randomised, controlled trial. *J Bone Joint Surg Br.* 2006;88:238-42.
- 65.** Keene G, Simpson D, Kalairajah Y. Limb alignment in computer-assisted minimally-invasive unicompartmental knee replacement. *J Bone Joint Surg Br.* 2006;88:44-8.
- 66.** Tsumura N, Yoshiya S, Chin T, Shiba R, Kohso K, Doita M. A prospective comparison of clamping the drain or post-operative salvage of blood in reducing blood loss after total knee arthroplasty. *J Bone Joint Surg Br.* 2006;88:49-53.
- 67.** Costa ML, MacMillan K, Halliday D, Chester R, Shepstone L, Robinson AH, Donell ST. Randomised controlled trials of immediate weight-bearing mobilisation for rupture of the tendo Achillis. *J Bone Joint Surg Br.* 2006;88:69-77.
- 68.**

- Toftdahl K, Nikolajsen L, Haraldsted V, Madsen F, Tønnesen EK, Søballe K. Comparison of peri- and intraarticular analgesia with femoral nerve block after total knee arthroplasty: a randomized clinical trial. *Acta Orthop.* 2007;78:172-9.
- 69.** Andersen KV, Pfeiffer-Jensen M, Haraldsted V, Søballe K. Reduced hospital stay and narcotic consumption, and improved mobilization with local and intraarticular infiltration after hip arthroplasty: a randomized clinical trial of an intraarticular technique versus epidural infusion in 80 patients. *Acta Orthop.* 2007;78:180-6.
- 70.** Andersen LJ, Poulsen T, Krogh B, Nielsen T. Postoperative analgesia in total hip arthroplasty: a randomized double-blinded, placebo-controlled study on peroperative and postoperative ropivacaine, ketorolac, and adrenaline wound infiltration. *Acta Orthop.* 2007;78:187-92.
- 71.** Rutten MJ, Maresch BJ, Jager GJ, de Waal Malefijt MC. Injection of the subacromial-subdeltoid bursa: blind or ultrasound-guided? *Acta Orthop.* 2007;78:254-7.
- 72.** van der Heide HJ, Rijnberg WJ, Van Sorge A, Van Kampen A, Schreurs BW. Similar effects of rofecoxib and indomethacin on the incidence of heterotopic ossification after hip arthroplasty. *Acta Orthop.* 2007;78:90-4.
- 73.** Grohs JG, Schmidt M, Wanivenhaus A. Selective COX-2 inhibitor versus indomethacin for the prevention of heterotopic ossification after hip replacement: a double-blind randomized trial of 100 patients with 1-year follow-up. *Acta Orthop.* 2007;78:95-8.
- 74.** Kim KJ, Iwase M, Kotake S, Itoh T. Effect of bone marrow grafting on the titanium porous-coated implant in bilateral total knee arthroplasty. *Acta Orthop.* 2007;78:116-22.
- 75.** Kroon M, Visser CP, Mootanah R, Brand R. Performance of 3 gelatine-based resorbable cement plugs: a study on 15 synthetic femurs and a prospective randomized study on 103 patients. *Acta Orthop.* 2006;77:893-8.
- 76.** Hilding M, Aspenberg P. Postoperative clodronate decreases prosthetic migration: 4-year follow-up of a randomized radiostereometric study of 50 total knee patients. *Acta Orthop.* 2006;77:912-6.
- 77.** Geerdink CH, Grimm B, Ramakrishnan R, Rondhuis J, Verburg AJ, Tonino AJ. Crosslinked polyethylene compared to conventional polyethylene in total hip replacement: pre-clinical evaluation, in-vitro testing and prospective clinical follow-up study. *Acta Orthop.* 2006;77:719-25.
- 78.** Carlsson LV, Albrektsson BE, Albrektsson BG, Albrektsson TO, Jacobsson CM, Macdonald W, Regnér L, Röslund T, Weidenhielm LR. Stepwise introduction of a bone-conserving osseointegrated hip arthroplasty using RSA and a randomized study: I. Preliminary investigations--52 patients followed for 3 years. *Acta Orthop.* 2006;77:549-58.
- 79.** Carlsson LV, Albrektsson T, Albrektsson BE, Jacobsson CM, Macdonald W, Regnér L, Weidenhielm LR. Stepwise introduction of a bone-conserving osseointegrated hip arthroplasty using RSA and a randomized study: II. Clinical proof of concept--40 patients followed for 2 years. *Acta Orthop.* 2006;77:559-66.
- 80.** Hsieh PH, Chang YH, Shih CH. Image-guided periacetabular osteotomy: computer-assisted navigation compared with the conventional technique: a randomized study of 36 patients followed for 2 years. *Acta Orthop.* 2006;77:591-7.
- 81.** Nygaard M, Elling F, Bastholm L, Søballe K, Borgwardt A. No difference in early cellular response of the pseudo-synovial membrane after total hip arthroplasty: comparison of 3 combinations of bearing materials. *Acta Orthop.* 2006;77:402-12.
- 82.** Atroshi I, Brogren E, Larsson GU, Kloow J, Hofer M, Berggren AM. Wrist-bridging versus non-bridging external fixation for displaced distal radius fractures: a randomized assessor-blind clinical trial of 38 patients followed for 1 year. *Acta Orthop.* 2006;77:445-53.
- 83.** Olivecrona C, Tidermark J, Hamberg P, Ponzer S, Cederfjäll C. Skin protection underneath the pneumatic tourniquet during total knee arthroplasty: a randomized controlled trial of 92 patients. *Acta Orthop.* 2006;77:519-23.
- 84.** Digas G, Kärrholm J, Thanner J. Different loss of BMD using uncemented press-fit and whole polyethylene cups fixed with cement: repeated DXA studies in 96 hips randomized to 3 types of fixation. *Acta Orthop.* 2006;77:218-26.
- 85.**

- Mattsson P, Larsson S. Calcium phosphate cement for augmentation did not improve results after internal fixation of displaced femoral neck fractures: a randomized study of 118 patients. *Acta Orthop.* 2006;77:251-6.
- 86.** Ström H, Kolstad K, Mallmin H, Sahlstedt B, Milbrink J. Comparison of the uncemented Cone and the cemented Bimetric hip prosthesis in young patients with osteoarthritis: an RSA, clinical and radiographic study. *Acta Orthop.* 2006;77:71-8.
- 87.** Hagsten B, Svensson O, Gardulf A. Health-related quality of life and self-reported ability concerning ADL and IADL after hip fracture: a randomized trial. *Acta Orthop.* 2006;77:114-9.
- 88.** Matsuda K, Nakamura S, Wakimoto N, Kobayashi M, Matsushita T. Drainage does not increase anemia after cementless total hip arthroplasty. *Clin Orthop Relat Res.* 2007;458:101-5.
- 89.** Ensini A, Catani F, Leardini A, Romagnoli M, Giannini S. Alignments and clinical results in conventional and navigated total knee arthroplasty. *Clin Orthop Relat Res.* 2007;457:156-62.
- 90.** de Jong M, Ray M, Crawford S, Whitehouse SL, Crawford RW. Platelet and leukocyte activation in salvaged blood and the effect of its reinfusion on the circulating blood. *Clin Orthop Relat Res.* 2007;456:238-42.
- 91.** Gioe TJ, Stroemer ES, Santos ER. All-polyethylene and metal-backed tibias have similar outcomes at 10 years: a randomized level I [corrected] evidence study. *Clin Orthop Relat Res.* 2007;455:212-8. Erratum in: *Clin Orthop Relat Res.* 2007;458:249.
- 92.** Ben-Galim P, Rosenblatt Y, Parnes N, Dekel S, Steinberg EL. Intramedullary fixation of tibial shaft fractures using an expandable nail. *Clin Orthop Relat Res.* 2007;455:234-40.
- 93.** Yagi M, Kuroda R, Nagamune K, Yoshiya S, Kurosaka M. Double-bundle ACL reconstruction can improve rotational stability. *Clin Orthop Relat Res.* 2007;454:100-7.
- 94.** Ström H, Nilsson O, Milbrink J, Mallmin H, Larsson S. Early migration pattern of the uncemented CLS stem in total hip arthroplasties. *Clin Orthop Relat Res.* 2007;454:127-32.
- 95.** Kraay MJ, Thomas RD, Rimnac CM, Fitzgerald SJ, Goldberg VM. Zirconia versus Co-Cr femoral heads in total hip arthroplasty: early assessment of wear. *Clin Orthop Relat Res.* 2006;453:86-90.
- 96.** Aglietti P, Baldini A, Sensi L. Quadriceps-sparing versus mini-subvastus approach in total knee arthroplasty. *Clin Orthop Relat Res.* 2006;452:106-11.
- 97.** Heiss C, Meissner SA, Hoesel LM, Pfeil J, Schnettler R. Posteromedial approach to proximal tibia for corticotomy in callus distractions. *Clin Orthop Relat Res.* 2006;451:182-8.
- 98.** Eneroth M, Olsson UB, Thorngren KG. Nutritional supplementation decreases hip fracture-related complications. *Clin Orthop Relat Res.* 2006;451:212-7.
- 99.** Rahme H, Mattsson P, Wikblad L, Larsson S. Cement and press-fit humeral stem fixation provides similar results in rheumatoid patients. *Clin Orthop Relat Res.* 2006;448:28-32.
- 100.** Zhou ZK, Li MG, Börlin N, Wood DJ, Nivbrant B. No increased migration in cups with ceramic-on-ceramic bearing: an RSA study. *Clin Orthop Relat Res.* 2006;448:39-45.
- 101.** Digas G, Kärrholm J, Thanner J. Addition of fluoride to acrylic bone cement does not improve fixation of a total hip arthroplasty stem. *Clin Orthop Relat Res.* 2006;448:58-66.
- 102.** Luites JW, Spruit M, Hellemond GG, Horstmann WG, Valstar ER. Failure of the uncoated titanium ProxiLock femoral hip prosthesis. *Clin Orthop Relat Res.* 2006;448:79-86.
- 103.** Henricson A, Dalén T, Nilsson KG. Mobile bearings do not improve fixation in cemented total knee arthroplasty. *Clin Orthop Relat Res.* 2006;448:114-21.
- 104.** van der Linde MJ, Garling EH, Valstar ER, Tonino AJ, Nelissen RG. Periapatite may not improve micromotion of knee prostheses in rheumatoid arthritis. *Clin Orthop Relat Res.* 2006;448:122-8.
- 105.**

- Nilsson KG, Henricson A, Norgren B, Dalén T. Uncemented HA-coated implant is the optimum fixation for TKA in the young patient. *Clin Orthop Relat Res.* 2006;448:129-39.
- 106.** Flivik G, Kristiansson I, Kesteris U, Ryd L. Is removal of subchondral bone plate advantageous in cemented cup fixation? A randomized RSA study. *Clin Orthop Relat Res.* 2006;448:164-72.
- 107.** Faes M, van den Akker B, de Lint JA, Kooloos JG, Hopman MT. Dynamic extensor brace for lateral epicondylitis. *Clin Orthop Relat Res.* 2006;442:149-57.
- 108.** Lang T, Barker R, Steinlechner B, Gustorff B, Puskas T, Gore O, Kober A. TENS relieves acute posttraumatic hip pain during emergency transport. *J Trauma.* 2007;62:184-8.
- 109.** Helttula I, Karanko M, Gullichsen E. Similar central hemodynamics but increased postoperative oxygen consumption in unreamed versus reamed intramedullary nailing of femoral fractures. *J Trauma.* 2006;61:1178-85.
- 110.** Stannard JP, Robinson JT, Anderson ER, McGwin G Jr, Volgas DA, Alonso JE. Negative pressure wound therapy to treat hematomas and surgical incisions following high-energy trauma. *J Trauma.* 2006;60:1301-6.
- 111.** Rajan GP, Fornaro J, Trentz O, Zellweger R. Cancellous allograft versus autologous bone grafting for repair of comminuted distal radius fractures: a prospective, randomized trial. *J Trauma.* 2006;60:1322-9.
- 112.** Harris IA, Kadir A, Donald G. Continuous compartment pressure monitoring for tibia fractures: does it influence outcome? *J Trauma.* 2006;60:1330-5.