Copyright @ by The Journal of Bone and Joint Surgery, Incorporated

MEFTAH ET AL.

 $Long-Term\ Results\ of\ Intralesional\ Curettage\ and\ Cryosurgery\ for\ Treatment\ of\ Low-Grade\ Chondrosarcoma\ http://dx.doi.org/10.2106/JBJS.L.00442$ 

Page 1 of 1

TABLE E-1 Published Studies with a Minimum One Year of Follow-up After Cryosurgery for Treatment of Low-Grade Chondrosarcoma Without Soft-Tissue Extension

Mandat Cont Fiscas Extension				
Study	Year	No. of Cases	Minimum/Mean Follow-up (yr)	Recurrence (no. [%])
Current study	2012	43	5/10	4* (9)
Mohler et al. <sup>7</sup>	2010	52	1.5/4	2 (4)
Donati et al. <sup>15</sup>	2010	15	5.5/13	2 (13)
Souna et al.8	2010	15	5/8	0 (0)
Aarons et al. <sup>5</sup>	2009	17	2/4.6	1 (6)
van der Geest et al. <sup>9</sup>	2008	55	2/5	0 (0)
Ahlmann et al. <sup>10</sup>	2006	10	2/3.2	0 (0)
Schreuder et al. <sup>22</sup>	1998	9	1.2/2.1	0 (0)
Marcove et al.31	1977	7	4.9/6.5	0 (0)

<sup>\*</sup>All four recurrences were in patients who had had soft-tissue extension of the tumor.