

TABLE E-I Patient Data

Patient	Diagnosis	Age at Onset (Years)	Follow-up (Months)	Sex	Hip Pain	Gait ($\pm$ LLD in cm)	HHS	IHR	Global PODCI	Stulberg Class	Ficat Stage
1	SCFE	15.3	51.9	M	Yes	Normal	65	83	77	IV	III
2	SCFE	13.3	27.4	M	No	Normal	87	89	78	III	III
3	SCFE	15.5	47.0	M	No	Normal (1.0)	100	100	100	II	I
4	SCFE	11.2	43.4	F	No	Normal	100	98	95	I	I
5	SCFE	11.3	25.6	F	No	Normal	90	82	86	III	III
6	SCFE	11.8	48.4	M	No	Normal (1.0)	75	66	86	II	I
7	SCFE	10.1	32.3	F	No	Normal	96	93	92	II	I
8	SCFE	13.8	30.2	M	No	Normal (1.0)	100	100	100	I	I
9	SCFE	12.3	26.1	M	No	Normal	100	100	100	I	I
10	SCFE	13.4	48.9	M	No	Trendelenburg	97	98	100	II	II
11	SCFE	12.3	58.3	M	No	Short leg (2.5)	97	96	100	III	III
12	SCFE	13.2	27.2	M	No	Normal	98	96	94	III	III
13	Femoral neck fracture	10.5	42.0	F	No	Normal (1.0)	96	93	94	I	I
14	Femoral neck fracture	12.2	51.6	M	Yes	Short leg (2.0)	70	86	81	IV	III
15	Femoral neck fracture	13.8	28.0	M	No	Short leg (2.0)	93	92	94	III	III
16	Femoral neck fracture	11.7	39.0	M	Yes	Normal	86	93	81	III	III
17	Hip dislocation	14.0	30.4	M	No	Normal	100	100	97	I	I

SCFE = Slipped capital femoral epiphysis; LLD = Limb-length discrepancy; HHS = Harris hip score; IHR = Iowa Hip Rating; PODCI = Pediatric Outcomes Data Collection Instrument.




Fig. E-1

Consensus Stulberg classification as applied in this study²⁰.