

Table E1. Data on the patients with septic arthritis of the glenohumeral joint

Case	Age (yr)/sex	Injection history	Symptom duration (wk)	Organism	Arthroscopic stage*	Associated lesion†	Underlying disease‡	Complications	Follow-up (mo)	Modified UCLA score ⁷
1	33/M	1	3	<i>Pseudomonas</i>	II			Recurrence	12	35
2	23/M	1	2	<i>Pseudomonas</i>	I	Labral tear			16	35
3	55/F	4	4	<i>S. aureus</i>	III	LHB deg	RA	Recurrence	16	26
4	56/F	1	2	No growth	III	RCT	DM		18	26
5	62/M	1	2	<i>S. aureus</i>	II	RCT	DM		12	24
6	73/M	1	3	<i>S. aureus</i>	III	RCT			16	18
7	59/M	1	8	<i>S. aureus</i>	IV	RCT	Hodgkin dis.	Recurrence→open débrid.	24	9
8	71/M	1	2	<i>S. aureus</i>	III	RCT	HF+DM		18	29
9	66/M	1	3	<i>Pseudomonas</i>	III				25	28
10	43/M	5	4	<i>S. aureus</i>	III	RCT	DM	Recurrence→open débrid	18	18
11	64/M		2	<i>Pneumococcus</i>	III	RCT			14	35
12	61/M	1	2	<i>Acinetobacter</i>	II	LHB deg	DM	Recurrence	12	35
13	57/M		3	No growth	II	LHB deg	Septic knee		14	28
14	70/M		3	No growth	II	RCT	DM		12	26
15	38/M	1	4	No growth	II	RCT	LC		16	24
16	69/M	30	6	Methicillin-sensitive <i>S. aureus</i>	IV		Tbc		14	16
17	68/M		1	No growth	III				29	29
18	85/M	1	2	No growth	II	RCT	Tbc		12	22
19	76/M	1	4	<i>E. coli</i>	III	RCT	HF		12	30
Average	59		3						16.4	26

*Arthroscopic staging based on the modified criteria of Gächter⁵. †LHB deg = long head of biceps degeneration, and RCT = rotator cuff tear. ‡RA = rheumatoid arthritis, DM = diabetes mellitus, LC = liver cirrhosis, Tbc = pulmonary tuberculous, and HF = heart failure.