COPYRIGHT © BY THE JOURNAL OF BONE AND JOINT SURGERY, INCORPORATED HUANG ET AL. INTRAVENOUS AND TOPICAL TRANEXAMIC ACID ALONE ARE SUPERIOR TO TOURNIQUET USE FOR PRIMARY TOTAL KNEE ARTHROPLASTY. A PROSPECTIVE, RANDOMIZED CONTROLLED TRIAL http://dx.doi.org/10.2106/JBJS.16.01525

Page 1

Appendix

TABLE E-1 VAS Pain Scores at Different Time Points

	Mean and Standard Deviation			P Value*				
Time Point†	Group A (N = 50)	Group B (N = 50)	Group C (N = 50)	A vs. B vs. C	A vs. B	A vs. C	B vs. C	
Preop.	4.28 ± 1.26	4.18 ± 1.41	4.08 ± 1.23	0.745	0.701	0.444	0.701	
POD 1	3.12 ± 0.87	2.84 ± 1.28	3.60 ± 1.34	0.006	0.239	0.044	0.002	
POD 2	2.66 ± 0.80	2.14 ± 1.13	3.12 ± 1.06	< 0.001	0.011	0.024	< 0.001	
POD 3	2.28 ± 0.67	1.88 ± 0.92	2.66 ± 0.99	< 0.001	0.021	0.029	< 0.001	
POD 5	1.74 ± 0.78	1.26 ± 0.49	2.02 ± 0.91	< 0.001	0.002	0.063	< 0.001	
1 mo	0.84 ± 0.65	0.68 ± 0.51	1.02 ± 0.92	0.061	0.263	0.208	0.018	
3 mo	0.52 ± 0.54	0.48 ± 0.50	0.64 ± 0.63	0.337	0.723	0.288	0.157	
6 mo	0.36 ± 0.48	0.32 ± 0.47	0.40 ± 0.73	0.785	0.728	0.728	0.487	

*P values were calculated using 1-way ANOVA and the Tukey post hoc multiple comparison test. †POD = postoperative day.

Copyright O by The Journal of Bone and Joint Surgery, Incorporated Huang et al.

INTRAVENOUS AND TOPICAL TRANEXAMIC ACID ALONE ARE SUPERIOR TO TOURNIQUET USE FOR PRIMARY TOTAL KNEE ARTHROPLASTY. A PROSPECTIVE, RANDOMIZED CONTROLLED TRIAL

http://dx.doi.org/10.2106/JBJS.16.01525

Page 2

TABLE E-2 Inflammator	v Biomarker Level	s at Different	Time Points
	y Diomarker Level	b at Different	1 me 1 omus

	Mean and Standard Deviation			P Value*				
Time Point+	Group A (N = 50)	Group B (N = 50)	Group C (N = 50)	A vs. B vs. C	A vs. B	A vs. C	B vs. C	
CRP								
Preop.	3.0 ± 2.0	3.0 ± 2.2	3.1 ± 1.8	0.992	0.924	0.903	0.980	
POD 1	35.7 ± 16.6	26.0 ± 17.5	48.7 ± 22.9	< 0.001	0.012	0.001	< 0.001	
POD 2	58.0 ± 24.0	42.4 ± 29.4	98.9 ± 48.8	< 0.001	0.031	< 0.001	< 0.001	
POD 3	43.1 ± 22.9	30.6 ± 20.6	80.4 ± 40.6	< 0.001	0.036	< 0.001	< 0.001	
POD 5	35.1 ± 17.6	25.8 ± 16.9	64.9 ± 34.8	< 0.001	0.059	< 0.001	< 0.001	
IL-6								
Preop.	4.4 ± 1.3	4.6 ± 1.2	4.3 ± 1.2	0.526	0.395	0.823	0.283	
POD 1	40.0 ± 16.0	28.5 ± 18.0	91.6 ± 23.4	< 0.001	0.004	< 0.001	< 0.001	
POD 2	33.6 ± 12.6	21.5 ± 14.7	58.7 ± 15.7	< 0.001	< 0.001	< 0.001	< 0.001	
POD 3	20.8 ± 9.6	14.1 ± 10.1	30.5 ± 9.6	< 0.001	0.001	< 0.001	< 0.001	
POD 5	12.5 ± 5.6	10.7 ± 6.6	18.6 ± 8.1	< 0.001	0.201	< 0.001	< 0.001	

*P values were calculated using 1-way ANOVA and the Tukey post hoc multiple comparison test. †POD = postoperative day.

COPYRIGHT © BY THE JOURNAL OF BONE AND JOINT SURGERY, INCORPORATED HUANG ET AL. INTRAVENOUS AND TOPICAL TRANEXAMIC ACID ALONE ARE SUPERIOR TO TOURNIQUET USE FOR PRIMARY TOTAL KNEE ARTHROPLASTY. A PROSPECTIVE, RANDOMIZED CONTROLLED TRIAL http://dx.doi.org/10.2106/JBJS.16.01525 Page 3

TABLE E-3 Complications

	No. of Patients			P Value*				
	Group A (N =	Group B (N =	Group C (N =	A vs. B vs.	A vs.	A vs.	B vs.	
Complications	50)	50)	50)	С	В	С	С	
DVT	0	0	0	—	—	—	—	
PE	0	0	0	—	—	—	—	
Intramuscular venous	6	4	3	0.672	0.741	0.487	1	
thrombosis								
Superficial infection	1	0	3	0.324	1	0.617	0.242	
Wound secretion	6	0	9	0.004	0.027	0.577	0.003	
Wound blistering	0	0	3	0.107		0.242	0.242	

*P values were calculated using the Pearson chi-square test or Fisher exact test.