
Copyright @ by The Journal of Bone and Joint Surgery, Incorporated Moewis et al.

RETENTION OF POSTERIOR CRUCIATE LIGAMENT ALONE MAY NOT ACHIEVE PHYSIOLOGICAL KNEE JOINT KINEMATICS AFTER TOTAL KNEE ARTHROPLASTY. A RETROSPECTIVE STUDY http://dx.doi.org/10.2106/JBJS.20.00024 Page 1

## The following content was supplied by the authors as supporting material and has not been copy-edited or verified by JBJS.


**Appendix Figure\_1**: Unloaded flexion-extension at the medial condyle.



Copyright  $\circledcirc$  by The Journal of Bone and Joint Surgery, Incorporated Moewis et al.

RETENTION OF POSTERIOR CRUCIATE LIGAMENT ALONE MAY NOT ACHIEVE PHYSIOLOGICAL KNEE JOINT KINEMATICS AFTER TOTAL KNEE ARTHROPLASTY. A RETROSPECTIVE STUDY http://dx.doi.org/10.2106/JBJS.20.00024 Page 2


Appendix Figure\_2: Unloaded flexion-extension at the lateral condyle.



Copyright @ by The Journal of Bone and Joint Surgery, Incorporated Moewis et al.

RETENTION OF POSTERIOR CRUCIATE LIGAMENT ALONE MAY NOT ACHIEVE PHYSIOLOGICAL KNEE JOINT KINEMATICS AFTER TOTAL KNEE ARTHROPLASTY. A RETROSPECTIVE STUDY http://dx.doi.org/10.2106/JBJS.20.00024 Page 3

## **Appendix Figure\_3**: Loaded lunge at the medial condyle.



Copyright @ by The Journal of Bone and Joint Surgery, Incorporated Moewis et al.

RETENTION OF POSTERIOR CRUCIATE LIGAMENT ALONE MAY NOT ACHIEVE PHYSIOLOGICAL KNEE JOINT KINEMATICS AFTER TOTAL KNEE ARTHROPLASTY. A RETROSPECTIVE STUDY http://dx.doi.org/10.2106/JBJS.20.00024 Page 4

## **Appendix Figure\_4**: Loaded lunge at the lateral condyle.

