STADHOUDER ET AL.

HIGH FAILURE RATES OF A UNILATERAL POSTERIOR PERI-APICAL DISTRACTION DEVICE (APIFIX) FOR FUSIONLESS TREATMENT OF ADOLESCENT IDIOPATHIC SCOLIOSIS

http://dx.doi.org/10.2106/JBJS.20.02176

| http://dx.doi.org/10.2100/3DJ3.20.02170 | | |
|--|--|-------------------------------------|
| Question Text | Author Response | Custom Submission Question ID |
| Please complete the following questions in accordance with the International Committee of Medical Journal Editors' recommendations on data sharing in clinical trials (guidelines and examples are available here Will individual participant data be available (including data dictionaries)? | yes | 54 |
| What data in particular will be shared? (Examples include all individual participant data after deidentification, only participant data that underlies the results, or not available.) | all data gathered after deidentification | 55 |
| What other documents will be available? | study protocol, informed consent, clinical study reports | 56 |
| When will data be available (start and end dates)? | 3 months to 5 years after publication | 57 |
| With whom? (Examples include anyone who wishes to access the data, researchers who provide a proposal, or not applicable.) | investigators whose proposed use of the data has been approved by an independent review committee | 58 |
| For what types of analyses? | for individual participant meta analysis | 59 |
| By what mechanism will data be made available? | proposals may be submitted up to 36 months following article publication. After 36 months data will be available at the University warehouse but without investigator support. | 60 |