The following content was supplied by the authors as supporting material and has not been copy-edited or verified by JBJS.

Appendix 1. Characteristics of the 37 patients who received treatment in infancy

Patient	Type of treatment received	Age when treatment	Age at study	Primary	Secondary
		was initiated (weeks)	radiograph (years)	outcome	outcome
1	Pavlik harness	3.3	4.5	Yes	Yes
2	Pavlik harness	3.6	5.4	Yes	Yes
3	Pavlik harness	8.7	3.2	Yes	Yes
4	Pavlik harness	1.4	5.3	No	Yes
5	Pavlik harness	3.4	4.0	No	Yes
6	Pavlik harness	6.0	5.3	No	Yes
7	Pavlik harness	7.3	3.2	No	Yes
8	Pavlik harness	7.9	4.6	No	Yes
9	Pavlik harness	10.4	4.2	No	Yes
10	Pavlik harness	12.7	5.5	No	Yes
11	Pavlik harness	14.6	5.1	No	Yes
12	Closed reduction, innominate osteotomy	27.9	6.0	No	Yes
13	Pavlik harness	0.3	2.5	No	No
14	Pavlik harness	0.9	5.0	No	No
15	Pavlik harness	1.4	5.6	No	No
16	Pavlik harness	2.7	6.1	No	No
17	Pavlik harness	2.7	3.9	No	No
18	Pavlik harness	4.6	6.5	No	No
19	Pavlik harness	6.9	4.4	No	No
20	Pavlik harness	7.3	4.5	No	No
21	Pavlik harness	8.3	4.2	No	No
22	Pavlik harness	8.6	5.6	No	No
23	Pavlik harness	8.6	5.3	No	No
24	Pavlik harness	9.7	5.0	No	No
25	Pavlik harness	10.6	6.3	No	No
26	Pavlik harness	11.3	3.9	No	No
27	Pavlik harness	11.9	5.0	No	No
28	Semi-rigid harness	7.0	4.5	No	No
29	Semi-rigid harness	7.1	2.0	No	No

Copyright © 2020 The Authors. Published by The Journal of Bone and Joint Surgery, Incorporated. HUMPHRY ET AL. PREDICTORS OF HIP DYSPLASIA AT 4 YEARS IN CHILDREN WITH PERINATAL RISK FACTORS http://dx.doi.org/10.2106/JBJS.OA.20.00108 Page 2

30	Semi-rigid harness	8.9	2.8	No	No
31	Semi-rigid and Pavlik harness	5.6	5.4	No	No
32	Semi-rigid and Pavlik harness	10.6	4.5	No	No
33	Pavlik harness, closed reduction, spica cast	4.4	4.6	No	No
34	Pavlik harness, closed reduction, spica cast	5.1	4.9	No	No
35	Pavlik harness, closed reduction, spica cast	12.1	6.3	No	No
36	Pavlik harness, closed reduction, spica cast	12.4	6.6	No	No
37	Closed reduction, spica cast	25.7	4.3	No	No

Copyright © 2020 The Authors. Published by The Journal of Bone and Joint Surgery, Incorporated. HUMPHRY ET AL. PREDICTORS OF HIP DYSPLASIA AT 4 YEARS IN CHILDREN WITH PERINATAL RISK FACTORS http://dx.doi.org/10.2106/JBJS.OA.20.00108 Page 3

Appendix 2. Comparing the characteristics of 1053 patients who received treatment in the postnatal period to those without a past medical history of DDH

Predictor		Not-treated (n=1016)	Treated (n=37)
Primary outcome	No	992 (97.6%)	34 (91.9%)
	Yes	24 (2.4%)	3 (8.1%)
Secondary outcome	No	882 (86.8%)	25 (67.6%)
	Yes	134 (13.2%)	12 (32.4%)
Gender	Male	495 (48.7%)	6 (16.2%)
	Female	521 (51.3%)	31 (83.8%)
Parity	Primiparous	540 (53.1%)	22 (59.5%)
	Multiparous	448 (44.1%)	12 (32.4%)
	Missing	28 (2.8%)	3 (8.1%)
Twin	No	856 (84.3%)	34 (91.9%)
	Yes	146 (14.4%)	-
	Missing	14 (1.4%)	3 (8.1%)
Mode of delivery	Cesarean	565 (55.6%)	14 (37.8%)
	Vaginal	436 (42.9%)	21 (56.8%)
	Missing	15 (1.5%)	2 (5.4%)
First degree relative with DDH	No	931 (91.6%)	32 (86.5%)
	Yes	85 (8.4%)	5 (13.5%)
Breech presentation	No	692 (68.1%)	26 (70.3%)
	Yes	324 (31.9%)	11 (29.7%)
Foot Deformity	No	986 (97%)	35 (94.6%)
	Yes	30 (3%)	2 (5.4%)
Ortolani or Barlow positive, abduction	No	990 (97.4%)	13 (35.1%)
asymmetry, leg length difference	Yes	26 (2.6%)	24 (64.9%)
Weight (kg)		3.22 (0.58)	3.39 (0.50)