
**Appendix A**


| **Reliability: Intraclass Correlation Coefficients for Intra-rater, Inter-rater, Test Retest 1-hour, and Test Retest 1-week Scores** | | | |
|---|---|---|---|
| **Intra-rater Reliability** | **ICC (2,1)** | **95% CI** | **p-value** |
| **Session 1, Trials 1 and 2** | | | |
| Rater A | 1.00 | 1.00 - 1.00 | <0.001 |
| Rater B | 1.00 | 1.00 - 1.00 | <0.001 |
| Rater C | 1.00 | 1.00 - 1.00 | <0.001 |
| **Session 2, Trials 1 and 2** | | | |
| Rater A | 1.00 | 1.00 - 1.00 | <0.001 |
| Rater B | 1.00 | 1.00 - 1.00 | <0.001 |
| Rater C | 1.00 | 1.00 - 1.00 | <0.001 |
| **Session 3, Trials 1 and 2** | | | |
| Rater A | 0.99 | 0.98 - 1.00 | <0.001 |
| **Inter-rater Reliability** | **ICC (2,1)** | **95% CI** | **p-value** |
| **Session 1, Trial 1** | | | |
| Raters A and B | 1.00 | 1.00 - 1.00 | <0.001 |
| Raters A and C | 1.00 | 1.00 - 1.00 | <0.001 |
| Raters B and C | 1.00 | 1.00 - 1.00 | <0.001 |
| **Session 1, Trial 2** | | | |
| Raters A and B | 1.00 | 1.00 - 1.00 | <0.001 |
| Raters A and C | 1.00 | 1.00 - 1.00 | <0.001 |
| Raters B and C | 1.00 | 1.00 - 1.00 | <0.001 |
| **Session 2, Trial 1** | | | |
| Raters A and B | 1.00 | 1.00 - 1.00 | <0.001 |
| Raters A and C | 1.00 | 1.00 - 1.00 | <0.001 |
| Raters B and C | 1.00 | 1.00 - 1.00 | <0.001 |
| **Session 2, Trial 2** | | | |
| Raters A and B | 1.00 | 1.00 - 1.00 | <0.001 |
| Raters A and C | 1.00 | 1.00 - 1.00 | <0.001 |
| Raters B and C | 1.00 | 1.00 - 1.00 | <0.001 |
| **1-hour Test Retest Reliability** | **ICC (2,2)** | **95% CI** | **p-value** |
| **Sessions 1 and 2 (n=22)** | | | |
| Rater A | 0.99 | 0.97 - 0.99 | <0.001 |
| Rater B | 0.99 | 0.97 - 0.99 | <0.001 |
| Rater C | 0.99 | 0.97 - 1.00 | <0.001 |
| **1-week Test Retest Reliability** | **ICC (2,2)** | **95% CI** | **p-value** |
| **Sessions 1 and 3 (n=21)** | | | |
| Rater A | 0.99 | 0.99 – 1.00 | <0.001 |

ICC (2,1) = SPSS, Model 2, repeated measures ANOVA, single measures
ICC (2,2) = SPSS, Model 2, repeated measures ANOVA, average measures, two trials per session

**Appendix B**

**Box plot analysis with interquartile range and outliers for 1-hour and 1 week difference in clonus duration**



**Appendix C**

**Test-retest reliability of drop test 10-sec oscillations (n = 22; ICC 2, 1)**

| TRIAL | 1 | 2 | 3 | 4 | 5 |
|---|---|---|---|---|---|
| **1** | **NA** | **0.98**<br>0.97-0.99 | **0.90**<br>0.76-0.96 | **0.90**<br>0.76-9.96 | **0.79**<br>0.52-0.91 |
| **2** | **0.98**<br>0.97-0.99 | **NA** | **0.96**<br>0.91-0.98 | **0.99**<br>0.99-1.00 | **0.85**<br>0.65-0.94 |
| **3** | **0.90**<br>0.76-0.96 | **0.96**<br>0.91-0.98 | **NA** | **0.97**<br>0.91-0.98 | **0.87**<br>0.70-0.94 |
| **4** | **0.90**<br>0.76-0.96 | **0.99**<br>0.99-1.00 | **0.97**<br>0.91-0.98 | **NA** | **0.88**<br>0.72-0.95 |
| **5** | **0.79**<br>0.52-0.91 | **0.85**<br>0.65-0.94 | **0.87**<br>0.70-0.94 | **0.88**<br>0.72-0.95 | **NA** |

ICC (2, 1) = SPSS, Model 2, repeated measures ANOVA, single measures; NA = not applicable; top bolded number = ICC (2, 1); bottom number = 95% Confidence Interval; all p-values < 0.001