

Supplemental Table 1. Demographics of enrolled patients.

	Control (200 mg/day)	DHA (1000 mg/day)	<i>p value</i>
Maternal			
Enrollment, <i>n</i>	16	11	
Lost prior to delivery, <i>n</i>	2	1	
Race, <i>n</i>			
Caucasian	10	6	0.910
African American	3	4	
Other/not disclosed	1	0	
Maternal BMI at enrollment	33.54 (30.63)	29.64 (28.62)	0.254
Mode of birth, <i>n</i>			
Cesarean	8	8	0.252
Vaginal	6	2	
Infant			
Enrollment, <i>n</i>	16	14	
GA at birth, <i>weeks</i>	25.88 (25.80)	26.33 (26.70)	0.331
Withdrew/Dropped	5	1	
Deceased, <i>n</i>	2	3	
	(1 completed before death)	(1 completed before death)	
Completed 4 weeks	8	8	
Infant sex, <i>n= males</i>	5	4	
Infant morbidities, <i>n</i>			
RDS/CLD	7	8	0.696
IVH	2	1	0.360
Sepsis	1	1	0.515
PDA	1	3	0.633
NEC	2	0	0.408

Data were analyzed by Mann-Whitney U-test or Kruskal-Wallace. No differences between groups were indicated. RDS/CLD, respiratory distress syndrome/chronic lung disease; IVH, intraventricular hemorrhage; PDA, patent ductus arteriosus; NEC, necrotizing enterocolitis.