etable 1. Individual patient phenotypes and investigations

r (Clinical syndrome	Autoimmune screen	Infection screen	CSF results	CoV-2 ab titres		recorded EDSS
-		2400 (POS)	1600 (POS)	STM	ANA, ENA, anti-neuronal ab negative, dsDNA and AQP4 NEG		WCC 0, protein 335, OCB absent, PCR NEG	N/A	257	1.5
g		3200 (POS)	1600 (POS)	LETM & STM, possible BS (ataxia)	N/A	HIV, Hep B and C & syphilis NEG	WCC 34, protein 660, OCB absent, PCR NEG	N/A	404	N/A
-			300 (LOW POS)		ANA, AQP4, phospholipid ab, ENA, anti-ribosomal P, Ro, SSA, SSB, centromere B, SM, anti-Sm, RNP-68, RNP-A, Scl-70, Jo NEG	Hep B&C, HIV, Quantiferon NEG	WCC 78, OCB absent	Anti-S 551.5 AU/ml. Anti-N NEG	237	3.0
:			200 (LOW POS)			Syphilis, Hep B&C, HIV NEG.	WCC 56, protein 1306, OCB absent, PCR NEG	N/A	356	2.0
-	-0	1600 (POS)	2400 (POS)	BS	RF, ANA and ANCA, complement, SPE, ACE, dsDNA and phospholipid ab NEG	Hep B&C, HIV NEG	WCC 30, protein 528, OCB matched, PCR NEG	N/A	222	3.0
	L8	1600	1200 (POS)	LETM	RF, ANA, phospholipid	Borrelia	WCC 150, protein	N/A	384	1.0

Ab	Antibody					
ANA	Anti-nuclear antibodies					
ANCA	Anti-neutrophil cytoplasmic antibo					
ASOT	Anti-streptolysin-O titre					
BS	Brainstem					
C3/4	Complement factor 3 & 4					
ССР	Anti-cyclic citrullinated peptide					
EBV	Epstein-Bar virus					
ENA	Extractable nuclear antigen antiboo					
GBS	Guillain-Barre syndrome					
Нер	Hepatitis					
HTLV	Human T-lymphotropic virus					
LETM	Longitudinally extensive transverse					
N/A	No test performed/ no results avail					
NEG	Negative					
OCB	Oligoclonal bands					
ONb	Optic neuritis (bilateral)					
Onu	Optic neuritis (unilateral)					
PCR	Polymerase chain reaction. Minimu					
virus, Enterovirus, Herpes simplex virus						
POS	Positive					
RF	Rheumatoid factor					
STM	Short segment myelitis					
Тохо	Toxoplasma					
WCC	White cell count					
CSF protein expressed as mg/L						
CSF WCC expressed as No. x 10 ⁶ /L						
•						

e table 2. Seronegative individuals' clinical syndromes according to vaccine type

BNT162b2	ChAdOx1S		
STM and asymptomatic periventricular lesions. OCB+	LETM and bilateral cerebral lesions		
Pure LETM	Pure LETM		
Bilateral sequential ON	LETM and STM with facial palsy		
Unilateral ON	ON and facial palsy		
STM and unilateral ON. OCB+	ON, LETM, brainstem & encephalopathy; possible GBS		
	STM, confirmed GBS		

GBS- Guillain-Barre syndrome

LETM- longitudinally extensive transverse myelitis

OCB- oligoclonal bands

ON- optic neuritis

STM- short segment myelitis

etable 3. Investigation and treatment of seronegative patients according to vaccine type

		BNT162b2 (n=5)	ChAdOx1S (n=6)
Immune screen	ANA	1/3	0/4
	ENAs	0/2	0/1
	ANCA	0/1	0/3
	Phospholipid antibodies	0/1	0/1
Infection screen	HIV	0/3	0/3
	Syphilis	0/2	0/3
	Lyme	0/1	0/3*
CSF parameters	Median CSF white cell count (x10 ⁶ /L)	6	4.5
	Median CSF protein (g/L)	282.5	1130
	CSF oligoclonal bands	2/3	0/5
Treatment	Oral steroid only	2/5	1/6**
	IVMP only	2/5	0/6
	IVIg only	0/5	1/6***
	IVMP + PLEX	1/5	1/6
	IVMP + PLEX + IVIg	0/5	2/5
	IVMP + PLEX + IVIg + CYCL	0/5	1/6

ANA- anti-nuclear antibodies

ANCA- anti-neutrophil cytoplasmic antibodies

CYCL- cyclophosphamide

ENA- extractable nuclear antigen antibodies

IVIG- intravenous immunoglobulins

IVMP- intravenous methylprednisolone


PLEX- plasma exchange


^{*2} patients' screening enzyme immunoassay positive, subsequently confirmed negative

^{**}Presumed Bell's palsy

^{***}GBS treatment

eFigure 1. Effect of season on attack onset pre- and post-introduction of vaccine





Ratchet circular scan test for seasonality pre-pandemic and after vaccine introduction March 2021-February 2022. 3 month peak (March-May 2021), 57.1% events, p< 0.025.