

Supplementary Table 2: Risk factors of genital *Chlamydia trachomatis* (CT) infection among male and female sexually-experienced participants Comarca Ngäbe-Bugle, Panama, 2018

	Yes n/N (%)	p-value	OR (95%CI)	p-value	AOR (95%CI)	p-value
Participant sex		0.03	1.77 (1.06-2.94)	0.03	2.02 (1.20-3.41)	<0.01
Female	39/224 (17.4)					
Male	31/291 (10.7)					
Age		0.37	1.23 (1.00-1.51)	0.05	1.30 (1.05-1.60)	0.02
14-15 years	6/71 (8.5)					
16 years	7/65 (10.8)					
17 years	13/98 (13.3)					
18 years	24/173 (13.9)					
19 years	20/108 (18.5)				3.0 (1.12-8.07)	
Comarcal Region		0.43	1.20 (0.91-1.59)	0.19	1.21 (0.91-1.60)	0.2
Nidrini (Chiriquí)	29/248 (11.7)					
Kädriri (Veraguas)	13/93 (14.0)					
Ño Kribo (Bocas del Toro)	28/174 (16.1)					
Forced sex¹		0.47				
No	46/355 (13.0)					
Yes	20/129 (15.5)		0.72 (0.24-2.11)	0.55		
Transactional sex²		0.86				
No	55/410 (13.4)					
Yes	10/79 (12.7)					
Number of partners in lifetime		0.24	1.20 (0.85-1.69)	0.31		
1	17/152 (11.2)					
2	13/66 (19.7)					
3 or more	17/112 (15.2)					
Same-sex sex partner		0.54	1.06 (0.78-1.44)	0.71		
No	52/350 (14.9)					
Yes	4/36 (11.1)					

AOR= adjusted odds ratio; CI = confidence interval; OR = odds ratio

¹Has been physically forced to engage in sex

²Has been offered something (money, food, housing, a better grade) in exchange for sex.