

Appendix 1

A: Low back pain trials

- Alcoff, J., Jones, E., Rust, P., & Newman, R. (1982). Controlled trial of imipramine for chronic low back pain. *J Fam Pract*, 14(5), 841-846.
- Amlie, E., Weber, H., & Holme, I. (1987). Treatment of acute low-back pain with piroxicam: results of a double-blind placebo-controlled trial. *Spine (Phila Pa 1976)*, 12(5), 473-476.
- Arbus, L., Fajadet, B., Aubert, D., Morre, M., & Goldberger, E. (1990). Activity of Tetraxepam (Myolastan) in Low-Back-Pain-a Double-Blind Trial V Placebo. *Clinical Trials Journal*, 27(4), 258-267.
- Atkinson, J. H., Slater, M. A., Capparelli, E. V., Wallace, M. S., Zisook, S., Abramson, I., . . . Garfin, S. R. (2007). Efficacy of noradrenergic and serotonergic antidepressants in chronic back pain: a preliminary concentration-controlled trial. *Journal of clinical psychopharmacology*, 27(2), 135-142.
- Atkinson, J. H., Slater, M. A., Wahlgren, D. R., Williams, R. A., Zisook, S., Pruitt, S. D., . . . Abramson, I. (1999). Effects of noradrenergic and serotonergic antidepressants on chronic low back pain intensity. *Pain*, 83(2), 137-145.
- Atkinson, J. H., Slater, M. A., Williams, R. A., Zisook, S., Patterson, T. L., Grant, I., . . . Garfin, S. R. (1998). A placebo-controlled randomized clinical trial of nortriptyline for chronic low back pain. *Pain*, 76(3), 287-296.
- Babej-Dolle, R., Freytag, S., Eckmeyer, J., Zerle, G., Schinzel, S., Schmeider, G., & Stankov, G. (1994). Parenteral dipyron versus diclofenac and placebo in patients with acute lumbago or sciatic pain: randomized observer-blind multicenter study. *Int J Clin Pharmacol Ther*, 32(4), 204-209.
- Baratta, R. R. (1976). A double-blind comparative study of carisoprodol, propoxyphene, and placebo in the management of low back syndrome. *Curr Ther Res Clin Exp*, 20(3), 233-240.
- Baron, R., Freynhagen, R., Tolle, T. R., Cloutier, C., Leon, T., Murphy, T. K., . . . Investigators, A. (2010). The efficacy and safety of pregabalin in the treatment of neuropathic pain associated with chronic lumbosacral radiculopathy. *Pain*, 150(3), 420-427. doi:10.1016/j.pain.2010.04.013
- Basmajian, J. (1978). Cyclobenzaprine hydrochloride effect on skeletal muscle spasm in the lumbar region and neck: two double-blind controlled clinical and laboratory studies. *Archives of physical medicine and rehabilitation*, 59(2), 58-63. [\(2 trials\)](#)
- Basmajian, J. V. (1989). Acute back pain and spasm. A controlled multicenter trial of combined analgesic and antispasm agents. *Spine*, 14(4), 438-439.
- Beaulieu, A. D., Peloso, P., Bensen, W., Clark, A. J., Watson, C. P. N., Gardner-Nix, J., . . . Harsanyi, Z. (2007). A randomized, double-blind, 8-week crossover study of once-daily controlled-release tramadol versus immediate-release tramadol taken as needed for chronic noncancer pain. *Clinical therapeutics*, 29(1), 49-60.
- Berry, H., Bloom, B., Hamilton, E. B., & Swinson, D. R. (1982). Naproxen sodium, diflunisal, and placebo in the treatment of chronic back pain. *Ann Rheum Dis*, 41(2), 129-132.
- Berry, H., & Hutchinson, D. (1988). A multicentre placebo-controlled study in general practice to evaluate the efficacy and safety of tizanidine in acute low-back pain. *Journal of international medical research*, 16(2), 75-82.
- Bianchi, M. (1978). Evaluation of cyclobenzaprine for skeletal muscle spasm of local origin. *Postgrad Med Comm*, 25-29.
- Birbara, C. A., Puopolo, A. D., Munoz, D. R., Sheldon, E. A., Mangione, A., Bohidar, N. R., . . . Group, E. P. S. (2003). Treatment of chronic low back pain with etoricoxib, a new cyclo-oxygenase-2 selective inhibitor: improvement in pain and disability—a randomized, placebo-controlled, 3-month trial. *The Journal of Pain*, 4(6), 307-315.

- Brötz, D., Maschke, E., Burkard, S., Engel, C., Mänz, C., Ernemann, U., . . . Weller, M. (2010). Is there a role for benzodiazepines in the management of lumbar disc prolapse with acute sciatica? *Pain*, 149(3), 470-475.
- Buynak, R., Shapiro, D. Y., Okamoto, A., Hove, I. V., Rauschkolb, C., Steup, A., . . . Etropolski, M. (2010). Efficacy and safety of tapentadol extended release for the management of chronic low back pain: results of a prospective, randomized, double-blind, placebo-and active-controlled Phase III study. *Expert opinion on pharmacotherapy*, 11(11), 1787-1804.
- Casale, R. (1988). Acute Low Back Pain. *The Clinical journal of pain*, 4(2), 81-88.
- Chu, L. F., D'Arcy, N., Brady, C., Zamora, A. K., Young, C. A., Kim, J. E., . . . Clark, J. D. (2012). Analgesic tolerance without demonstrable opioid-induced hyperalgesia: a double-blinded, randomized, placebo-controlled trial of sustained-release morphine for treatment of chronic nonradicular low-back pain. *Pain®*, 153(8), 1583-1592.
- Cloutier, C., Taliano, J., O'Mahony, W., Csanadi, M., Cohen, G., Sutton, I., . . . Robinson, L. (2013). Controlled-release oxycodone and naloxone in the treatment of chronic low back pain: a placebo-controlled, randomized study. *Pain Research and Management*, 18(2), 75-82.
- Coats, T. L., Borenstein, D. G., Nangia, N. K., & Brown, M. T. (2004). Effects of valdecoxib in the treatment of chronic low back pain: results of a randomized, placebo-controlled trial. *Clinical therapeutics*, 26(8), 1249-1260.
- Dapas, F., Hartman, S. F., Martinez, L., Northrup, B., Nussdorf, R. T., Silberman, H. M., & Gross, H. (1985). Baclofen for the Treatment of Acute Low-Back Syndrome: A Double-Blind Comparison with Placebo. *Spine*, 10(4), 345-349.
- Dickens, C., Jayson, M., Sutton, C., & Creed, F. (2000). The relationship between pain and depression in a trial using paroxetine in sufferers of chronic low back pain. *Psychosomatics*, 41(6), 490-499.
- Dreiser, R.-L., Le Parc, J., Velicitat, P., & Lleu, P. (2001). Oral meloxicam is effective in acute sciatica: two randomised, double-blind trials versus placebo or diclofenac. *Inflammation research*, 50(1), 17-23.
- Dreiser, R., Marty, M., Ionescu, E., Gold, M., & Liu, J. (2003). Relief of acute low back pain with diclofenac-K 12.5 mg tablets: a flexible dose, ibuprofen 200 mg and placebo-controlled clinical trial. *Int J Clin Pharmacol Ther*, 41(9), 375-385.
- Eskin, B., Shih, R. D., Fiesseler, F. W., Walsh, B. W., Allegra, J. R., Silverman, M. E., . . . Troncoso, A. (2014). Prednisone for emergency department low back pain: a randomized controlled trial. *The Journal of emergency medicine*, 47(1), 65-70.
- Finckh, A., Zufferey, P., Schurch, M.-A., Balagué, F., Waldburger, M., & So, A. K. (2006). Short-term efficacy of intravenous pulse glucocorticoids in acute discogenic sciatica. A randomized controlled trial. *Spine*, 31(4), 377-381.
- Friedman, B. W., Esses, D., Solorzano, C., Choi, H. K., Cole, M., Davitt, M., . . . Gallagher, E. (2008). A randomized placebo-controlled trial of single-dose IM corticosteroid for radicular low back pain. *Spine*, 33(18), E624.
- Friedman, B. W., Holden, L., Esses, D., Bijur, P. E., Choi, H. K., Solorzano, C., . . . Gallagher, E. J. (2006). Parenteral corticosteroids for Emergency Department patients with non-radicular low back pain. *The Journal of emergency medicine*, 31(4), 365-370.
- Goldie, I. (1968). A clinical trial with indomethacin (indomee®) in low back pain and sciatica. *Acta orthopaedica Scandinavica*, 39(1-3), 117-128.
- Goodkin, K., Gullion, C. M., & Agras, W. S. (1990). A Randomized, Double-Blind, Placebo-Controlled: Trial of Trazodone Hydrochloride in Chronic Low Back: Pain Syndrome. *Journal of clinical psychopharmacology*, 10(4), 269-278.
- Gordon, A., Callaghan, D., Spink, D., Cloutier, C., Dzungowski, P., O'Mahony, W., . . . Cohen, G. (2010). Buprenorphine transdermal system in adults with chronic low back pain: a randomized, double-blind, placebo-controlled crossover study, followed by an open-label extension phase. *Clinical therapeutics*, 32(5), 844-860.

- Haimovic, I. C., & Beresford, H. R. (1986). Dexamethasone is not superior to placebo for treating lumbosacral radicular pain. *Neurology*, 36(12), 1593-1593.
- Hale, M., Khan, A., Kutch, M., & Li, S. (2010). Once-daily OROS hydromorphone ER compared with placebo in opioid-tolerant patients with chronic low back pain. *Current medical research and opinion*, 26(6), 1505-1518.
- Hale, M. E., Ahdieh, H., Ma, T., Rauck, R., & Group, O. E. S. (2007). Efficacy and safety of OPANA ER (oxymorphone extended release) for relief of moderate to severe chronic low back pain in opioid-experienced patients: a 12-week, randomized, double-blind, placebo-controlled study. *The Journal of Pain*, 8(2), 175-184.
- Hale, M. E., Dvergsten, C., & Gimbel, J. (2005). Efficacy and safety of oxymorphone extended release in chronic low back pain: results of a randomized, double-blind, placebo-and active-controlled phase III study. *The Journal of Pain*, 6(1), 21-28.
- Hancock, M. J., Maher, C. G., Latimer, J., McLachlan, A. J., Cooper, C. W., Day, R. O., . . . McAuley, J. H. (2007). Assessment of diclofenac or spinal manipulative therapy, or both, in addition to recommended first-line treatment for acute low back pain: a randomised controlled trial. *The Lancet*, 370(9599), 1638-1643.
- Hedeboe, J., Buhl, M., & Ramsing, P. (1982). Effects of using dexamethasone and placebo in the treatment of prolapsed lumbar disc. *Acta neurologica Scandinavica*, 65(1), 6-10.
- Herrmann, W., & Geertsens, M. (2009). Efficacy and safety of lornoxicam compared with placebo and diclofenac in acute sciatica/lumbo-sciatica: an analysis from a randomised, double-blind, multicentre, parallel-group study. *International journal of clinical practice*, 63(11), 1613-1621.
- Hindle III, T. H. (1972). Comparison of carisoprodol, butabarbital, and placebo in treatment of the low back syndrome. *California medicine*, 117(2), 7.
- Hingorani, K. (1966). Diazepam in backache. A double-blind controlled trial. *Ann Phys Med*, 8(8), 303-306.
- Holve, R. L., & Barkan, H. (2008). Oral steroids in initial treatment of acute sciatica. *The Journal of the American Board of Family Medicine*, 21(5), 469-474.
- Jacobs, J., & Grayson, M. (1968). Trial of an anti-inflammatory agent (indomethacin) in low back pain with and without radicular involvement. *Br Med J*, 3(5611), 158-160.
- Jenkins, D., Ebbutt, A., & Evans, C. (1976). Tofranil in the treatment of low back pain. *The Journal of international medical research*, 4(2 Suppl), 28-40.
- Katz, J., Pennella-Vaughan, J., Hetzel, R. D., Kanazi, G. E., & Dworkin, R. H. (2005). A randomized, placebo-controlled trial of bupropion sustained release in chronic low back pain. *The Journal of Pain*, 6(10), 656-661.
- Katz, N., Ju, W. D., Krupa, D. A., Sperling, R. S., Rodgers, D. B., Gertz, B. J., . . . Nabizadeh, S. (2003). Efficacy and safety of rofecoxib in patients with chronic low back pain: results from two 4-week, randomized, placebo-controlled, parallel-group, double-blind trials. *Spine*, 28(9), 851-858.
- Katz, N., Rauck, R., Ahdieh, H., Ma, T., van der Hoop, R. G., Kerwin, R., & Podolsky, G. (2007). A 12-week, randomized, placebo-controlled trial assessing the safety and efficacy of oxymorphone extended release for opioid-naïve patients with chronic low back pain. *Current medical research and opinion*, 23(1), 117-128.
- Khoromi, S., Cui, L., Nackers, L., & Max, M. B. (2007). Morphine, nortriptyline and their combination vs. placebo in patients with chronic lumbar root pain. *Pain*, 130(1), 66-75.
- Khoromi, S., Patsalides, A., Parada, S., Salehi, V., Meegan, J. M., & Max, M. B. (2005). Topiramate in chronic lumbar radicular pain. *The Journal of Pain*, 6(12), 829-836.
- Lacey, P., Dodd, G., & Shannon, D. (1984). A double blind, placebo controlled study of piroxicam in the management of acute musculoskeletal disorders. *European journal of rheumatology and inflammation*, 7(3), 95-104.
- Lee, J. H., Lee, C.-S., & Group, U. E. S. (2013). A randomized, double-blind, placebo-controlled, parallel-group study to evaluate the efficacy and safety of the extended-release tramadol

- hydrochloride/acetaminophen fixed-dose combination tablet for the treatment of chronic low back pain. *Clinical therapeutics*, 35(11), 1830-1840.
- Lepisto, P. (1979). A comparative trial of ds 103-283 and placebo in the treatment of acute skeletal muscle spasms due to disorders of the back. *Current therapeutic research -clinical and experimental*, 26(4), 454-459.
- Markman, J. D., Frazer, M. E., Rast, S. A., McDermott, M. P., Gewandter, J. S., Chowdhry, A. K., . . . Dworkin, R. H. (2015). Double-blind, randomized, controlled, crossover trial of pregabalin for neurogenic claudication. *Neurology*, 84(3), 265-272.
- McCleane, G. J. (2001). Does gabapentin have an analgesic effect on background, movement and referred pain? A randomised, double-blind, placebo controlled study. *The Pain Clinic*, 13(2), 103-107.
- Moll, W. (1973). [Therapy of acute lumbovertebral syndromes through optimal muscle relaxation using diazepam. Results of a double-blind study on 68 cases]. *Med Welt*, 24(45), 1747-1751.
- Muehlbacher, M., Nickel, M. K., Kettler, C., Tritt, K., Lahmann, C., Leiberich, P. K., . . . Rother, W. K. (2006). Topiramate in treatment of patients with chronic low back pain: a randomized, double-blind, placebo-controlled study. *The Clinical journal of pain*, 22(6), 526-531.
- Nalamachu, S., Hale, M., & Khan, A. (2014). Hydromorphone extended release for neuropathic and non-neuropathic/nociceptive chronic low back pain: a post hoc analysis of data from a randomized, multicenter, double-blind, placebo-controlled clinical trial. *J Opioid Manag*, 10(5), 311-322. doi:10.5055/jom.2014.0221
- Peloso, P. M., Fortin, L., Beaulieu, A., Kamin, M., Rosenthal, N., & Group, P. T.-C.-S. (2004). Analgesic efficacy and safety of tramadol/acetaminophen combination tablets (Ultracet) in treatment of chronic low back pain: a multicenter, outpatient, randomized, double blind, placebo controlled trial. *The Journal of rheumatology*, 31(12), 2454-2463.
- Pheasant, H., Bursk, A., Goldfarb, J., Azen, S. P., Weiss, J. N., & Borelli, L. (1983). Amitriptyline and chronic low-back pain: a randomized double-blind crossover study. *Spine*, 8(5), 552-557.
- Porsman, O., & Friis, H. (1979). Prolapsed lumbar disc treated with intramuscularly administered dexamethasonephosphate: a prospectively planned, double-blind. Controlled clinical trial in 52 patients. *Scandinavian journal of rheumatology*, 8(3), 142-144.
- Pratzel, H., Alken, R.-G., & Ramm, S. (1996). Efficacy and tolerance of repeated oral doses of tolperisone hydrochloride in the treatment of painful reflex muscle spasm: results of a prospective placebo-controlled double-blind trial. *Pain*, 67(2), 417-425.
- Preuper, H. R. S., Geertzen, J. H., van Wijhe, M., Boonstra, A. M., Molmans, B. H., Dijkstra, P. U., & Reneman, M. F. (2014). Do analgesics improve functioning in patients with chronic low back pain? An explorative triple-blinded RCT. *European Spine Journal*, 23(4), 800-806.
- Ralph, L., Look, M., Wheeler, W., & Sacks, H. (2008). Double-blind, placebo-controlled trial of carisoprodol 250-mg tablets in the treatment of acute lower-back spasm. *Current medical research and opinion*, 24(2), 551-558.
- Rauck, R. L., Nalamachu, S., Wild, J. E., Walker, G. S., Robinson, C. Y., Davis, C. S., & Farr, S. J. (2014). Single-Entity Hydrocodone Extended-Release Capsules in Opioid-Tolerant Subjects with Moderate-to-Severe Chronic Low Back Pain: A Randomized Double-Blind, Placebo-Controlled Study. *Pain Medicine*, 15(6), 975-985.
- Roth, S. H., Fleischmann, R. M., Burch, F. X., Dietz, F., Bockow, B., Rapoport, R. J., . . . Lacouture, P. G. (2000). Around-the-clock, controlled-release oxycodone therapy for osteoarthritis-related pain: placebo-controlled trial and long-term evaluation. *Archives of Internal Medicine*, 160(6), 853-860.
- Ruoff, G. E., Rosenthal, N., Jordan, D., Karim, R., & Kamin, M. (2003). Tramadol/acetaminophen combination tablets for the treatment of chronic lower back pain: a multicenter, randomized, double-blind, placebo-controlled outpatient study. *Clinical therapeutics*, 25(4), 1123-1141.
- Salzmann, E., Pforringer, W., Paal, G., & Gierend, M. (1992). Treatment of chronic low-back syndrome with tetrazepam in a placebo controlled double-blind trial. *J Drug Dev*, 4(4), 219-228.

- Schnitzer, T. J., Gray, W. L., Paster, R. Z., & Kamin, M. (2000). Efficacy of tramadol in treatment of chronic low back pain. *The Journal of rheumatology*, 27(3), 772-778.
- Skljarevski, V., Desai, D., Liu-Seifert, H., Zhang, Q., Chappell, A. S., Detke, M. J., . . . Backonja, M. (2010). Efficacy and safety of duloxetine in patients with chronic low back pain. *Spine*, 35(13), E578-E585.
- Skljarevski, V., Ossanna, M., Liu-Seifert, H., Zhang, Q., Chappell, A., Iyengar, S., . . . Backonja, M. (2009). A double-blind, randomized trial of duloxetine versus placebo in the management of chronic low back pain. *European journal of neurology*, 16(9), 1041-1048.
- Skljarevski, V., Zhang, S., Desai, D., Alaka, K. J., Palacios, S., Miazgowski, T., & Patrick, K. (2010). Duloxetine versus placebo in patients with chronic low back pain: a 12-week, fixed-dose, randomized, double-blind trial. *The Journal of Pain*, 11(12), 1282-1290.
- Steiner, D. J., Sitar, S., Wen, W., Sawyerr, G., Munera, C., Ripa, S. R., & Landau, C. (2011). Efficacy and safety of the seven-day buprenorphine transdermal system in opioid-naïve patients with moderate to severe chronic low back pain: an enriched, randomized, double-blind, placebo-controlled study. *Journal of pain and symptom management*, 42(6), 903-917.
- Szpalski, M., & Hayez, J. (1994). Objective functional assessment of the efficacy of tenoxicam in the treatment of acute low back pain. A double-blind placebo-controlled study. *Rheumatology*, 33(1), 74-78.
- Trèves, R., Montaine, d. I. R. P., Dumond, J., Bertin, P., Arnaud, M., & Desproges-Gotteron, R. (1990). Prospective study of the analgesic action of clomipramine versus placebo in refractory lumbosciatica (68 cases). *Revue du rhumatisme et des maladies osteo-articulaires*, 58(7), 549-552.
- Vorsanger, G., Xiang, J., Gana, T., Pascual, M., & Fleming, R. (2008). Extended-release tramadol (tramadol ER) in the treatment of chronic low back pain. *Journal of opioid management*, 4(2), 87-97.
- Weber, H., & Aasand, G. (1980). The effect of phenylbutazone on patients with acute lumbago-sciatica. A double blind trial. *J Oslo City Hosp*, 30(5), 69-72.
- Weber, H., Holme, I., & Amlie, E. (1993). The natural course of acute sciatica with nerve root symptoms in a double-blind placebo-controlled trial evaluating the effect of piroxicam. *Spine*, 18(11), 1433-1438.
- Webster, L. R., Butera, P. G., Moran, L. V., Wu, N., Burns, L. H., & Friedmann, N. (2006). Oxytrex minimizes physical dependence while providing effective analgesia: a randomized controlled trial in low back pain. *The Journal of Pain*, 7(12), 937-946.
- Williams, C. M., Maher, C. G., Latimer, J., McLachlan, A. J., Hancock, M. J., Day, R. O., & Lin, C.-W. C. (2014). Efficacy of paracetamol for acute low-back pain: a double-blind, randomised controlled trial. *The Lancet*, 384(9954), 1586-1596.
- Überall, M. A., Mueller-Schwefe, G. H., & Terhaag, B. (2012). Efficacy and safety of flupirtine modified release for the management of moderate to severe chronic low back pain: results of SUPREME, a prospective randomized, double-blind, placebo-and active-controlled parallel-group phase IV study. *Current medical research and opinion*, 28(10), 1617-1634.
- Yildirim, K., Şişecioğlu, M., Karatay, S., Erdal, A., Levent, A., Uğur, M., & Şenel, K. (2003). The effectiveness of gabapentin in patients with chronic radiculopathy. *The Pain Clinic*, 15(3), 213-218. doi:10.1163/156856903767650718

B: Neuropathic pain trials

- Agrawal, R., Choudhary, R., Sharma, P., Sharma, S., Beniwal, R., Kaswan, K., & Kochar, D. (2007). Glyceryl trinitrate spray in the management of painful diabetic neuropathy: a randomized

- double blind placebo controlled cross-over study. *Diabetes research and clinical practice*, 77(2), 161-167.
- Agrawal, R., Goswami, J., Jain, S., & Kochar, D. (2009). Management of diabetic neuropathy by sodium valproate and glyceryl trinitrate spray: a prospective double-blind randomized placebo-controlled study. *Diabetes research and clinical practice*, 83(3), 371-378.
- Apalla, Z., Sotiriou, E., Lallas, A., Lazaridou, E., & Ioannides, D. (2013). Botulinum toxin A in postherpetic neuralgia: a parallel, randomized, double-blind, single-dose, placebo-controlled trial. *The Clinical journal of pain*, 29(10), 857-864.
- Arbaiza, D., & Vidal, O. (2007). Tramadol in the treatment of neuropathic cancer pain. *Clinical drug investigation*, 27(1), 75-83.
- Arezzo, J. C., Rosenstock, J., LaMoreaux, L., & Pauer, L. (2008). Efficacy and safety of pregabalin 600 mg/d for treating painful diabetic peripheral neuropathy: a double-blind placebo-controlled trial. *BMC neurology*, 8(1), 33.
- Atli, A., & Dogra, S. (2005). Zonisamide in the treatment of painful diabetic neuropathy: a randomized, double-blind, placebo-controlled pilot study. *Pain Medicine*, 6(3), 225-234.
- Backonja, M., Beydoun, A., Edwards, K. R., Schwartz, S. L., Fonseca, V., Hes, M., . . . Group, G. D. N. S. (1998). Gabapentin for the symptomatic treatment of painful neuropathy in patients with diabetes mellitus: a randomized controlled trial. *Jama*, 280(21), 1831-1836.
- Backonja, M., Wallace, M. S., Blonsky, E. R., Cutler, B. J., Malan, P., Rauck, R., . . . Group, N.-C. S. (2008). NGX-4010, a high-concentration capsaicin patch, for the treatment of postherpetic neuralgia: a randomised, double-blind study. *The Lancet Neurology*, 7(12), 1106-1112.
- Backonja, M. M., Malan, T. P., Vanhove, G. F., & Tobias, J. K. (2010). NGX-4010, a High-Concentration Capsaicin Patch, for the Treatment of Postherpetic Neuralgia: A Randomized, Double-Blind, Controlled Study with an Open-Label Extension. *Pain Medicine*, 11(4), 600-608.
- Bernstein, J. E., Korman, N. J., Bickers, D. R., Dahl, M. V., & Millikan, L. E. (1989). Topical capsaicin treatment of chronic postherpetic neuralgia. *Journal of the American Academy of Dermatology*, 21(2), 265-270.
- Beydoun, A., Shaibani, A., Hopwood, M., & Wan, Y. (2006). Oxcarbazepine in painful diabetic neuropathy: results of a dose-ranging study. *Acta neurologica Scandinavica*, 113(6), 395-404.
- Bone, M., Critchley, P., & Buggy, D. J. (2002). Gabapentin in postamputation phantom limb pain: a randomized, double-blind, placebo-controlled, cross-over study. *Regional anesthesia and pain medicine*, 27(5), 481-486.
- Boureau, F., Legallier, P., & Kabir-Ahmadi, M. (2003). Tramadol in post-herpetic neuralgia: a randomized, double-blind, placebo-controlled trial. *Pain*, 104(1), 323-331.
- Breuer, B., Pappagallo, M., Knotkova, H., Guleyupoglu, N., Wallenstein, S., & Portenoy, R. K. (2007). A randomized, double-blind, placebo-controlled, two-period, crossover, pilot trial of lamotrigine in patients with central pain due to multiple sclerosis. *Clinical therapeutics*, 29(9), 2022-2030.
- Byas-Smith, M. G., Max, M. B., Muir, J., & Kingman, A. (1995). Transdermal clonidine compared to placebo in painful diabetic neuropathy using a two-stage 'enriched enrollment' design. *Pain*, 60(3), 267-274. (2 trials)
- Campbell, C. M., Kipnes, M. S., Stouch, B. C., Brady, K. L., Kelly, M., Schmidt, W. K., . . . Campbell, J. N. (2012). Randomized control trial of topical clonidine for treatment of painful diabetic neuropathy. *Pain*, 153(9), 1815-1823.
- Cardenas, D. D., Nieshoff, E. C., Suda, K., Goto, S.-i., Sanin, L., Kaneko, T., . . . Yang, R. (2013). A randomized trial of pregabalin in patients with neuropathic pain due to spinal cord injury. *Neurology*, 80(6), 533-539.
- Chabal, C., Jacobson, L., Mariano, A., Chaney, E., & Britell, C. W. (1992). The use of oral mexiletine for the treatment of pain after peripheral nerve injury. *Anesthesiology*, 76(4), 513-517.
- Chad, D. A., Aronin, N., Lundstrom, R., McKeon, P., Ross, D., Molitch, M., . . . Tarsy, D. (1990). Does capsaicin relieve the pain of diabetic neuropathy? *Pain*, 42(3), 387-388.

- Cheville, A. L., Sloan, J. A., Northfelt, D. W., Jillella, A. P., Wong, G. Y., Bearden III, J. D., . . . Christensen, B. J. (2009). Use of a lidocaine patch in the management of postsurgical neuropathic pain in patients with cancer: a phase III double-blind crossover study (N01CB). *Supportive care in cancer*, 17(4), 451-460.
- Chiou-Tan, F. Y., Tuel, S. M., Johnson, J. C., Priebe, M. M., Hirsh, D. D., & Strayer, J. R. (1996). Effect of mexiletine on spinal cord injury dysesthetic pain. *Am J Phys Med Rehabil*, 75(2), 84-87.
- Clifford, D. B., Simpson, D. M., Brown, S., Moyle, G., Brew, B. J., Conway, B., . . . Group, N.-C. S. (2012). A randomized, double-blind, controlled study of NGX-4010, a capsaicin 8% dermal patch, for the treatment of painful HIV-associated distal sensory polyneuropathy. *JAIDS Journal of Acquired Immune Deficiency Syndromes*, 59(2), 126-133.
- Demant, D. T., Lund, K., Vollert, J., Maier, C., Segerdahl, M., Finnerup, N. B., . . . Sindrup, S. H. (2014). The effect of oxcarbazepine in peripheral neuropathic pain depends on pain phenotype: a randomised, double-blind, placebo-controlled phenotype-stratified study. *Pain®*, 155(11), 2263-2273.
- Dogra, S., Beydoun, S., Mazzola, J., Hopwood, M., & Wan, Y. (2005). Oxcarbazepine in painful diabetic neuropathy: A randomized, placebo-controlled study. *European journal of pain*, 9(5), 543-543.
- Drewes, A., Andreasen, A., & Poulsen, L. (1994). Valproate for treatment of chronic central pain after spinal cord injury. A double-blind cross-over study. *Spinal Cord*, 32(8), 565-569.
- Dworkin, R., Corbin, A., Young, J., Sharma, U., LaMoreaux, L., Bockbrader, H., . . . Poole, R. (2003). Pregabalin for the treatment of postherpetic neuralgia A randomized, placebo-controlled trial. *Neurology*, 60(8), 1274-1283.
- Eisenberg, E., Kleiser, A., Dortort, A., Haim, T., & Yarnitsky, D. (1998). The NMDA (N-methyl-D-aspartate) receptor antagonist memantine in the treatment of postherpetic neuralgia: a double-blind, placebo-controlled study. *European journal of pain*, 2(4), 321-327.
- Eisenberg, E., Lurie, Y., Braker, C., Daoud, D., & Ishay, A. (2001). Lamotrigine reduces painful diabetic neuropathy A randomized, controlled study. *Neurology*, 57(3), 505-509.
- Ellison, N., Loprinzi, C. L., Kugler, J., Hatfield, A. K., Miser, A., Sloan, J. A., . . . Cascino, T. L. (1997). Phase III placebo-controlled trial of capsaicin cream in the management of surgical neuropathic pain in cancer patients. *Journal of Clinical Oncology*, 15(8), 2974-2980.
- Evans, S. R., Simpson, D. M., Kitch, D. W., King, A., Clifford, D. B., Cohen, B. A., . . . Group, A. C. T. (2007). A randomized trial evaluating Prosaptide™ for HIV-associated sensory neuropathies: Use of an electronic diary to record neuropathic pain. *PLoS One*, 2(7), e551.
- Falah, M., Madsen, C., Holbech, J., & Sindrup, S. (2012). A randomized, placebo-controlled trial of levetiracetam in central pain in multiple sclerosis. *European journal of pain*, 16(6), 860-869.
- Finnerup, N., Grydehøj, J., Bing, J., Johannesen, I. L., Biering-Sørensen, F., Sindrup, S., & Jensen, T. S. (2009). Levetiracetam in spinal cord injury pain: a randomized controlled trial. *Spinal Cord*, 47(12), 861.
- Finnerup, N. B., Sindrup, S. H., Bach, F. W., Johannesen, I. L., & Jensen, T. S. (2002). Lamotrigine in spinal cord injury pain: a randomized controlled trial. *Pain*, 96(3), 375-383.
- Freeman, R., Raskin, P., Hewitt, D. J., Vorsanger, G. J., Jordan, D. M., Xiang, J., . . . Group, C.-S. (2007). Randomized study of tramadol/acetaminophen versus placebo in painful diabetic peripheral neuropathy. *Current medical research and opinion*, 23(1), 147-161.
- Freyenhagen, R., Strojek, K., Griesing, T., Whalen, E., & Balkenohl, M. (2005). Efficacy of pregabalin in neuropathic pain evaluated in a 12-week, randomised, double-blind, multicentre, placebo-controlled trial of flexible-and fixed-dose regimens. *Pain*, 115(3), 254-263.
- Gao, Y., Ning, G., Jia, W.-p., Zhou, Z.-g., Xu, Z.-r., Liu, Z.-m., . . . Cheng, L.-l. (2010). Duloxetine versus placebo in the treatment of patients with diabetic neuropathic pain in China. *Chinese medical journal*, 123(22), 3184-3192.
- Gilron, I., Bailey, J. M., Tu, D., Holden, R. R., Weaver, D. F., & Houlden, R. L. (2005). Morphine, gabapentin, or their combination for neuropathic pain. *New England Journal of Medicine*, 352(13), 1324-1334.

- Gilron, I., Booher, S., Rowan, J., Smoller, B., & Max, M. (2000). A randomized, controlled trial of high-dose dextromethorphan in facial neuralgias. *Neurology*, 55(7), 964-971.
- Gimbel, J. S., Richards, P., & Portenoy, R. K. (2003). Controlled-release oxycodone for pain in diabetic neuropathy A randomized controlled trial. *Neurology*, 60(6), 927-934.
- Goldstein, D. J., Lu, Y., Detke, M. J., Lee, T. C., & Iyengar, S. (2005). Duloxetine vs. placebo in patients with painful diabetic neuropathy. *Pain*, 116(1), 109-118.
- Goldstein, D. J., Wang, O., Gitter, B. D., & Iyengar, S. (2001). Dose-response study of the analgesic effect of lanepitant in patients with painful diabetic neuropathy. *Clinical neuropharmacology*, 24(1), 16-22.
- Gordh, T. E., Stubhaug, A., Jensen, T. S., Arnèr, S., Biber, B., Boivie, J., . . . Kalso, E. (2008). Gabapentin in traumatic nerve injury pain: a randomized, double-blind, placebo-controlled, cross-over, multi-center study. *Pain*, 138(2), 255-266.
- Gorson, K. C., Schott, C., Herman, R., Ropper, A. H., & Rand, W. M. (1999). Gabapentin in the treatment of painful diabetic neuropathy: a placebo controlled, double blind, crossover trial. *Journal of Neurology, Neurosurgery & Psychiatry*, 66(2), 251-252.
- Graff-Radford, S. B., Shaw, L. R., & Naliboff, B. N. (2000). Amitriptyline and fluphenazine in the treatment of postherpetic neuralgia. *The Clinical journal of pain*, 16(3), 188-192.
- Grosskopf, J., Mazzola, J., Wan, Y., & Hopwood, M. (2006). A randomized, placebo-controlled study of oxcarbazepine in painful diabetic neuropathy. *Acta neurologica Scandinavica*, 114(3), 177-180.
- Group, C. S. (1992). Effect of treatment with capsaicin on daily activities of patients with painful diabetic neuropathy. *Diabetes Care*, 15(2), 159-165.
- Guan, Y., Ding, X., Cheng, Y., Fan, D., Tan, L., Wang, Y., . . . Pan, X. (2011). Efficacy of pregabalin for peripheral neuropathic pain: results of an 8-week, flexible-dose, double-blind, placebo-controlled study conducted in China. *Clinical therapeutics*, 33(2), 159-166.
- Guida, G., De Martino, M., De Fabiani, A., Cantieri, L., Alexandre, A., & Vassallo, G. (2010). PEA (Normast) in chronic neuropathic pain caused by compression lumbosciatalgia: a multicentric clinical trial. *DOLOR*, 25, 35-42.
- Hahn, K., Arendt, G., Braun, J., von Giesen, H.-J., Husstedt, I., Maschke, M., . . . Group, G. N.-A. W. (2004). A placebo-controlled trial of gabapentin for painful HIV-associated sensory neuropathies. *Journal of neurology*, 251(10), 1260-1266.
- Harati, Y., Gooch, C., Swenson, M., Edelman, S., Greene, D., Raskin, P., . . . Siu, C. (1998). Double-blind randomized trial of tramadol for the treatment of the pain of diabetic neuropathy. *Neurology*, 50(6), 1842-1846.
- Holbech, J. V., Otto, M., Bach, F. W., Jensen, T. S., & Sindrup, S. H. (2011). The anticonvulsant levetiracetam for the treatment of pain in polyneuropathy: A randomized, placebo-controlled, cross-over trial. *European journal of pain*, 15(6), 608-614.
- Irving, G., Jensen, M., Cramer, M., Wu, J., Chiang, Y.-K., Tark, M., & Wallace, M. (2009). Efficacy and tolerability of gastric-retentive gabapentin for the treatment of postherpetic neuralgia: results of a double-blind, randomized, placebo-controlled clinical trial. *The Clinical journal of pain*, 25(3), 185-192.
- Irving, G. A., Backonja, M. M., Duntzman, E., Blonsky, E. R., Vanhove, G. F., Lu, S.-P., & Tobias, J. (2011). A multicenter, randomized, double-blind, controlled study of NGX-4010, a high-concentration capsaicin patch, for the treatment of postherpetic neuralgia. *Pain Medicine*, 12(1), 99-109.
- Jungehulsing, G., Israel, H., Safar, N., Taskin, B., Nolte, C., Brunecker, P., . . . Villringer, A. (2013). Levetiracetam in patients with central neuropathic post-stroke pain—a randomized, double-blind, placebo-controlled trial. *European journal of neurology*, 20(2), 331-337.
- Kalliomäki, J., Attal, N., Jonzon, B., Bach, F. W., Huizar, K., Ratcliffe, S., . . . Bouhassira, D. (2013). A randomized, double-blind, placebo-controlled trial of a chemokine receptor 2 (CCR2) antagonist in posttraumatic neuralgia. *Pain®*, 154(5), 761-767.

- Kalliomäki, J., Jonzon, B., Huizar, K., O'Malley, M., Andersson, A., & Simpson, D. M. (2013). Evaluation of a novel chemokine receptor 2 (CCR2)-antagonist in painful diabetic polyneuropathy. *Scandinavian Journal of Pain*, 4(2), 77-83.
- Kalso, E., Tasmuth, T., & Neuvonen, P. J. (1996). Amitriptyline effectively relieves neuropathic pain following treatment of breast cancer. *Pain*, 64(2), 293-302.
- Kemper, C. A., Kent, G., Burton, S., & Deresinski, S. C. (1998). Mexiletine for HIV-infected patients with painful peripheral neuropathy: a double-blind, placebo-controlled, crossover treatment trial. *JAIDS Journal of Acquired Immune Deficiency Syndromes*, 19(4), 367-372.
- Kiebertz, K., Simpson, D., Yiannoutsos, C., Max, M., Hall, C., Ellis, R., . . . Dal Pan, G. (1998). A randomized trial of amitriptyline and mexiletine for painful neuropathy in HIV infection. *Neurology*, 51(6), 1682-1688.
- Kim, J. S., Bashford, G., Murphy, T. K., Martin, A., Dror, V., & Cheung, R. (2011). Safety and efficacy of pregabalin in patients with central post-stroke pain. *Pain*, 152(5), 1018-1023.
- Kishore-kumar, R., Max, M. B., Schafer, S. C., Gaughan, A. M., Smoller, B., Gracely, R. H., & Dubner, R. (1990). Desipramine relieves postherpetic neuralgia. *Clinical Pharmacology & Therapeutics*, 47(3), 305-312.
- Kochar, D., Garg, P., Bumb, R., Kochar, S., Mehta, R., Beniwal, R., & Rawat, N. (2005). Divalproex sodium in the management of post-herpetic neuralgia: a randomized double-blind placebo-controlled study. *Qjm*, 98(1), 29-34.
- Kochar, D., Jain, N., Agarwal, R., Srivastava, T., Agarwal, P., & Gupta, S. (2002). Sodium valproate in the management of painful neuropathy in type 2 diabetes—a randomized placebo controlled study. *Acta neurologica Scandinavica*, 106(5), 248-252.
- Kochar, D., Rawat, N., Agrawal, R., Vyas, A., Beniwal, R., Kochar, S., & Garg, P. (2004). Sodium valproate for painful diabetic neuropathy: a randomized double-blind placebo-controlled study. *Qjm*, 97(1), 33-38.
- Kulkantrakorn, K., Lorsuwansiri, C., & Meesawatsom, P. (2013). 0.025% Capsaicin Gel for the Treatment of Painful Diabetic Neuropathy: A Randomized, Double-Blind, Crossover, Placebo-Controlled Trial. *Pain Practice*, 13(6), 497-503.
- Langford, R., Mares, J., Novotna, A., Vachova, M., Novakova, I., Notcutt, W., & Ratcliffe, S. (2013). A double-blind, randomized, placebo-controlled, parallel-group study of THC/CBD oromucosal spray in combination with the existing treatment regimen, in the relief of central neuropathic pain in patients with multiple sclerosis. *Journal of neurology*, 260(4), 984-997.
- Leijon, G., & Boivie, J. (1989). Central post-stroke pain—a controlled trial of amitriptyline and carbamazepine. *Pain*, 36(1), 27-36.
- Lesser, H., Sharma, U., LaMoreaux, L., & Poole, R. (2004). Pregabalin relieves symptoms of painful diabetic neuropathy A randomized controlled trial. *Neurology*, 63(11), 2104-2110.
- Levendoglu, F., Ögün, C. Ö., Özerbil, Ö., Ögün, T. C., & Ugurlu, H. (2004). Gabapentin is a first line drug for the treatment of neuropathic pain in spinal cord injury. *Spine*, 29(7), 743-751.
- Low, P. A., Opfer-Gehrking, T. L., Dyck, P. J., Litchy, W. J., & O'brien, P. C. (1995). Double-blind, placebo-controlled study of the application of capsaicin cream in chronic distal painful polyneuropathy. *Pain*, 62(2), 163-168.
- Lynch, M. E., Clark, A. J., Sawynok, J., & Sullivan, M. J. (2005). Topical 2% Amitriptyline and 1% Ketamine in Neuropathic Pain SyndromesA Randomized, Double-blind, Placebo-controlled Trial. *Anesthesiology: The Journal of the American Society of Anesthesiologists*, 103(1), 140-146.
- Maier, C., Dertwinkel, R., Mansourian, N., Hosbach, I., Schwenkreis, P., Senne, I., . . . Tegenthoff, M. (2003). Efficacy of the NMDA-receptor antagonist memantine in patients with chronic phantom limb pain—results of a randomized double-blinded, placebo-controlled trial. *Pain*, 103(3), 277-283.
- Max, M., Culnane, M., Schafer, S., Gracely, R., Walther, D., Smoller, B., & Dubner, R. (1987). Amitriptyline relieves diabetic neuropathy pain in patients with normal or depressed mood. *Neurology*, 37(4), 589-589.

- Max, M., Schafer, S., Culnane, M., Smoller, B., Dubner, R., & Gracely, R. (1988). Amitriptyline, but not lorazepam, relieves postherpetic neuralgia. *Neurology*, 38(9), 1427-1427.
- Max, M. B., Kishore-Kumar, R., Schafer, S. C., Meister, B., Gracely, R. H., Smoller, B., & Dubner, R. (1991). Efficacy of desipramine in painful diabetic neuropathy: a placebo-controlled trial. *Pain*, 45(1), 3-9.
- Max, M. B., Lynch, S. A., Muir, J., Shoaf, S. E., Smoller, B., & Dubner, R. (1992). Effects of desipramine, amitriptyline, and fluoxetine on pain in diabetic neuropathy. *New England Journal of Medicine*, 326(19), 1250-1256.
- McArthur, J., Yiannoutsos, C., Simpson, D., Adornato, B., Singer, E., Hollander, H., . . . Tucker, T. (2000). A phase II trial of nerve growth factor for sensory neuropathy associated with HIV infection. *Neurology*, 54(5), 1080-1088.
- McCleane, G. (1999). 200 mg daily of lamotrigine has no analgesic effect in neuropathic pain: a randomised, double-blind, placebo controlled trial. *Pain*, 83(1), 105-107.
- McCleane, G. (2000). Topical application of doxepin hydrochloride, capsaicin and a combination of both produces analgesia in chronic human neuropathic pain: a randomized, double-blind, placebo-controlled study. *British journal of clinical pharmacology*, 49(6), 574-579.
- Mishra, S., Bhatnagar, S., Goyal, G. N., Rana, S. P. S., & Upadhy, S. P. (2012). A comparative efficacy of amitriptyline, gabapentin, and pregabalin in neuropathic cancer pain: a prospective randomized double-blind placebo-controlled study. *American Journal of Hospice and Palliative Medicine*®, 29(3), 177-182.
- Moon, D. E., Lee, D. I., Lee, S. C., Song, S. O., Yoon, D. M., Yoon, M. H., . . . Lee, P. B. (2010). Efficacy and tolerability of pregabalin using a flexible, optimized dose schedule in Korean patients with peripheral neuropathic pain: a 10-week, randomized, double-blind, placebo-controlled, multicenter study. *Clinical therapeutics*, 32(14), 2370-2385.
- Motilal, S., & Maharaj, R. G. (2013). Nutmeg extracts for painful diabetic neuropathy: a randomized, double-blind, controlled study. *The Journal of Alternative and Complementary Medicine*, 19(4), 347-352.
- Nelson, K. A., Park, K. M., Robinovitz, E., Tsigos, C., & Max, M. B. (1997). High-dose oral dextromethorphan versus placebo in painful diabetic neuropathy and postherpetic neuralgia. *Neurology*, 48(5), 1212-1218.
- Nikolajsen, L., Gottrup, H., Kristensen, A. G., & Jensen, T. S. (2000). Memantine (a N-methyl-D-aspartate receptor antagonist) in the treatment of neuropathic pain after amputation or surgery: a randomized, double-blinded, cross-over study. *Anesthesia & Analgesia*, 91(4), 960-966.
- Norrbrink, C., & Lundeborg, T. (2009). Tramadol in neuropathic pain after spinal cord injury: a randomized, double-blind, placebo-controlled trial. *The Clinical journal of pain*, 25(3), 177-184.
- Nurmikko, T. J., Serpell, M. G., Hoggart, B., Toomey, P. J., Morlion, B. J., & Haines, D. (2007). Sativex successfully treats neuropathic pain characterised by allodynia: a randomised, double-blind, placebo-controlled clinical trial. *Pain*®, 133(1), 210-220.
- Oskarsson, P., Ljunggren, J.-G., & Lins, P.-E. (1997). Efficacy and safety of mexiletine in the treatment of painful diabetic neuropathy. The Mexiletine Study Group. *Diabetes Care*, 20(10), 1594-1597.
- Ostenfeld, T., Krishen, A., Lai, R., Bullman, J., Baines, A., Green, J., . . . Kelly, M. (2013). Analgesic efficacy and safety of the novel p38 MAP kinase inhibitor, losmapimod, in patients with neuropathic pain following peripheral nerve injury: a double-blind, placebo-controlled study. *European journal of pain*, 17(6), 844-857.
- Otto, M., Bach, F. W., Jensen, T. S., Brøsen, K., & Sindrup, S. H. (2008). Escitalopram in painful polyneuropathy: a randomized, placebo-controlled, cross-over trial. *Pain*, 139(2), 275-283.
- Otto, M., Bach, F. W., Jensen, T. S., & Sindrup, S. H. (2004). Valproic acid has no effect on pain in polyneuropathy A randomized, controlled trial. *Neurology*, 62(2), 285-288.

- Panerai, A., Monza, G., Movilia, P., Bianchi, M., Francucci, B., & Tiengo, M. (1990). A randomized, within-patient, cross-over, placebo-controlled trial on the efficacy and tolerability of the tricyclic antidepressants chlorimipramine and nortriptyline in central pain. *Acta neurologica Scandinavica*, 82(1), 34-38.
- Pickering, G., Morel, V., Simen, E., Cardot, J.-M., Moustafa, F., Delage, N., . . . Dubray, C. (2011). Oral magnesium treatment in patients with neuropathic pain: a randomized clinical trial. *Magnesium research*, 24(2), 28-35.
- Raja, S., Haythornthwaite, J., Pappagallo, M., Clark, M., Trivison, T., Sabeen, S., . . . Max, M. (2002). Opioids versus antidepressants in postherpetic neuralgia A randomized, placebo-controlled trial. *Neurology*, 59(7), 1015-1021.
- Ranoux, D., Attal, N., Morain, F., & Bouhassira, D. (2008). Botulinum toxin type A induces direct analgesic effects in chronic neuropathic pain. *Annals of neurology*, 64(3), 274-283.
- Raskin, J., Pritchett, Y. L., Wang, F., D'souza, D. N., Waninger, A. L., Iyengar, S., & Wernicke, J. F. (2005). A double-blind, randomized multicenter trial comparing duloxetine with placebo in the management of diabetic peripheral neuropathic pain. *Pain Medicine*, 6(5), 346-356.
- Raskin, P., Donofrio, P., Rosenthal, N., Hewitt, D., Jordan, D., Xiang, J., . . . Group, C.-S. (2004). Topiramate vs placebo in painful diabetic neuropathy Analgesic and metabolic effects. *Neurology*, 63(5), 865-873.
- Rauck, R., Makumi, C. W., Schwartz, S., Graff, O., Meno-Tetang, G., Bell, C. F., . . . McClung, C. L. (2013). A randomized, controlled trial of gabapentin enacarbil in subjects with neuropathic pain associated with diabetic peripheral neuropathy. *Pain Practice*, 13(6), 485-496.
- Rauck, R. L., Shaibani, A., Biton, V., Simpson, J., & Koch, B. (2007). Lacosamide in painful diabetic peripheral neuropathy: a phase 2 double-blind placebo-controlled study. *The Clinical journal of pain*, 23(2), 150-158.
- Rice, A., Maton, S., & Group1UK, P. N. S. (2001). Gabapentin in postherpetic neuralgia: a randomised, double blind, placebo controlled study. *Pain*, 94(2), 215-224.
- Rice, A. S., Dworkin, R. H., McCarthy, T. D., Anand, P., Bountra, C., McCloud, P. I., . . . Desem, N. (2014). EMA401, an orally administered highly selective angiotensin II type 2 receptor antagonist, as a novel treatment for postherpetic neuralgia: a randomised, double-blind, placebo-controlled phase 2 clinical trial. *The Lancet*, 383(9929), 1637-1647.
- Richter, R. W., Portenoy, R., Sharma, U., Lamoreaux, L., Bockbrader, H., & Knapp, L. E. (2005). Relief of painful diabetic peripheral neuropathy with pregabalin: a randomized, placebo-controlled trial. *The Journal of Pain*, 6(4), 253-260.
- Rintala, D. H., Holmes, S. A., Courtade, D., Fiess, R. N., Tastard, L. V., & Loubser, P. G. (2007). Comparison of the effectiveness of amitriptyline and gabapentin on chronic neuropathic pain in persons with spinal cord injury. *Archives of physical medicine and rehabilitation*, 88(12), 1547-1560.
- Robinson, L. R., Czerniecki, J. M., Ehde, D. M., Edwards, W. T., Judish, D. A., Goldberg, M. L., . . . Jensen, M. P. (2004). Trial of amitriptyline for relief of pain in amputees: results of a randomized controlled study. *Archives of physical medicine and rehabilitation*, 85(1), 1-6.
- Rog, D. J., Nurmikko, T. J., Friede, T., & Young, C. A. (2005). Randomized, controlled trial of cannabis-based medicine in central pain in multiple sclerosis. *Neurology*, 65(6), 812-819.
- Rosenstock, J., Tuchman, M., LaMoreaux, L., & Sharma, U. (2004). Pregabalin for the treatment of painful diabetic peripheral neuropathy: a double-blind, placebo-controlled trial. *Pain*, 110(3), 628-638.
- Rowbotham, M., Harden, N., Stacey, B., Bernstein, P., Magnus-Miller, L., & Group, G. P. N. S. (1998). Gabapentin for the treatment of postherpetic neuralgia: a randomized controlled trial. *Jama*, 280(21), 1837-1842.
- Rowbotham, M. C., Arslanian, A., Nothaft, W., Duan, W. R., Best, A. E., Pritchett, Y., . . . Stacey, B. R. (2012). Efficacy and safety of the $\alpha 4 \beta 2$ neuronal nicotinic receptor agonist ABT-894 in patients with diabetic peripheral neuropathic pain. *Pain*, 153(4), 862-868. (2 trials)

- Rowbotham, M. C., Duan, W. R., Thomas, J., Nothaft, W., & Backonja, M.-M. (2009). A randomized, double-blind, placebo-controlled trial evaluating the efficacy and safety of ABT-594 in patients with diabetic peripheral neuropathic pain. *Pain*, 146(3), 245-252.
- Rowbotham, M. C., Goli, V., Kunz, N. R., & Lei, D. (2004). Venlafaxine extended release in the treatment of painful diabetic neuropathy: a double-blind, placebo-controlled study. *Pain*, 110(3), 697-706.
- Sabatowski, R., Gálvez, R., Cherry, D. A., Jacquot, F., Vincent, E., Maisonobe, P., . . . Group, -. S. (2004). Pregabalin reduces pain and improves sleep and mood disturbances in patients with post-herpetic neuralgia: results of a randomised, placebo-controlled clinical trial. *Pain*, 109(1), 26-35.
- Sandercock, D., Cramer, M., Biton, V., & Cowles, V. E. (2012). A gastroretentive gabapentin formulation for the treatment of painful diabetic peripheral neuropathy: efficacy and tolerability in a double-blind, randomized, controlled clinical trial. *Diabetes research and clinical practice*, 97(3), 438-445.
- Sang, C. N., Booher, S., Gilron, I., Parada, S., & Max, M. B. (2002). Dextromethorphan and Memantine in Painful Diabetic Neuropathy and Postherpetic Neuralgia Efficacy and Dose-Response Trials. *Anesthesiology: The Journal of the American Society of Anesthesiologists*, 96(5), 1053-1061.
- Sang, C. N., Sathyanarayana, R., Sweeney, M., & Investigators, D.-S. (2013). Gastroretentive gabapentin (G-GR) formulation reduces intensity of pain associated with postherpetic neuralgia (PHN). *The Clinical journal of pain*, 29(4), 281-288.
- Satoh, J., Yagihashi, S., Baba, M., Suzuki, M., Arakawa, A., Yoshiyama, T., & Shoji, S. (2011). Efficacy and safety of pregabalin for treating neuropathic pain associated with diabetic peripheral neuropathy: a 14 week, randomized, double-blind, placebo-controlled trial. *Diabetic Medicine*, 28(1), 109-116.
- Scheffler, N., Sheitel, P., & Lipton, M. (1991). Treatment of painful diabetic neuropathy with capsaicin 0.075%. *Journal of the American Podiatric Medical Association*, 81(6), 288-293.
- Selvarajah, D., Gandhi, R., Emery, C. J., & Tesfaye, S. (2010). Randomized placebo-controlled double-blind clinical trial of cannabis-based medicinal product (Sativex) in painful diabetic neuropathy. *Diabetes Care*, 33(1), 128-130.
- Semenchuk, M. R., Sherman, S., & Davis, B. (2001). Double-blind, randomized trial of bupropion SR for the treatment of neuropathic pain. *Neurology*, 57(9), 1583-1588.
- Serpell, M., & Group, N. P. S. (2002). Gabapentin in neuropathic pain syndromes: a randomised, double-blind, placebo-controlled trial. *Pain*, 99(3), 557-566.
- Shackelford, S., Rauck, R., Quessy, S., Blum, D., Hodge, R., & Philipson, R. (2009). A randomized, double-blind, placebo-controlled trial of a selective COX-2 inhibitor, GW406381, in patients with postherpetic neuralgia. *The Journal of Pain*, 10(6), 654-660.
- Shaibani, A., Fares, S., Selam, J.-L., Arslanian, A., Simpson, J., Sen, D., & Bongardt, S. (2009). Lacosamide in painful diabetic neuropathy: an 18-week double-blind placebo-controlled trial. *The Journal of Pain*, 10(8), 818-828.
- Shaibani, A. I., Pope, L. E., Thisted, R., & Hepner, A. (2012). Efficacy and safety of dextromethorphan/quinidine at two dosage levels for diabetic neuropathic pain: a double-blind, placebo-controlled, multicenter study. *Pain Medicine*, 13(2), 243-254.
- Shembalkar, P., Täubel, J., Abadias, M., Arezina, R., Hammond, K., & Anand, P. (2001). Cizolirtine citrate (E-4018) in the treatment of chronic neuropathic pain. *Current medical research and opinion*, 17(4), 262-266.
- Siddall, P. J., Cousins, M., Otte, A., Griesing, T., Chambers, R., & Murphy, T. (2006). Pregabalin in central neuropathic pain associated with spinal cord injury A placebo-controlled trial. *Neurology*, 67(10), 1792-1800.
- Simpson, D., Dorfman, D., Olney, R., McKinley, G., Dobkin, J., So, Y., . . . Friedman, B. (1996). Peptide T in the treatment of painful distal neuropathy associated with AIDS Results of a placebo-controlled trial. *Neurology*, 47(5), 1254-1259.

- Simpson, D., Olney, R., McArthur, J., Khan, A., Godbold, J., Ebel-Frommer, K., & Group, L. H. N. S. (2000). A placebo-controlled trial of lamotrigine for painful HIV-associated neuropathy. *Neurology*, 54(11), 2115-2119.
- Simpson, D. A. (2001). Gabapentin and venlafaxine for the treatment of painful diabetic neuropathy. *Journal of clinical neuromuscular disease*, 3(2), 53-62.
- Simpson, D. M., Brown, S., Tobias, J., & Group, N.-C. S. (2008). Controlled trial of high-concentration capsaicin patch for treatment of painful HIV neuropathy. *Neurology*, 70(24), 2305-2313.
- Simpson, D. M., McArthur, J., Olney, R., Clifford, D., So, Y., Ross, D., . . . Team, L. H. N. S. (2003). Lamotrigine for HIV-associated painful sensory neuropathies A placebo-controlled trial. *Neurology*, 60(9), 1508-1514.
- Simpson, D. M., Rice, A. S., Emir, B., Landen, J., Semel, D., Chew, M. L., & Sporn, J. (2014). A randomized, double-blind, placebo-controlled trial and open-label extension study to evaluate the efficacy and safety of pregabalin in the treatment of neuropathic pain associated with human immunodeficiency virus neuropathy. *Pain*, 155(10), 1943-1954.
- Simpson, D. M., Schifitto, G., Clifford, D., Murphy, T., Durso-De Cruz, E., Glue, P., . . . Freeman, R. (2010). Pregabalin for painful HIV neuropathy A randomized, double-blind, placebo-controlled trial. *Neurology*, 74(5), 413-420.
- Sindrup, S. H., Andersen, G., Madsen, C., Smith, T., Brøsen, K., & Jensen, T. S. (1999). Tramadol relieves pain and allodynia in polyneuropathy: a randomised, double-blind, controlled trial. *Pain*, 83(1), 85-90.
- Sindrup, S. H., Bach, F. W., Madsen, C., Gram, L., & Jensen, T. (2003). Venlafaxine versus imipramine in painful polyneuropathy A randomized, controlled trial. *Neurology*, 60(8), 1284-1289.
- Sindrup, S. H., Konner, R., Lehmann, R., Meier, T., Winkel, M., Ashworth, J., . . . Jensen, T. (2012). Randomized controlled trial of the combined monoaminergic and opioid investigational compound GRT9906 in painful polyneuropathy. *European journal of pain*, 16(6), 849-859.
- Sindrup, S. H., Madsen, C., Bach, F. W., Gram, L. F., & Jensen, T. S. (2001). St. John's wort has no effect on pain in polyneuropathy. *Pain*, 91(3), 361-365.
- Smith, D. G., Ehde, D. M., Hanley, M. A., & Campbell, K. M. (2005). Efficacy of gabapentin in treating chronic phantom limb and residual limb pain. *Journal of rehabilitation research and development*, 42(5), 645.
- Smith, E. M. L., Pang, H., Cirrincione, C., Fleishman, S., Paskett, E. D., Ahles, T., . . . Le-Lindqwister, N. (2013). Effect of duloxetine on pain, function, and quality of life among patients with chemotherapy-induced painful peripheral neuropathy: a randomized clinical trial. *Jama*, 309(13), 1359-1367.
- Smith, T., DiBernardo, A., Shi, Y., Todd, M. J., Brashear, H. R., & Ford, L. M. (2014). Efficacy and Safety of Carisbamate in Patients with Diabetic Neuropathy or Postherpetic Neuralgia: Results from 3 Randomized, Double-Blind Placebo-Controlled Trials. *Pain Practice*, 14(4), 332-342 (3 trials)
- Stacey, B. R., Barrett, J. A., Whalen, E., Phillips, K. F., & Rowbotham, M. C. (2008). Pregabalin for postherpetic neuralgia: placebo-controlled trial of fixed and flexible dosing regimens on allodynia and time to onset of pain relief. *The Journal of Pain*, 9(11), 1006-1017.
- Staughton, R., Chir, B., & Good, J. (1990). Double-blind, placebo-controlled clinical trial of a mixture of gangliosides ('Cronassial') in post-herpetic neuralgia. *Current medical research and opinion*, 12(3), 169-176.
- Stracke, H., Meyer, U. E., Schumacher, H. E., & Federlin, K. (1992). Mexiletine in the treatment of diabetic neuropathy. *Diabetes Care*, 15(11), 1550-1555.
- Surman, O. S., Flynn, T., Schooley, R. T., Baer, L., Parker, S., Hirsch, M. S., & Davis, L. G. (1990). A double-blind, placebo-controlled study of oral acyclovir in postherpetic neuralgia. *Psychosomatics*, 31(3), 287-292.
- Svendsen, K. B., Jensen, T. S., & Bach, F. W. (2004). Does the cannabinoid dronabinol reduce central pain in multiple sclerosis? Randomised double blind placebo controlled crossover trial. *Bmj*, 329(7460), 253.

- Tandan, R., Lewis, G. A., Krusinski, P. B., Badger, G. B., & Fries, T. J. (1992). Topical capsaicin in painful diabetic neuropathy: controlled study with long-term follow-up. *Diabetes Care*, 15(1), 8-14.
- Tasmuth, T., Härtel, B., & Kalso, E. (2002). Venlafaxine in neuropathic pain following treatment of breast cancer. *European journal of pain*, 6(1), 17-24.
- Thienel, U., Neto, W., Schwabe, S., & Vijapurkar, U. (2004). Topiramate in painful diabetic polyneuropathy: findings from three double-blind placebo-controlled trials. *Acta neurologica Scandinavica*, 110(4), 221-231.
- Tölle, T., Freynhagen, R., Versavel, M., Trostmann, U., & Young, J. P. (2008). Pregabalin for relief of neuropathic pain associated with diabetic neuropathy: a randomized, double-blind study. *European journal of pain*, 12(2), 203-213.
- Van Seventer, R., Bach, F. W., Toth, C. C., Serpell, M., Temple, J., Murphy, T. K., & Nimour, M. (2010). Pregabalin in the treatment of post-traumatic peripheral neuropathic pain: a randomized double-blind trial. *European journal of neurology*, 17(8), 1082-1089.
- van Seventer, R., Feister, H. A., Young Jr, J. P., Stoker, M., Versavel, M., & Rigaudy, L. (2006). Efficacy and tolerability of twice-daily pregabalin for treating pain and related sleep interference in postherpetic neuralgia: a 13-week, randomized trial. *Current medical research and opinion*, 22(2), 375-384.
- Vestergaard, K., Andersen, G., Gottrup, H., Kristensen, B., & Jensen, T. S. (2001). Lamotrigine for central poststroke pain a randomized controlled trial. *Neurology*, 56(2), 184-190.
- Vilholm, O. J., Cold, S., Rasmussen, L., & Sindrup, S. H. (2008). Effect of levetiracetam on the postmastectomy pain syndrome. *European journal of neurology*, 15(8), 851-857.
- Vinik, A. I., Shapiro, D. Y., Rauschkolb, C., Lange, B., Karcher, K., Pennett, D., & Etropolski, M. S. (2014). A randomized withdrawal, placebo-controlled study evaluating the efficacy and tolerability of tapentadol extended release in patients with chronic painful diabetic peripheral neuropathy. *Diabetes Care*, 37(8), 2302-2309.
- Vinik, A. I., Tuchman, M., Safirstein, B., Corder, C., Kirby, L., Wilks, K., . . . White, J. (2007). Lamotrigine for treatment of pain associated with diabetic neuropathy: results of two randomized, double-blind, placebo-controlled studies. *Pain*, 128(1), 169-179. (2 trials)
- Vranken, J., Dijkgraaf, M., Kruis, M., Van der Vegt, M., Hollmann, M., & Heesen, M. (2008). Pregabalin in patients with central neuropathic pain: a randomized, double-blind, placebo-controlled trial of a flexible-dose regimen. *Pain*, 136(1), 150-157.
- Vranken, J., Hollmann, M., Van Der Vegt, M., Kruis, M., Heesen, M., Vos, K., . . . Dijkgraaf, M. (2011). Duloxetine in patients with central neuropathic pain caused by spinal cord injury or stroke: a randomized, double-blind, placebo-controlled trial. *Pain®*, 152(2), 267-273.
- Vrethem, M., Boivie, J., Arnqvist, H., Holmgren, H., Lindström, T., & Thorell, L.-H. (1997). A comparison of amitriptyline and maprotiline in the treatment of painful polyneuropathy in diabetics and nondiabetics. *The Clinical journal of pain*, 13(4), 313-323.
- Wallace, M. S., Irving, G., & Cowles, V. E. (2010). Gabapentin extended-release tablets for the treatment of patients with postherpetic neuralgia. *Clinical drug investigation*, 30(11), 765-776.
- Watson, C. P., Evans, R. J., Reed, K., Merskey, H., Goldsmith, L., & Warsh, J. (1982). Amitriptyline versus placebo in postherpetic neuralgia. *Neurology*, 32(6), 671-671.
- Watson, C. P. N., & Babul, N. (1998). Efficacy of oxycodone in neuropathic pain A randomized trial in postherpetic neuralgia. *Neurology*, 50(6), 1837-1841.
- Watson, C. P. N., & Evans, R. J. (1992). The postmastectomy pain syndrome and topical capsaicin: a randomized trial. *Pain*, 51(3), 375-379.
- Watson, C. P. N., Moulin, D., Watt-Watson, J., Gordon, A., & Eisenhoffer, J. (2003). Controlled-release oxycodone relieves neuropathic pain: a randomized controlled trial in painful diabetic neuropathy. *Pain*, 105(1), 71-78.
- Watson, C. P. N., Tyler, K., Bickers, D., Millikan, L., Smith, S., & Coleman, E. (1993). A randomized vehicle-controlled trial of topical capsaicin in the treatment of postherpetic neuralgia. *Clinical therapeutics*, 15, 510-510.

- Webster, L. R., Malan, T. P., Tuchman, M. M., Mollen, M. D., Tobias, J. K., & Vanhove, G. F. (2010). A multicenter, randomized, double-blind, controlled dose finding study of NGX-4010, a high-concentration capsaicin patch, for the treatment of postherpetic neuralgia. *The Journal of Pain*, 11(10), 972-982.
- Webster, L. R., Tark, M., Rauck, R., Tobias, J. K., & Vanhove, G. F. (2010). Effect of duration of postherpetic neuralgia on efficacy analyses in a multicenter, randomized, controlled study of NGX-4010, an 8% capsaicin patch evaluated for the treatment of postherpetic neuralgia. *BMC neurology*, 10(1), 92.
- Wernicke, J., Pritchett, Y., D'souza, D., Waninger, A., Tran, P., Iyengar, S., & Raskin, J. (2006). A randomized controlled trial of duloxetine in diabetic peripheral neuropathic pain. *Neurology*, 67(8), 1411-1420.
- Windebank, A. J., Sorenson, E. J., Civil, R., & O'brien, P. C. (2004). Role of insulin-like growth factor-I in the treatment of painful small fiber predominant neuropathy. *Journal of the Peripheral Nervous System*, 9(3), 183-189.
- Wright, J. M., Oki, J. C., & Graves III, L. (1997). Mexiletine in the symptomatic treatment of diabetic peripheral neuropathy. *Annals of Pharmacotherapy*, 31(1), 29-34.
- Wu, C.-J., Lian, Y.-J., Zheng, Y.-K., Zhang, H.-F., Chen, Y., Xie, N.-C., & Wang, L.-J. (2012). Botulinum toxin type A for the treatment of trigeminal neuralgia: results from a randomized, double-blind, placebo-controlled trial. *Cephalalgia*, 32(6), 443-450.
- Wu, C. L., Agarwal, S., Tella, P. K., Klick, B., Clark, M. R., Haythornthwaite, J. A., . . . Raja, S. N. (2008). Morphine versus Mexiletine for Treatment of Postamputation Pain A Randomized, Placebo-controlled, Crossover Trial. *Anesthesiology: The Journal of the American Society of Anesthesiologists*, 109(2), 289-296.
- Wymer, J. P., Simpson, J., Sen, D., Bongardt, S., & Group, L. S. S. (2009). Efficacy and safety of lacosamide in diabetic neuropathic pain: an 18-week double-blind placebo-controlled trial of fixed-dose regimens. *The Clinical journal of pain*, 25(5), 376-385.
- Xiao, L., Mackey, S., Hui, H., Xong, D., Zhang, Q., & Zhang, D. (2010). Subcutaneous injection of botulinum toxin a is beneficial in postherpetic neuralgia. *Pain Medicine*, 11(12), 1827-1833.
- Yasuda, H., Hotta, N., Nakao, K., Kasuga, M., Kashiwagi, A., & Kawamori, R. (2011). Superiority of duloxetine to placebo in improving diabetic neuropathic pain: results of a randomized controlled trial in Japan. *Journal of diabetes investigation*, 2(2), 132-139.
- Yuan, R.-Y., Sheu, J.-J., Yu, J. M., Chen, W. T., Tseng, I., Chang, H. H., & Hu, C.-J. (2009). Botulinum toxin for diabetic neuropathic pain A randomized double-blind crossover trial. *Neurology*, 72(17), 1473-1478.
- Yucel, A., Ozyalcin, S., Talu, G. K., Kiziltan, E., Yucel, B., Andersen, O. K., . . . Disci, R. (2005). The effect of venlafaxine on ongoing and experimentally induced pain in neuropathic pain patients: a double blind, placebo controlled study. *European journal of pain*, 9(4), 407-407.
- Yuen, K. C., Baker, N. R., & Rayman, G. (2002). Treatment of chronic painful diabetic neuropathy with isosorbide dinitrate spray. *Diabetes Care*, 25(10), 1699-1703.
- Zeigler, D., Lynch, S. A., Muir, J., Benjamin, J., & Max, M. B. (1992). Transdermal clonidine versus placebo in painful diabetic neuropathy. *Pain*, 48(3), 403-408.
- Ziegler, D., Hidvégi, T., Gurieva, I., Bongardt, S., Freynhagen, R., Sen, D., . . . Group, L. S. S. (2010). Efficacy and safety of lacosamide in painful diabetic neuropathy. *Diabetes Care*, 33(4), 839-841.
- Österberg A., B. J. (2005). *Central Pain in Multiple sclerosis- A double-blind placebo-controlled trial of amitriptyline and carbamazepine. In: Österberg A, editor. Central Pain in multiple sclerosis – clinical characteristics, sensory abnormalities, and treatment. Thesis., Linköpings Universitet.*