

Supplemental Table 1. Cohort characteristics

Characteristic	HIV-Infected (n=105)	HIV-Uninfected	P-value
Age (years)	49 (45-51)	50 (46-54)	0.06
Female (%)	51	50	0.84
BMI (kg/m ²)	22.8±4	22.9±5	0.47
Ever-smoker (%)	36	50	0.06
Median Non-HDL cholesterol mg/dL (IQR)	107 (89 – 137)	112.5 (93 – 136)	0.74
Median Hemoglobin A1c (IQR)	5.3 (5.0 – 5.7)	5.6 (5.3 – 5.9)	0.86
ART Regimen (%)	80 (85)	N/A	
AZT/3TC/NVP	68 (65)		
AZT/3TC/EFV	21 (20)		
TDF/3TC/LPV/R	9 (9)		
Other	7 (7)		
Median ART Duration (IQR)	7.0 (6.4 – 7.5)	N/A	
Median CD4 Count (IQR)	430 (334 – 546)	N/A	
Undetectable Viral Load (%)	80 (85)	N/A	
Ankle-brachial index (%)			0.01
<0.9	1 (1)	0 (0)	
0.9 – 1.2	77 (74)	90 (90)	
>1.2	27 (26)	10 (10)	

Figure S1. Prevalence of arterial stiffness among older-aged individuals with and without HIV infection in rural Uganda. Error bars represent the upper bound of the 95% confidence interval of the estimate.

