

Fig S1. Kawamoto et al.




Fig S2. Kawamoto et al.




Fig S3. Kawamoto et al.




Fig S4. Kawamoto et al.




Fig S5. Kawamoto et al.




Fig S6. Kawamoto et al.




Fig S7. Kawamoto et al.




Fig S8. Kawamoto et al.




Fig S9. Kawamoto et al.




Fig S10. Kawamoto et al.




Fig S11. Kawamoto et al.




Fig S12. Kawamoto et al.




Fig S13. Kawamoto et al.




Fig S14. Kawamoto et al.




Fig S15. Kawamoto et al.




Table S1. Clinical parameters of the pneumonia- and non pneumonia-etiology subgroups in sepsis

	Sepsis with pneumonia (n=12)	Sepsis without pneumonia (n=15)	p-Value
Age, years	62.6±20.6	73.2±15.3	0.152
Female/Male	3/9	8/7	0.217
TP, g/dL	5.5±1.4	5.5±1.2	0.721
Alb, g/dL	2.6±0.8	2.8±0.6	0.878
BUN, mg/dL	29.1±16.7	26.5±17.7	0.719
Cre, mg/dL	0.8±0.4	1.3±1.1	0.167
AST, U/L	110.3±137.8	160.0±309.0	0.516
ALT, U/L	70.0±87.5	120.5±245.9	0.399
T-bil, mg/dL	1.8±4.4	1.2±0.9	0.27
CRP, mg/dL	2.6±6.5	12.7±8.8	0.943
WBC count, ×10 ⁹ /L	12.1±7.8	16.9±10.0	0.277
Hb, g/dL	10.8±2.3	10.6±1.90	0.648
Hct, %	32.6±6.4	32.1±5.4	0.581
Neutrophil, ×10 ⁹ /L	11.1±7.7	14.1±8.5	0.467
Lymphocyte, ×10 ⁹ /L	0.6±0.7	0.5±0.4	0.959
Monocyte, ×10 ⁹ /L	0.6±0.5	0.5±0.4	0.851
Platelet, ×10 ³ /μL	212.8±150.0	145.6±81.5	0.277
SOFA score	7.5±4.8	9.0±4.8	0.399
P/F ratio	252.3±96.4	267.2±86.3	0.905
mBP, mmHg	60.1±12.2	61.4±12.5	0.755
GCS	9.5±4.4	9.8±4.0	0.905
28-day mortality	7 (58.3%)	3 (20.0%)	0.093

Results are expressed as mean +/- SEM.