

Supplemental Digital Content (SDC)

Table 1 Immunosuppressive regimen at 1-year and 3-year visits^a.

	Tac/Bas n = 143		Tac/MMF n = 139		Triple therapy n = 139	
	Year 1	Year 3	Year 1	Year 3	Year 1	Year 3
Tacrolimus mean dose (SD) [mg]	5.9 (2.99)	4.9 (2.22)	5.6 (2.96)	4.5 (2.24)	5.5 (2.87)	4.4 (2.21)
Tacrolimus mean trough levels (SD) [ng/mL]	9.34 (3.66)	8.3 (3.01)	9.19 (2.69)	7.9 (2.46)	9.24 (3.10)	8.1 (2.69)
Tacrolimus monotherapy ^b (%)	58.2	48.5	5.6	9.3	0.8	3.0
Steroid free ^b (%)	67.5	64.0	72.6	77.6	16.7	33.7
Prednisolone mean dose (SD) [mg]	9.6 (6.90)	5.2 (1.75)	5.0 (3.54)	6.3 (2.50)	4.3 (1.14)	5.7 (2.83)
MMF use ^b (%)	22.8	29.0	83.9	79.4	88.3	79.2
MMF mean dose (SD) [g]	1.2 (0.49)	1.1 (0.44)	0.91 (0.26)	0.92 (0.33)	0.91 (0.21)	0.93 (0.32)

^aPatients providing 3-year follow-up data

^bp<0.01 at 1 year and at 3 years, Chi-squared test

SD = standard deviation