

Table S1: Characteristics of patients with and without an echocardiogram

	Patients with echocardiogram (n=142)	Patients without echocardiogram (n=79)	P value
Age, years*	16.9 (13.6-19.4)	17 (14-19.5)	NS
Age ≥18, n (%)	58 (41)	30 (37)	NS
Male Gender (%)	88 (62)	48 (61)	NS
African-American/Hispanic, n (%)	35/130 (27)	24/72 (33)	NS
Age at Transplantation, years*	12 (7.1-14.8)	12.1 (7.4-15.6)	NS
Time post Transplantation, years*	3.9 (1.2-7)	3.2 (1.4-7)	NS
Acquired Glomerular Disease, n (%)	29 (20)	22 (28)	NS
First Transplant, n (%)	132 (93)	75 (95)	NS
Living Donors, n (%)	85 (60)	45 (60)	NS
Prior Dialysis, n (%)	90 (64)	55 (70)	NS
History of Rejection, n (%)	37 (26)	26 (32)	NS
Immunosuppression, n (%)			
Calcineurin Inhibitors	113 (80)	63 (84)	NS
Steroids	67 (48)	27 (36)	NS
Mycophenolate Mofetil	109 (77)	62 (82)	NS
Overweight/Obese, n (%)	52 (37)	26 (34)	NS
Hemoglobin, g/dL*	11.9 (10.9-13.3)	12.8 (11.6-13.5)	0.018
Anemia, n (%)	38 (27)	14 (17)	NS
Treated for hypertension, n (%)	94 (66)	61 (75)	NS
≥2 BP medications, n (%)	36 (25)	23 (28)	NS
eGFR, Schwartz/MDRD (ml/min/1.73m ²)*	66 (50-85)	63 (47-80)	NS
eGFR <60 ml/min/1.73m ² , n (%)	61 (43)	38 (48)	NS
Abnormal ABP, n (%)	64 (45)	37 (47)	NS