

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article

Table S1: Baseline characteristics of patients in cohort 2 on the NKTWL 2000-2010, stratified by suspension status.

	Not suspended in first year		Suspended in first year		Total
N	8946	(100.0%)	3292	(100.0%)	
Sex					
Male	5628	(62.9%)	1960	(59.5%)	0.002
Female	3311	(37.0%)	1332	(40.5%)	
Unknown	7	(0.1%)	0	(0.0%)	
Ethnicity Group					
White	7266	(81.2%)	2356	(71.6%)	<0.0001
Asian	562	(6.3%)	535	(16.3%)	
Black	504	(5.6%)	312	(9.5%)	
Other	204	(2.3%)	87	(2.6%)	
Unknown	10	(0.1%)	0	(0.0%)	
Age Group					
18-34	1593	(17.8%)	441	(13.4%)	<0.0001
35-49	3221	(36.0%)	1098	(33.4%)	
50-59	2326	(26.0%)	1029	(31.3%)	
60-69	1568	(17.5%)	641	(19.5%)	
70+	238	(2.7%)	83	(2.5%)	
Disease Group					
Hypertension/Renovascular Disease	511	(5.7%)	250	(7.6%)	<0.0001
Diabetes	629	(7.0%)	285	(8.7%)	
Polycystic kidneys	1257	(14.1%)	469	(14.2%)	
Pyelonephritis/Interstitial Nephritis	710	(7.9%)	226	(6.9%)	
Glomerulonephritis	1504	(16.8%)	513	(15.6%)	
Other	1245	(13.9%)	460	(14.0%)	
Not Reported	3090	(34.5%)	1089	(33.1%)	
Highly Sensitised					
Not highly sensitized	8832	(98.7%)	3221	(97.8%)	0.0004
Highly Sensitized	114	(1.3%)	71	(2.2%)	