

E-REFERENCES

- e1. Zheng ZS, Reggente N, Lutkenhoff E, Owen AM, Monti MM. Disentangling disorders of consciousness: Insights from diffusion tensor imaging and machine learning. *Hum Brain Mapp* 2017;38:431-443.
- e2. Naro A, Russo M, Leo A, et al. Cortical responsiveness to nociceptive stimuli in patients with chronic disorders of consciousness: do C-fiber laser evoked potentials have a role? *PLoS One* 2015;10:e0144713.
- e3. Sazbon L, Zagreba F, Ronen J, Solzi P, Costeff H. Course and outcome of patients in vegetative state of nontraumatic aetiology. *J Neurol Neurosurg Psychiatry* 1993;56:407-409.
- e4. Cavinato M, Freo U, Ori C, et al. Post-acute P300 predicts recovery of consciousness from traumatic vegetative state. *Brain Inj* 2009;23:973-980.
- e5. Sara M, Pistoia F, Pasqualetti P, Sebastiano F, Onorati P, Rossini PM. Functional isolation within the cerebral cortex in the vegetative state: A nonlinear method to predict clinical outcomes. *Neurorehabil Neural Repair* 2011;25:35-42.
- e6. Luaute J, Maucort-Boulch D, Tell L, et al. Long-term outcomes of chronic minimally conscious and vegetative states. *Neurology* 2010;75:246-252.
- e7. Skandsen T, Lund TI, Fredriksli O, Vik A. Global outcome, productivity and epilepsy 3-8 years after severe head injury. The impact of injury severity. *Clin Rehab* 2008;22:653-662.
- e8. Noe E, Olaya J, Navarro MD, et al. Behavioral recovery in disorders of consciousness: a prospective study with the Spanish version of the Coma Recovery Scale-Revised. *Arch Phys Med Rehabil* 2012;93:428-433.
- e9. Danze F, Veys B, Lebrun T, et al. Prognostic factors of post-traumatic vegetative states: 522 cases. *Neurochirurgie* 1994;40:348-357; discussion 357-348.

- e10. Sazbon L, Grosswasser Z. Outcome in 134 patients with prolonged posttraumatic unawareness. Part 1: Parameters determining late recovery of consciousness. *J Neurosurg* 1990;72:75-80.
- e11. Pascarella A, Trojano L, Loreto V, Bilo L, Moretta P, Estraneo A. Long-term outcome of patients with disorders of consciousness with and without epileptiform activity and seizures: a prospective single centre cohort study. *J Neurol* 2016;263:2048-2056.
- e12. Bagnato S, Boccagni C, Sant'Angelo A, Fingelkurt AA, Fingelkurt AA, Galardi G. Longitudinal assessment of clinical signs of recovery in patients with unresponsive wakefulness syndrome after traumatic or nontraumatic brain injury. *J Neurotrauma* 2017;34:535-539.
- e13. Mateen FJ, Niu JW, Gao S, et al. Causes and outcomes of persistent vegetative state in a Chinese versus American referral hospital. *Neurocrit Care* 2013;18:266-270.
- e14. Estraneo A, Moretta P, Loreto V, et al. Predictors of recovery of responsiveness in prolonged anoxic vegetative state. *Neurology* 2013;80:464-470.
- e15. Tsubokawa T, Yamamoto T, Katayama Y. Prediction of outcome of prolonged coma caused by brain damage. *Brain Inj* 1990;4:329-337.
- e16. Pape TL, Heinemann AW, Kelly JP, Hurder AG, Lundgren S. A measure of neurobehavioral functioning after coma. Part I: Theory, reliability, and validity of Disorders of Consciousness Scale. *J Rehabil Res Dev* 2005;42:1-17.
- e17. Dolce G, Lucca LF, Candelieri A, Rogano S, Pignolo L, Sannita WG. Visual pursuit in the severe disorder of consciousness. *J Neurotrauma* 2011;28:1149-1154.
- e18. Wilson SL, Gill-Thwaites H. Early indication of emergence from vegetative state derived from assessments with the SMART--a preliminary report. *Brain Inj* 2000;14:319-331.

- e19. Beis JM, Seyer JL, Brugeronne B, et al. Care protocol for persistent vegetative states (PVS) and minimally conscious state (MSC) in Lorraine: retrospective study over an 18-year period. Ann Phys Rehabil Med 2009;52:374-381.
- e20. Bagnato S, Minafra L, Bravata V, et al. Brain-derived neurotrophic factor (Val66Met) polymorphism does not influence recovery from a post-traumatic vegetative state: a blinded retrospective multi-centric study. J Neurotrauma 2012;29:2050-2059.
- e21. Kotchoubey B, Lang S, Mezger G, et al. Information processing in severe disorders of consciousness: Vegetative state and minimally conscious state. Clin Neurophysiol 2005;116:2441-2453.
- e22. Giacino JT, Kalmar K. The vegetative and minimally conscious states: a comparison of clinical features and functional outcome. J Head Trauma Rehabilitation 1997;12:36-51.
- e23. Estraneo A, Moretta P, Loreto V, Lanzillo B, Santoro L, Trojano L. Late recovery after traumatic, anoxic, or hemorrhagic long-lasting vegetative state. Neurology 2010;75:239-245.
- e24. Giacino JT. Disorders of consciousness: differential diagnosis and neuropathologic features. Semin Neurol 1997;17:105-111.
- e25. Groswasser Z, Sazbon L. Outcome in 134 patients with prolonged posttraumatic unawareness. Part 2: Functional outcome of 72 patients recovering consciousness. J Neurosurg 1990;72:81-84.
- e26. Nayak P, Mahapatra A. Single photon emission computed tomography scanning: A predictor of outcome in vegetative state of head injury. J Neurosci Rural Pract 2011;2:12-16.
- e27. Goodwin L. Use of MRI in prediction of recovery from persistent vegetative state. J Insur Med 1998;30:113-114.

- e28. Wang F, Di H, Hu X, et al. Cerebral response to subject's own name showed high prognostic value in traumatic vegetative state. *BMC Med* 2015;13:83.
- e29. Whyte J, Katz D, Long D, et al. Predictors of outcome in prolonged posttraumatic disorders of consciousness and assessment of medication effects: A multicenter study. *Arch Phys Med Rehabil* 2005;86:453-462.
- e30. Sazbon L, Fuchs C, Costeff H. Prognosis for recovery from prolonged post-traumatic unawareness: Logistic analysis. *Journal of Neurology Neurosurgery and Psychiatry* 1991;54:149-152.
- e31. Zheng X, Chen M, Li J, Cao F. Prognosis in prolonged coma patients with diffuse axonal injury assessed by somatosensory evoked potentia. *Neural Regen Res* 2013;8:948-954.
- e32. Vogel D, Markl A, Yu T, Kotchoubey B, Lang S, Muller F. Can mental imagery functional magnetic resonance imaging predict recovery in patients with disorders of consciousness? *Arch Phys Med Rehabil* 2013;94:1891-1898.
- e33. Ganesh S, Guernon A, Chalcraft L, Harton B, Smith B, Louise-Bender Pape T. Medical comorbidities in disorders of consciousness patients and their association with functional outcomes. *Arch Phys Med Rehabil* 2013;94:1899-1907.
- e34. Arnaldi D, Terzaghi M, Cremascoli R, et al. The prognostic value of sleep patterns in disorders of consciousness in the sub-acute phase. *Clin Neurophysiol* 2016;127:1445-1451.
- e35. Pape TL, Tang C, Guernon A, et al. Predictive value of the Disorders of Consciousness Scale (DOCS). *PMR* 2009;1:152-161.
- e36. Heindl UT, Laub MC. Outcome of persistent vegetative state following hypoxic or traumatic brain injury in children and adolescents. *Neuropediatrics* 1996;27:94-100.

- e37. Kriel RL, Krach LE, Jones-Saete C. Outcome of children with prolonged unconsciousness and vegetative states. *Pediat Neurol* 1993;9:362-368.
- e38. Giacino JT, Whyte J, Bagiella E, et al. Placebo-controlled trial of amantadine for severe traumatic brain injury. *N Engl J Med* 2012;366:819-826.
- e39. Krewer C, Luther M, Koenig E, Muller F. Tilt table therapies for patients with severe disorders of consciousness: a randomized, controlled trial. *PLoS One* 2015;10:e0143180.
- e40. Herrold AA, Pape TL, Guernon A, Mallinson T, Collins E, Jordan N. Prescribing multiple neurostimulants during rehabilitation for severe brain injury. *ScientificWorldJournal* 2014;2014:964578.
- e41. Nakase-Richardson R, Whyte J, Giacino JT, et al. Longitudinal outcome of patients with disordered consciousness in the NIDRR TBI Model Systems Programs. *J Neurotrauma* 2012;29:59-65.
- e42. Katz DI, Polyak M, Coughlan D, Nichols M, Roche A. Natural history of recovery from brain injury after prolonged disorders of consciousness: outcome of patients admitted to inpatient rehabilitation with 1-4 year follow-up. *Prog Brain Res* 2009;177:73-88.