

Table e-1. The genotyping results of known *C9orf72* genetic modifiers (*TMEM106B* rs1990622, *TMEM106B* rs3173615 and *ATXN2* CAG repeats).

Sample	TMEM106B rs1990622	TMEM106B rs3173615	ATXN2 CAG repeats
9548	AA	CC	22/22
9686	AG	CG	22/22

Note: Intermediate *ATXN2* alleles (27–33 CAG-repeats) were reported as modifiers in *C9orf72* carriers, rendering susceptibility to ALS ⁷. The homozygosity for the minor allele (G) of rs3173615 in *TMEM106B* was reported to protect against developing FTD in *C9orf72* patients ⁸. The major allele (A) of rs1990622 (conferring risk for developing FTLD) was associated to later age of onset and age of death in *C9orf72* patients ⁹. These results were not able to explain the phenotype difference between 9548 and 9686.

Table e-2. The CpG methylation status in the 5' region close to the *C9orf72* repeats in the tissues of 9686. M represents methylated site; U represents unmethylated site.

CpG site	Frontal cortex	Motor cortex	Occipital cortex	Hippocampus	Thalamus	Brain stem	Temporal cortex	Cerebellum	Heart	Liver	Kidney	Lung	Tongue	Testicles	Epididymis	Cervical spinal cord	Thoracic spinal cord	Lumbar spinal cord
1	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
2	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
3	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
4	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
5	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
6	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
7	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
8	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
9	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
10	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
11	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
12	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
13	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
14	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
15	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
16	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
17	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
18	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
19	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
20	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
21	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
22	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
23	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
24	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
25	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
26	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	

Table e-3. The CpG methylation status in the 3' region close to the *C9orf72* repeats in the tissues of 9686. M represents methylated site; U represents unmethylated site.

CpG site	Frontal cortex	Motor cortex	Occipital cortex	Hippocampus	Thalamus	Brain stem	Temporal cortex	Cerebellum	Heart	Liver	Kidney	Lung	Tongue	Testicles	Epididymis	Cervical spinal cord	Thoracic spinal cord	Lumbar spinal cord
1	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
2	U	U	U	U	U	M	U	U	U	U	U	U	U	U	U	U	U	
3	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
4	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
5	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
6	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
7	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
8	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
9	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	M	U	
10	U	U	U	M	U	U	U	U	M	U	M	U	U	U	U	M	U	
11	U	U	M	M	U	U	U	U	U	M	U	U	U	U	U	U	U	
12	U	U	M	U	U	U	U	U	U	M	U	M	U	U	U	M	M	
13	U	U	U	U	U	U	U	U	U	M	U	U	U	U	U	U	U	
14	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	M	
15	U	U	U	U	U	M	U	U	U	U	U	U	U	U	U	U	U	
16	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
17	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
18	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
19	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	M	
20	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
21	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	
22	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	

Table e-4. *C9orf72* repeat size in multiple tissues of small expansion carriers according to previous studies. NA represents not available.

Studies [#]	Sample ID	Pheno-type	Age of onset, years	Duration, years	Blood	Frontal cortex	Temporal cortex	Cerebellum	Spinal cord	Occipital	Parietal lobe	Spleen	Heart	Muscle	Kidney	Dura Mater	Cauda Equina	RNA foci	DPR	TDP43
Van Blitterswijk et al. 2013 ¹⁰	P20	MND-FTD	NA	NA	small	large(smear)	Large (smear)	large (dominant)	small/ large(smear)	small/ large(smear)	small/ large	small/ medium	Small	small/ medium						
Fratta et al. 2015 ¹¹	61776	ALS	57	6	small (90)	small/large		large (dominant)	small/large						small	small	SF and ASF: FC, HC, CBL, HC	HC, CBL	SC	
Gijsselsink et al. 2016 ¹²	DR439.1	FTLD	54	15	small (56)	small(weak)/ large(smear)	small(weak)/ large(smear)	large (dominant)									FC, HC, CBL	HC, FC		
Nordin et al. 2015 ¹³	P9	ALS	63	2	small (61)	small(weak)/ large		large (dominant)	small/large		small/ large	small/ large	small	small	small					
Nordin et al. 2015 ¹³	P11	MND	75	5	small (92)	small(weak)/ large		large (dominant)	small(weak)/large				small							

Only those studies that had both CNS and non-CNS tissues were included.

Reference:

1. Xi Z, van Blitterswijk M, Zhang M, et al. Jump from pre-mutation to pathologic expansion in *C9orf72*. American journal of human genetics 2015;96:962-970.
2. Xi Z, Zhang M, Bruni AC, et al. The *C9orf72* repeat expansion itself is methylated in ALS and FTLD patients. Acta neuropathologica 2015;129:715-727.
3. Xiao S, MacNair L, McGoldrick P, et al. Isoform-specific antibodies reveal distinct subcellular localizations of *C9orf72* in amyotrophic lateral sclerosis. Annals of neurology 2015;78:568-583.
4. Freibaum BD, Taylor JP. The Role of Dipeptide Repeats in C9ORF72-Related ALS-FTD. Frontiers in molecular neuroscience 2017;10:35.
5. Mackenzie IR, Frick P, Grasser FA, et al. Quantitative analysis and clinico-pathological correlations of different dipeptide repeat protein pathologies in *C9orf72* mutation carriers. Acta neuropathologica 2015;130:845-861.

6. Davidson Y, Robinson AC, Liu X, et al. Neurodegeneration in frontotemporal lobar degeneration and motor neurone disease associated with expansions in C9orf72 is linked to TDP-43 pathology and not associated with aggregated forms of dipeptide repeat proteins. *Neuropathology and applied neurobiology* 2016;42:242-254.
7. van Blitterswijk M, Mullen B, Heckman MG, et al. Ataxin-2 as potential disease modifier in C9ORF72 expansion carriers. *Neurobiology of aging* 2014;35:2421 e2413-2427.
8. van Blitterswijk M, Mullen B, Nicholson AM, et al. TMEM106B protects C9ORF72 expansion carriers against frontotemporal dementia. *Acta neuropathologica* 2014;127:397-406.
9. Gallagher MD, Suh E, Grossman M, et al. TMEM106B is a genetic modifier of frontotemporal lobar degeneration with C9orf72 hexanucleotide repeat expansions. *Acta neuropathologica* 2014;127:407-418.
10. van Blitterswijk M, DeJesus-Hernandez M, Niemantsverdriet E, et al. Association between repeat sizes and clinical and pathological characteristics in carriers of C9ORF72 repeat expansions (Xpansize-72): a cross-sectional cohort study. *The Lancet Neurology* 2013;12:978-988.
11. Fratta P, Polke JM, Newcombe J, et al. Screening a UK amyotrophic lateral sclerosis cohort provides evidence of multiple origins of the C9orf72 expansion. *Neurobiology of aging* 2015;36:546 e541-547.
12. Gijselinck I, Van Mossevelde S, van der Zee J, et al. The C9orf72 repeat size correlates with onset age of disease, DNA methylation and transcriptional downregulation of the promoter. *Molecular psychiatry* 2016;21:1112-1124.
13. Nordin A, Akimoto C, Wuolikainen A, et al. Extensive size variability of the GGGGCC expansion in C9orf72 in both neuronal and non-neuronal tissues in 18 patients with ALS or FTD. *Human molecular genetics* 2015;24:3133-3142.