

Table e4. Correlations^a between visual function testing and magnetic resonance imaging parameters at each time point.

	Baseline		52 weeks		104 weeks	
	HCVA (100%)	LCVA (2.5%)	HCVA (100%)	LCVA (2.5%)	HCVA (100%)	LCVA (2.5%)
T1-hypointense lesion volume (mm ³)	-0.15 (<i>p</i> <0.0001)	-0.17 (<i>p</i> <0.0001)	-0.18 (<i>p</i> <0.0001)	-0.19 (<i>p</i> <0.0001)	-0.16 (<i>p</i> <0.0001)	-0.14 (<i>p</i> <0.0001)
T2 lesion volume (mm ³)	-0.13 (<i>p</i> <0.0001)	-0.21 (<i>p</i> <0.0001)	-0.13 (<i>p</i> <0.0001)	-0.16 (<i>p</i> <0.0001)	-0.16 (<i>p</i> <0.0001)	-0.18 (<i>p</i> <0.0001)
Gd+ lesion volume (mm ³)	-0.02 (<i>p</i> =0.5)	-0.10 (<i>p</i> <0.01)	-0.02 (<i>p</i> =0.5)	-0.06 (<i>p</i> =0.07)	-0.01 (<i>p</i> =0.76)	-0.04 (<i>p</i> =0.23)
No. new T1 lesions	NA	NA	-0.04 (<i>p</i> =0.22)	-0.09 (<i>p</i> =0.01)	-0.001 (<i>p</i> =0.67)	-0.07 (<i>p</i> =0.06)
No. new T2 lesions	NA	NA	-0.06 (<i>p</i> =0.07)	-0.10 (<i>p</i> <0.01)	-0.03 (<i>p</i> =0.459)	-0.06 (<i>p</i> =0.08)
No. Gd+ lesions	-0.03 (<i>p</i> =0.4)	-0.13 (<i>p</i> <0.0001)	-0.04 (<i>p</i> =0.268)	-0.06 (<i>p</i> =0.07)	-0.02 (<i>p</i> =0.5)	-0.06 (<i>p</i> =0.07)
BPF value	0.06 (<i>p</i> =0.07)	0.10 (<i>p</i> <0.01)	0.03 (<i>p</i> =0.32)	0.10 (<i>p</i> <0.01)	0.08 (<i>p</i> =0.02)	0.13 (<i>p</i> <0.0001)

Correlations^a between visual function testing and magnetic resonance imaging parameters at each time point.