

Table E1. Demographics and clinical manifestations of patients and normal subjects

	AD	MCI	Control	P Value
n	12	19	21	
M:F	5F7M	12F7M	14F7M	NS
Age (yrs)	73.3 ± 9.6	69.6 ± 9.8	67.6 ± 8.3	NS
SBP (mmHg)	124.8 ± 19.8	121.7 ± 14.0	129.9 ± 16.0	NS
DBP (mmHg)	77.2 ± 14.0	75.4 ± 8.2	79.5 ± 11.5	NS
HR (beats per minute)	63.2 ± 16.0	65.9 ± 10.0	63.6 ± 10.1	NS
MMSE	19.9 ± 5.6	25.7 ± 2.5	29.5 ± 0.8*	P < 0.01**
Disease Duration (yrs)	4.0 ± 1.5	3.5 ± 2.8		NS
Smoking	0	0	0	NS
Hypertension	5	12	13	NS
Dyslipidemia	6	11	10	NS
Type 2 Diabetes	1	1	2	NS
OCTA Image Strength ^Ψ	7.9 ± 1.6	7.7 ± 1.4	7.5 ± 1.3	NS

NS: non-significance; SBP: Systolic blood pressure; DBP: Diastolic blood pressure; HR: hear rate

MMSE: Mini-Mental State Examination; Disease Duration: years after onset of symptoms

*MMSE was not available from one control subject; therefore, data from 20 controls were used.

** Post hoc tests indicated significant differences between each pair.

^Ψ ranged from 5 to 10.