

TABLE 2. Orbits displaying optic canal narrowing

Orbit	Patient	Time since diagnosis (years)	Side	Walls				Total	Visual Function	
				Lateral	Floor	Medial	Roof		Visual Acuity	Visual Fields
1	1	6	R	+	+	+	+	4	20/20	N
2	1	-	L	+	-	+	+	3	NLP	NLP
3	2	5	R	+	+	+	+	4	20/20	N
4	2	-	L	+	+	+	+	4	20/20	N
5	4	1	R	+	-	+	-	2	20/30	N
6	7	2	R	-	-	-	-	0	20/20	N
7	7	-	L	+	-	+	+	3	20/20	RCS
8	9	2	R	-	-	+	+	2	20/20	DLS
9	12	1	R	-	-	+	+	2	20/20	N
10	12	-	L	-	-	-	+	1	20/20	N
11	13	4	L	+	+	+	+	4	20/20	N
12	14	8	R	+	+	-	-	2	20/20	N
13	14	-	L	+	+	+	+	4	20/20	RCS
14	15	4	R	+	+	+	+	4	20/20	N
15	15	-	L	+	-	+	+	3	20/20	N
16	17	13	R	+	-	-	+	2	20/20	N
17	17	-	L	+	-	-	+	2	20/20	N
18	20	20	R	+	+	+	+	4	20/20	N
19	20	6	L	-	-	-	-	0	20/20	N

NLP, no light perception; RCS, relative central scotoma; DLS, diffuse loss of sensibility.