

## **Appendix. Acknowledgements**

This study was made possible through the co-operation and collective efforts of the Nordic transvaginal mesh group, listed in alphabetical order:

### **Denmark**

Susanne Axelsen, Karl Möller Bek, Skejby Hospital, Skejby, Jens Prien Larsen, Nyköbing Hospital, Nyköbing.

### **Finland**

Pentti Kiilholma, Riikka Aaltonen, Turku University Hospital, Turku, Teuvo Takala, Tiina Vaari, Päijät-Häme Central Hospital, Lahti, Tapi Väyrynen, Helsinki University Central Hospital, Jorvi Hospital, Espoo, Hans-Anders Unnérus, Helsinki University Central Hospital, Lohja Hospital, Lohja.

### **Norway**

Marie Elström Engh, Akershus University Hospital, Ahus, Kolbjörn Kasin, Kongsberg Hospital, Kongsberg, Rolf Kirschner, Rikshospitalet, Oslo, Anne-Grethe Paulsen, Haukeland Hospital, Bergen, Sigurd Kulseng Hansen, Asker & Bærum Hospital, Bærum, Margareta Verelst, Regional Hospital in Tromsø, Tromsø.

### **Sweden**

Daniel Altman, Danderyd Hospital, Stockholm, Bo Anzén, Danderyd Hospital, Stockholm, Lena Clementsson, Kristiansstad Hospital, Kristiansstad, Christian Falconer, Danderyd Hospital, Stockholm, Folke Flam, S:t Göran Hospital, Stockholm, Eva Fornell, Linköping University Hospital, Linköping, Jonas Gunnarsson, Sahlgrenska University Hospital, Göteborg, Björn Holm, Capio Läkargruppen, Örebro, Kenneth Krohn, Vrinnevi Hospital, Norrköping, Nina Kyrklund, Danderyd Hospital, Stockholm, Gregor Larson, Västerås Hospital, Västerås, P-G

© 2009 American College of Obstetricians and Gynecologists.

Larsson, Skaraborg Hospital Skövde, Skövde, Paula Löwenadler, Ystad Hospital, Ystad, Mazoumeh Rezapour, Uppsala University Hospital, Uppsala, Gunilla Tegerstedt, South Hospital, Stockholm, Vasiliki Tsaknakis, Halmstad Hospital, Halmstad, Jan Zetterström, Danderyd Hospital, Stockholm.