"Video" Accompanying: Measure to Quantify the Influence of Time from End of Surgery to Tracheal Extubation on Operating Room Workflow



Franklin Dexter, M.D., Ph.D. Director, Division of Management Consulting Professor, Department of Anesthesia University of Iowa Franklin-Dexter@UIowa.edu www.FranklinDexter.net





This "video" lasts 18 minutes
 – Extubation 15.0 minutes after end of surgery



- This "video" lasts 18 minutes
  - Extubation 15.0 minutes after end of surgery
- One person remaining in the OR will be idle for at least one minute during the 15 minutes





- This "video" lasts 18 minutes
  - Extubation 15.0 minutes after end of surgery
- One person remaining in the OR will be idle for at least one minute during the 15 minutes
   – Representative situation (see Table 1)



- This "video" lasts 18 minutes
  - Extubation 15.0 minutes after end of surgery
- One person remaining in the OR will be idle for at least one minute during the 15 minutes
   – Representative situation (see Table 1)
- To learn how to evaluate if there is an idle person, do not speed up the video



- This "video" lasts 18 minutes
  - Extubation 15.0 minutes after end of surgery
- One person remaining in the OR will be idle for at least one minute during the 15 minutes
   – Representative situation (see Table 1)
- To learn how to evaluate if there is an idle person, do not speed up the video
  - 15 minutes is a long time, which is why evaluating if there has been idle person is easy







 Having perfect (magical) retrospective knowledge that extubation will occur at 15.0 minutes, it is necessary only to have observed for the last 2 minutes





- Having perfect (magical) retrospective knowledge that extubation will occur at 15.0 minutes, it is necessary only to have observed for the last 2 minutes
  - Reason is that people tend to move around until they start to be idle



- Having perfect (magical) retrospective knowledge that extubation will occur at 15.0 minutes, it is necessary only to have observed for the last 2 minutes
  - Reason is that people tend to move around until they start to be idle
- Looking for idleness is straightforward because usually all that is effectively being done is watching just before extubation



- Having perfect (magical) retrospective knowledge that extubation will occur at 15.0 minutes, it is necessary only to have observed for the last 2 minutes
  - Reason is that people tend to move around until they start to be idle

 Looking for idleness is straightforward because usually all that is effectively being done is watching just before extubation

- "Rewind" to 13.5 minutes emphasizes point







All figures are identical in appearance

 You will lose track of who each person is and his/her role, and that is *realistic*



- You will lose track of who each person is and his/her role, and that is *realistic*
  - Goal is not to track activities



- You will lose track of who each person is and his/her role, and that is *realistic*
  - Goal is not to track activities
  - Goal is to evaluate if a person has been idle for at least one minute



- You will lose track of who each person is and his/her role, and that is *realistic*
  - Goal is not to track activities
  - Goal is to evaluate if a person has been idle for at least one minute
- Movements in this video are realistic



- All figures are identical in appearance
  - You will lose track of who each person is and his/her role, and that is *realistic*
    - Goal is not to track activities
    - Goal is to evaluate if a person has been idle for at least one minute
- Movements in this video are realistic
  - When surgery resident starts being idle he remains idle until extubation, which is realistic
































































































.





.


























































































































## **Use of Stick Figures**

All figures are identical in appearance

- You will lose track of who each person is and his/her role, and that is *realistic* 
  - Goal is not to track activities
  - Goal is to evaluate if a person has been idle for at least one minute

Movements in this video are realistic

 When surgery resident starts being idle he remains idle until extubation, which is realistic



























































































































































































































































































## When Finished, the Video Rewinds to 13.5 Minutes

- Having perfect (magical) retrospective knowledge that extubation will occur at 15.0 minutes, it is necessary only to have observed for the last 2 minutes
  - Reason is that people tend to move around until they start to be idle

 Looking for idleness is straightforward because usually all that is effectively being done is watching just before extubation

– "Rewind" to 13.5 minutes emphasizes point





















































## Script

- Danielle Masursky
  Franklin Dexter
- Animation
  - Jodi Kazerani
- Editing

   Franklin Dexter

## Additional Information on Operating Room Management

## www.FranklinDexter.net/education.htm

- Example reports with calculations
- Lectures on day of surgery decision making, PACU staffing, OR allocation and staffing, anesthesia staffing, financial analysis, comparing surgical services among hospitals, strategic decision making, and drug costs
- www.FranklinDexter.net
  - Comprehensive bibliography of peer reviewed articles in operating room and anesthesia group management