

**Supplemental Figure 1: Cleveland Clinic Phase II Discharge Tool**

Phase II Discharge Scoring Tool	PRE	POST	POST	POST
Time				
Level of consciousness 2 = Fully awake 1 = Arousable with verbal or tactile stimulation 0 = Unresponsive				
Physical activity (Return to preanesthetic level) 2 = Steady gait, no dizziness or meets pre-op level 1 = Requires assistance 0 = Unable to ambulate				
Blood Pressure 2 = Systolic blood pressure +/- $\leq$ 20% of preanesthetic/procedural sedation level 1 = Systolic blood pressure +/- $\leq$ 30% of preanesthetic/procedural sedation level 0 = Systolic blood pressure +/- $>$ 30% of preanesthetic/procedural sedation level	BP			
Heart Rate 2 = 60-100 or if outside range, $\leq$ to 10% change from baseline HR 1 = Outside of the range of 60-100 and the change from baseline is $>$ 10% and $\leq$ 20% 0 = Outside of the range of 60-100 and the change from baseline is $>$ 20%	HR			
Oxygen saturation status 2 = Saturation $\geq$ 92% on room air, or pre preanesthetic/procedural sedation level 1 = Saturation $\geq$ 92% on supplemental oxygen not involving IV PCA 0 = Saturation $<$ 92% on supplemental oxygen	O <sub>2</sub> SAT			
Pain 2 = None or mild pain with/without analgesics 1 = Moderate pain controlled with analgesics 0 = Persistent severe pain uncontrolled with analgesics				
Postoperative nausea / vomiting 2 = None or mild nausea with no active vomiting 1 = Moderate nausea or transient vomiting 0 = Persistent severe nausea or vomiting				
Bleeding 2 = Minimal or no dressing changes 1 = Moderate – up to two dressing changes required 0 = Severe – $\geq$ three dressing changes				

Acceptable discharge score is considered achieved at  $\geq 14$  provided that no individual score is 0. Any score  $<$ 14 would result in either re-evaluation after achieving improvement in the suboptimal parameter (hence the multiple “post” columns), or physician evaluation

*Reprinted with permission, Cleveland Clinic Foundation © 2022. All rights reserved.*