710 1.23 [0.81,1.88   997 0.85 [0.44,1.63   73 1.35 [0.62,2.95	-	⊣ 0.20
0.85 [0.44,1.63	-	→ 0.20
•	3] [	
73 1.35 [0.62 2 9		6
73 1.35 [0.62 2 9		
10 100 [0102,2100	5] — —	0.74
26 0.97 [0.52,1.80	0]	-
009 1.20 [0.72,2.02	2] ⊢∎	
1.30 [0.66,2.56	6] — —	0.96
1.00 [0.59,1.70	0]	4
1.17 [0.61,2.26	6] — —	
-	-	
563 0.87 [0.49,1.54	4]	0.98
145 1.31 [0.84,2.04	4] ⊢-∎-	-
-	-	
15 0.58 [0.28,1.20	0] ←	0.22
346 1.54 [0.95,2.50	0]	<b> </b>
	-	
1.20 [0.77, 1.85	5] 🗕 🗖 🗖	⊣ 0.49
0.99 [0.56, 1.76	6] — —	4
-	- T	
023 1.11 [0.76,1.64	4] 🗕 🗖	0.98
1.13 [0.47,2.71	1]	i
528 1.22 [0.71,2.10	0] 🛏 💻	- 0.82
180 1.00 [0.63,1.59	9] — —	
		→ 0.44
-	6] 🗕 🛏 🗖 🔤 🛶	
1		28 1.16 [0.65,2.06]

Figure S7, Supplemental Digital Content 11. Subgroup analyses of regular use of PPIs and risk of external cause mortality.

Estimated effects were based on the fully adjusted model.

Abbreviation: BMI, body mass index; NSAID, non-steroidal anti-inflammatory drug; PPI, proton pump inhibitors; HR, hazard ratio; CI, confidence interval.