**Table S3.** Baseline characteristics of patients diagnosed with Celiac disease <18 years of age and sibling comparators

Characteristic	Celiac disease	Siblings
	(n=6,703)	(n=9,089)
Females, no. (%)	4 262 (63.6%)	4 379 (48.2%)
Males, no (%)	2 441 (36.4%)	4 710 (51.8%)
Age		
Mean (SD)	8.6 (4.6)	8.6 (4.8)
Median (IQR)	8.5 (4.6-12.4)	8.4 (4.7-12.5)
Range, min-max	0.0-18.0	0.0-18.0
Categories, no. (%)		
<5y	1 862 (27.8%)	2 473 (27.2%)
5y - <10y	2 180 (32.5%)	2 999 (33.0%)
10y - <15y	1 951 (29.1%)	2 550 (28.1%)
15y - <18y	710 (10.6%)	1 067 (11.7%)
Country of birth, no (%)		
Nordic country	6 570 (98.0%)	8 791 (96.7%)
Other	132 (2.0%)	297 (3.3%)
Missing	1 (0.0%)	1 (0.0%)
Start year of follow-up		
2004-2009	3 537 (52.8%)	4 770 (52.5%)
2010-2017	3 166 (47.2%)	4 319 (47.5%)
Disease history ever before start of follow-up, no. (%)		
Type 1 diabetes	465 (6.9%)	81 (0.9%)
Autoimmune thyroid disease	69 (1.0%)	19 (0.2%)
Inflammatory bowel disease	64 (1.0%)	17 (0.2%)
Systemic lupus erythematosus	0	0
Follow-up, years		
Mean (SD)	7.4 (3.7)	7.0 (3.8)
Median (IQR)	7.1 (4.7-9.9)	6.8 (4.2-9.6)
Range, min-max	0.0-16.0	0.0-15.9