

Practice Guidelines for Obstetrical Anesthesia: Bibliography
American Society of Anesthesiologists

1. Abboud TK, Afrasiabi A, Zhu J, Mantilla M, Reyes A, D'Onofrio L, Khoo N, Mosaad P, Richardson M, Kalra M, et al.: Epidural morphine or butorphanol augments bupivacaine analgesia during labor. *Reg Anesth* 14:115-120, 1989
2. Abboud TK, Artal R, Henriksen EH, Earl S, Kammula RK: Effects of spinal anesthesia on maternal circulating catecholamines. *Am J Obstet Gynecol* 142:252-254, 1982
3. Abboud TK, D'Onofrio L, Reyes A, Mosaad P, Zhu J, Mantilla M, Gangolly J, Crowell D, Cheung M, Afrasiabi A, et al.: Isoflurane or halothane for cesarean section: comparative maternal and neonatal effects. *Acta Anaesthesiol Scand* 33:578-581, 1989
4. Abboud TK, Khoo SS, Miller F, Doan T, Henriksen EH: Maternal, fetal, and neonatal responses after epidural anesthesia with bupivacaine, 2-chloroprocaine, or lidocaine. *Anesth Analg* 61:638-644, 1982
5. Abboud TK, Kim KC, Noueihed R, Kuhnert BR, DerMardirossian N, Moumdjian J, Sarkis F, Nagappala S: Epidural bupivacaine, chloroprocaine, or lidocaine for cesarean section--maternal and neonatal effects. *Anesth Analg* 62:914-919, 1983
6. Abboud TK, Kim SH, Henriksen EH, Chen T, Eisenman R, Levinson G, Shnider SM: Comparative maternal and neonatal effects of halothane and enflurane for cesarean section. *Acta Anaesthesiol Scand* 29:663-668, 1985
7. Abboud TK, Nagappala S, Murakawa K, David S, et al.: Comparison of the effects of general and regional anesthesia for cesarean section on neonatal neurologic and adaptive capacity scores. *Anesth Analg* 64:996-1000, 1985
8. Abboud TK, Raya J, Noueihed R, Daniel J: Intrathecal morphine for relief of labor pain in a parturient with severe pulmonary hypertension. *Anesthesiology* 59:477-479, 1983
9. Abboud TK, Zhu J, Reyes A, Miller H, Steffens Z, Afrasiabi K, Afrasiabi A, Sherman G, Emershad B: Effect of subarachnoid morphine on the incidence of spinal headache. *Reg Anesth* 17:34-36, 1992
10. Abboud TK, Zhu J, Richardson M, Peres da Silva E, Donovan M: Desflurane: a new volatile anesthetic for cesarean section. Maternal and neonatal effects. *Acta Anaesthesiol Scand* 39:723-726, 1995
11. Abboud TK, Zhu J, Richardson M, Peres da Silva E, Donovan M: Intravenous propofol vs thiamylal-isoflurane for caesarean section, comparative maternal and neonatal effects. *Acta Anaesthesiol Scand* 39:205-209, 1995
12. Abboud TK, Zhu J, Sharp R, LaGrange C, Rosa C, Kassells B: The efficacy of intrathecal injection of sufentanil using a microspinal catheter for labor analgesia. *Acta Anaesthesiol Scand* 40:210-215, 1996
13. Abouleisch EI: Postpartum tubal ligation requires more bupivacaine for spinal anesthesia than does cesarean section. *Anesth Analg* 65:897-900, 1986
14. Abouleish A, Abouleish E, Camann W: Combined spinal-epidural analgesia in advanced labour. *Can J Anaesth* 41:575-578, 1994
15. Abouleish E, Abboud T, Lechevalier T, Zhu J, Chalian A, Alford K: Rocuronium (Org 9426) for caesarean section [see comments]. *Br J Anaesth* 73:336-341, 1994
16. Abouleish E, Rawal N, Fallon K, Hernandez D: Combined intrathecal morphine and bupivacaine for cesarean section. *Anesth Analg* 67:370-374, 1988
17. Abouleish E, Rawal N, Rashad MN: The addition of 0.2 mg subarachnoid morphine to hyperbaric bupivacaine for cesarean delivery: a prospective study of 856 cases. *Reg Anesth* 16:137-140, 1991
18. Abouleish E, Rawal N, Shaw J, Lorenz T, Rashad MN: Intrathecal morphine 0.2 mg versus epidural bupivacaine 0.125% or their combination: effects on parturients. *Anesthesiology* 74:711-716, 1991
19. Abouleish E, Rawal N, Tobon Randall B, Rivera Weiss M, Meyer B, Wu A, Rashad MN: A clinical and laboratory study to compare the addition of 0.2 mg of morphine, 0.2 mg of epinephrine, or their combination to hyperbaric bupivacaine for spinal anesthesia in cesarean section [see comments]. *Anesth Analg* 77:457-462, 1993
20. Abouleish E: Apnoea associated with the intrathecal administration of morphine in obstetrics. A case report. *Br J Anaesth* 60:592-594, 1988
21. Abouleish E: Caudal analgesia for quadruplet delivery. *Anesth Analg* 55:61-66, 1976
22. Abouleish E: Foetal bradycardia during caudal analgesia: A discussion of possible causative factors. *Br J Anaesth* 48:481-484, 1976
23. Ackerman WE, Juneja MM, Knapp RK: Maternal paraparesis after epidural anesthesia and cesarean section. *South Med J* 83:695-697, 1990
24. Adams AP, Morgan M, Jones BC, McCormick PW: A case of massive aspiration of gastric contents during obstetric anaesthesia. Treatment by tracheostomy and prolonged intermittent positive pressure ventilation. *Br J Anaesth* 41:176-183, 1969
25. Alahuhta S, Rasanen J, Jouppila R, Jouppila P, Kangas Saarela T, Hollmen AI: Uteroplacental and fetal haemodynamics during extradural anaesthesia for caesarean section. *Br J Anaesth* 66:319-323, 1991
26. Ammar T, Beilin Y, Bernstein HH: Successful regional anesthesia for a woman with a single ventricle presenting for labor and delivery. *J Cardiothorac Vasc Anesth* 10:640-642, 1996

27. Andrews WW, Ramin SM, Maberry MC, Shearer V, Black S, Wallace DH: Effect of type of anesthesia on blood loss at elective repeat cesarean section. *Am J Perinatol* 9:197-200, 1992
28. Angel JL, Knuppel RA, Lake M. Sinusoidal fetal heart rate pattern associated with intravenous butorphanol administration: a case report. *Am J Obstet Gynecol* 149:465-467, 1984
29. Anstey JT, Sheldon GW, Blythe JG: Infectious morbidity after primary cesarean sections in a private institution. *Am J Obstet Gynecol* 136:205-210, 1980
30. Antal M, Benko M: Introduction of anaesthesia in caesarean section by using ketamine and fazadinium. *Acta Chir Hung* 27:19-25, 1986
31. Antoine C, Young BK, Silverman F, Greco MA, Alvarez SP. Sinusoidal fetal heart rate pattern with vasa previa in twin pregnancy. *J Reprod Med* 27:295-300, 1982
32. Antoine C, Young BK: Fetal lactic acidosis with epidural anesthesia. *Am J Obstet Gynecol* 142:55-59, 1982
33. Arkoosh VA, Sharkey SJ, Jr., Norris MC, Isaacson W, Honet JE, Leighton BL: Subarachnoid labor analgesia. Fentanyl and morphine versus sufentanil and morphine. *Reg Anesth* 19:243-246, 1994
34. Asling JH, Schnider SM, Margolis AJ, Wilkinson GL, Way EL. Paracervical block anesthesia in obstetrics. *Am J Obstet Gynecol* 107:626-634, 1970
35. Bach V, Carl P, Ravlo O, Crawford ME, et al.: A randomized comparison between midazolam and thiopental for elective cesarean section anesthesia: III. Placental transfer and elimination in neonates. *Anesth Analg* 68:238-242, 1989
36. Bader AM, Datta S, Arthur GR, Benvenuti E, Courtney M, Hauch M: Maternal and fetal catecholamines and uterine Incision-to-delivery interval during elective cesarean. *Obstet Gynecol* 75:600-603, 1990
37. Baheti DK: Epidural analgesia with left lateral tilt for Caesarean section. *Anaesthesia* 30:396-401, 1975
38. Bajorek J, Kornacki Z, Ronin Walknowska E: Influence of the method of anesthesia for cesarean section on the clinical state of the newborn. *Mater Med Pol* 4:28-31, 1972
39. Baraka A, Louis F, Noueihid R, Diab M, Dabbous A, Sibai A: Awareness following different techniques of general anaesthesia for caesarean section. *Br J Anaesth* 62:645-648, 1989
40. Baraka A, Maktabi M, Noueihid R: Epidural meperidine-bupivacaine for obstetric analgesia. *Anesth Analg* 61:652-656, 1982
41. Baraka A, Noueihid R, Hajj S: Intrathecal injection of morphine for obstetric analgesia. *Anesthesiology* 54:136-140, 1981
42. Baraka A, O'Brien M, Aslanian E, Saade R: Propanidid versus thiopentone for induction of general anaesthesia in elective Caesarean section. *Br J Anaesth* 43:609-612, 1971
43. Barrett JM, Boehm FH: Fetal heart rate responses to meperidine alone and in combination with propiomazine. *South Med J* 76:1480-1483, 1983
44. Barton ACH, Spielman FJ, Onder R, Mayer DC: Timing of postpartum sterilization: Complications and cost-savings of early tubal ligations. *Anesthesiology* 85(A896):3A, 1996
45. Baskett TF, Koh KS. Sinusoidal fetal heart pattern: a sign of fetal hypoxia. *Obstet Gynecol* 44(3):379-382, 1979
46. Baumann H, Alon E, Atanassoff P, Pasch T, Huch A, Huch R: Effect of epidural anesthesia for cesarean delivery on maternal femoral arterial and venous, uteroplacental, and umbilical blood flow velocities and waveforms. *Obstet Gynecol* 75:194-198, 1990
47. Belfrage P, Irestedt L, Raabe N, Arn'er S: General anaesthesia or lumbar epidural block for caesarean section? Effects on the foetal heart rate. *Acta Anaesthesiol Scand* 21:67-70, 1977
48. Benedetti TJ, Cotton DB, Read JC, Miller FC. Hemodynamic observations in severe pre-eclampsia with a flow-directed pulmonary artery catheter. *Am J Obstet Gynecol* 136:465-470, 1980
49. Benson RC, Berendes H, Weiss W: Fetal compromise during elective cesarean section. II. A report from the Collaborative Project. *Am J Obstet Gynecol* 105:579-588, 1969
50. Bernstein K, Gisselsson L, et al.: Influence of two different anaesthetic agents on the newborn and the correlation between foetal oxygenation and induction-delivery time in elective caesarean section. *Acta Anaesthesiol Scand* 29:157-160, 1985
51. Berryhill RE, Benumof JL, Saidman LJ, Smith PC, Plumer MH: Anesthetic management of emergency cesarean section in a patient with epidermolysis bullosa dystrophica polydysplastica. *Anesth Analg* 57:281-283, 1978
52. Blanche R, Eisenach JC, Tuttle R, Dewan DM: Previous wet tap does not reduce success rate of labor epidural analgesia. *Anesth Analg* 79:291-294, 1994
53. Blankenbaker WL, DiFazio CA, Berry FA, Jr. Lidocaine and its metabolites in the newborn. *Anesthesiology* 42:325-330, 1975
54. Bobrowski RA, Bottoms SF: Underappreciated risks of the elderly multipara. *Am J Obstet Gynecol* 172:1764-7; 1995
55. Boehm FH, Woodruff LF, Jr., Growdon JH, Jr. The effect of lumbar epidural anesthesia on fetal heart rate baseline variability. *Anesth Analg* 54:779-782, 1975

56. Bofill JA, Vincent RD, Road EI, Martin RW, Norman PF, Werham CF, Morrison JC: Nulliparous active labor, epidural analgesia, and cesarean delivery for dystocia. *Am J Obstet Gyn* 177:1465-1470, 1997
57. Bonnardot JP, Maillet M, Colau JC, Millot F, Deligne P: Maternal and fetal concentration of morphine after intrathecal administration during labour. *Br J Anaesth* 54:487-489, 1982
58. Boyle R: Caesarean section anaesthesia and the Apgar score. *Aust N Z J Obstet Gynaecol* 33:282-284, 1993
59. Boys JE: Failed intubation in obstetric anaesthesia. A case report. *Br J Anaesth* 55:187-188, 1983
60. Bradford EM, Moir DD: Anaesthesia for Caesarean section: a comparison of thiopentone and propanidid. *Br J Anaesth* 41:274, 1969
61. Breen TW, Janzen JA: Epidural fentanyl and caesarean section: when should fentanyl be given? [see comments]. *Can J Anaesth* 39:317-322, 1992
62. Brian JE, Jr., Seifen AB, Clark RB, Robertson DM, Quirk JG: Aortic stenosis, cesarean delivery, and epidural anesthesia [see comments]. *J Clin Anesth* 5:154-157, 1993
63. Brizgys RV, Shnider SM: Hyperbaric intrathecal morphine analgesia during labor in a patient with Wolff-Parkinson-White syndrome. *Obstet Gynecol* 64:44S-46S, 1984
64. Brownridge P: Spinal anaesthesia revisited: an evaluation of subarachnoid block in obstetrics. *Anaesth Intensive Care* 12:334-342, 1984
65. Burns AM, Dorje P, Lawes EG, Nielsen MS: Anaesthetic management of caesarean section for a mother with pre-eclampsia, the Klippel-Feil syndrome and congenital hydrocephalus. *Br J Anaesth* 61:350-354, 1988
66. Caldwell LE, Rosen MA, Shnider SM: Subarachnoid morphine and fentanyl for labor analgesia. Efficacy and adverse effects. *Reg Anesth* 19:2-8, 1994
67. Camann WR, Denney RA, Holby ED, Datta S: A comparison of intrathecal, epidural, and intravenous sufentanil for labor analgesia. *Anesthesiology* 77:884-887, 1992
68. Camann WR, Hortvet LA, Hughes N, Bader AM, Datta S: Maternal temperature regulation during extradural analgesia for labour. *Br J Anaesth* 67:565-568, 1991
69. Camann WR, Minzter BH, Denney RA, Datta S: Intrathecal sufentanil for labor analgesia. Effects of added epinephrine. *Anesthesiology* 78:870-4; 1993
70. Campbell DC, Banner R, Crone LA, GoreHickman W, Yip RW: Addition of epinephrine to intrathecal bupivacaine and sufentanil for ambulatory labor analgesia. *Anesthesiology* 86:525-531, 1997
71. Campbell DC, Camann WR, Datta S: Combined spinal-epidural for labor analgesia: Comparison of intrathecal sufentanil vs bupivacaine vs sufentanil plus bupivacaine. *Anesthesiology* 81(3A):A1145, 1994
72. Campbell DC, Camann WR, Datta S: The addition of bupivacaine to intrathecal sufentanil for labor analgesia. *Anesth Analg* 81:305-309, 1995
73. Capogna G, Celleno D, Costantino P, Muratori F, Sebastiani M, Baldassini M: Alkalinization improves the quality of lidocaine-fentanyl epidural anaesthesia for caesarean section. *Can J Anaesth* 40:1425-1430, 1993
74. Capogna G, Celleno D, Varrassi G, Emanuelli M, Sebastiani M, Muratori F, Cipriani G, Tomasetti M: Epidural mepivacaine for cesarean section: effects of a pH-adjusted solution. *J Clin Anesth* 3:211-4; 1991
75. Caritis SN, Abouleish E, Edelstone DI, Mueller Heubach E: Fetal acid-base state following spinal or epidural anesthesia for cesarean section. *Obstet Gynecol* 56:610-615, 1980
76. Carlsson BM, Johansson M, Westin B: Fetal heart rate pattern before and after paracervical anesthesia. A prospective study. *Acta Obstet Gynecol Scand Suppl* 66:391-395, 1987
77. Carlsson C, Nybell-Lindahl G, Ingemarsson I: Extradural block in patients who have previously undergone caesarean section. *Br J Anaesth* 52:827-829, 1980
78. Carton EG, McDonald N, McCarthy JR: Intrathecal morphine in labour--efficacy and side effects. *Ir J Med Sci* 156:323-327, 1987
79. Cascio M, Heath G: Meningitis following a combined spinal-epidural technique in a labouring term parturient. *Can J Anaesth* 43:399-402, 1996
80. Cascio M, Pygenon B, Bennett C, Ramanathan S: Labour analgesia with intrathecal fentanyl decreases maternal stress. *Can J Anaesth* 44(6):605-609, 1997
81. Celleno D, Capogna G, Emanuelli M, Varrassi G, Muratori F, Costantino P, Sebastiani M: Which induction drug for cesarean section? A comparison of thiopental sodium, propofol, and midazolam. *J Clin Anesth* 5:284-288, 1993
82. Celleno D, Capogna G, Sebastiani M, Costantino P, Muratori F, Cipriani G, Emanuelli M: Epidural analgesia during and after cesarean delivery. Comparison of five opioids. *Reg Anesth* 16:79-83, 1991
83. Celleno D, Capogna G, Tomasetti M, Costantino P, Di Feo G, Nisini R: Neurobehavioural effects of propofol on the neonate following elective caesarean section. *Br J Anaesth* 62:649-654, 1989
84. Celleno D, Capogna G: Epidural fentanyl plus bupivacaine 0.125 per cent for labour: analgesic effects. *Can J Anaesth* 35:375-378, 1988

85. Chadwick HS, Posner K, Caplan RA, Ward RJ, Cheney FW: A comparison of obstetric and nonobstetric anesthesia malpractice claims. *Anesthesiology* 74:242-249, 1991
86. Chan AS, Ananthanarayan C, Rolbin SH: Alternating nitroglycerin and syntocinon to facilitate uterine exploration and removal of an adherent placenta. *Can J Anaesth* 42:335-337, 1995
87. Chestnut DH, Noe AL: Effect of anesthesia for primary cesarean section on postoperative infectious morbidity. *Obstet Gynecol* 68:667-670, 1986
88. Chestnut DH, Owen CL, Bates JN, Ostman LG, Choi WW, Geiger MW: Continuous infusion epidural analgesia during labor: a randomized, double-blind comparison of 0.0625% bupivacaine/0.0002% fentanyl versus 0.125% bupivacaine. *Anesthesiology* 68:754-759, 1988
89. Chestnut DH, Owen CL, Brown CK, Vandewalker GE, Weiner CP: Does labor affect the variability of maternal heart rate during induction of epidural anesthesia? *Anesthesiology* 68:622-625, 1988
90. Chestnut DH: Effect of anesthesia for repeat cesarean section on postoperative infectious morbidity. *Obstet Gynecol* 66:199-202, 1985
91. Chung CJ, Bae SH, Chao KY, Chin YJ: Spinal anaesthesia with 0.25% hyperbaric bupivacaine for Caesarean section: Effects of volume. *Br J Anaesth* 77:145-149, 1996
92. Cibils LA: Clinical significance of fetal heart rate patterns during labor. VI. Early decelerations. *Am J Obstet Gynecol* 136:392-398, 1980
93. Cigarini I, Kaba A, Bonnet F, Brohon E, Dutz F, Damas F, Hans P: Epidural clonidine combined with bupivacaine for analgesia in labor. Effects on mother and neonate. *Reg Anesth* 20:113-120, 1995
94. Clark A, Carr D, Loyd G, Cook V, Spinnato J: The influence of epidural anesthesia on cesarean delivery rates: a randomized prospective clinical trial. *Am J Obstet Gynecol* 179:1527-1533, 1998
95. Clark SL, Greenspoon JS, Aldahl D, Phelan JP: Severe preeclampsia with persistent oliguria: management of hemodynamic subsets. *Am J Obstet Gynecol* 154:490-494, 1986
96. Clarke VT, Smiley RM, Finster M: Uterine hyperactivity after intrathecal injection of fentanyl for analgesia during labor: a cause of fetal bradycardia? [letter]. *Anesthesiology* 81:1083, 1994
97. Cohen S, Amar D, Nagashima H: Extradural block in obstetric patients: review of experience with gravity administration. *Acta Anaesthesiol Scand* 35:676-679, 1991
98. Cohen SE, Cherry CM, Holbrook RH, Jr., el Sayed YY, Gibson RN, Jaffe RA: Intrathecal sufentanil for labor analgesia--sensory changes, side effects, and fetal heart rate changes. *Anesth Analg* 77:1155-1160, 1993
99. Coleman AJ, Downing JW: Enflurane anesthesia for cesarean section. *Anesthesiology* 43:354-357, 1975
100. Collea JV, Rabin SC, Weghorst GR, Quilligan EJ: The randomized management of term frank breech presentation: vaginal delivery vs. cesarean section. *Am J Obstet Gynecol* 131:186-195, 1978
101. Collins KM, Bevan DR, Beard RW: Fluid loading to reduce abnormalities of fetal heart rate and maternal hypotension during epidural analgesia in labour. *Br Med J* 2:1460-1461, 1978
102. Collis RE, Baxandall ML, Srikantharajah ID, Edge G, Kadim MY, Morgan BM: Combined spinal epidural analgesia with ability to walk throughout labour. *Lancet* 341:767-768, 1993
103. Collis RE, Davies DW, Aveling W: Randomised comparison of combined spinal-epidural and standard epidural analgesia in labour [see comments]. *Lancet* 345:1413-1416, 1995
104. Combs CA, Laros RK, Jr. Prolonged third stage of labor: morbidity and risk factors. *Obstet Gynecol* 77:863-867, 1991
105. Combs CA, Murphy EL, Laros RK, Jr. Factors associated with hemorrhage in cesarean deliveries. *Obstet Gynecol* 77:77-82, 1991
106. Cooperstein M: The effects of fetal monitoring on oxytocin inductions. *J Am Osteopath Assoc* 74:654-660, 1975
107. Copel JA, Harrison D, Whittemore R, Hobbins JC: Intrathecal morphine analgesia for vaginal delivery in a woman with a single ventricle. A case report. *J Reprod Med* 31:274-276, 1986
108. Cosmi EV, Marx GF: Acid-base status of the fetus and clinical condition of the newborn following cesarean section. *Am J Obstet Gynecol* 102:378-382, 1968
109. Courtney MA, Bader AM, Hartwell B, Hauch M, Grennan MJ, Datta S: Perioperative analgesia with subarachnoid sufentanil administration. *Reg Anesth* 17:274-278, 1992
110. Craft JB, Levinson G, Shnider SM: Anaesthetic considerations in caesarean section for quadruplets. *Can Anaesth Soc J* 25:236-239, 1978
111. Crawford JS, Davies P, Lewis M: Some aspects of epidural block provided for elective caesarean section. *Anaesthesia* 41:1039-1046, 1986
112. Crawford JS, James FM3, Davies P, Crawley M: A further study of general anaesthesia for Caesarean section. *Br J Anaesth* 48:661-667, 1976
113. Crawford JS: A comparison of spinal analgesia and general anesthesia for elective cesarean section. Interim report. *Am J Obstet Gynecol* 94:858-860, 1966

114. Crosby ET: Obstetrical anaesthesia for patients with the syndrome of haemolysis, elevated liver enzymes and low platelets [see comments]. *Can J Anaesth* 38:227-233, 1991
115. Cruikshank DP, Laube DW, De-Backer LJ: Intraperitoneal lidocaine anesthesia for postpartum tubal ligation. *Obstet Gynecol* 42(1):127-130, 1973
116. Dailey PA, Fisher DM, Scnider SM, Baysinger CL, Shinohara Y, Miller RD, Abboud TK, Kim KC: Pharmacokinetics, placental transfer, and neonatal effects of vecuronium and pancuronium administered during cesarean section. *Anesthesiology* 60:569-574, 1984
117. Dailland P, Cockshott ID, Lirzin JD, et al.: Intravenous propofol during cesarean section: placental transfer, concentrations in breast milk, and neonatal effects. A preliminary study [see comments]. *Anesthesiology* 71:827-834, 1989
118. D'Angelo R, Anderson MT, Philip J, Eisenach JC: Intrathecal sufentanil compared to epidural bupivacaine for labor analgesia. *Anesthesiology* 80:1209-1215, 1994
119. D'Angelo, Gerancher JC, Eisenach JC, Raphael BL: Epidural fentanyl produces labor analgesia by a spinal mechanism. *Anesthesiology* 88:1519-1523, 1998
120. Daniels JC, Hehre FW: Anesthetic considerations for complicated obstetrics. I. A retrospective study of 527 twin deliveries. *Anesth Analg* 46:527-539, 1967
121. Darby S, Hunter DJ: Extradural analgesia in labour when the breech presents. *Br J Obstet Gynaecol* 83:35-38, 1976
122. Das Gupta D: Propanidid versus thiopentone for induction of general anaesthesia in elective Caesarean section. *Br J Anaesth* 44:290, 1972
123. Datta S, Alper MH, Ostheimer GW, Brown WU, Jr., Weiss JB: Effects of maternal position on epidural anesthesia for cesarean section, acid-base status, and bupivacaine concentrations at delivery. *Anesthesiology* 50:205-209, 1979
124. Datta S, Brown WU, Jr. Acid-base status in diabetic mothers and their infants following general or spinal anesthesia for cesarean section. *Anesthesiology* 47:272-276, 1977
125. Datta S, Camann W, Bader A, Vanderburgh L: Clinical effects and maternal and fetal plasma concentrations of epidural ropivacaine versus bupivacaine for cesarean section. *Anesthesiology* 82:1346-1352, 1995
126. Datta S, Houle GL, Fox GS: Concentration of lidocaine hydrochloride in newborn gastric fluid after elective caesarean section and vaginal delivery with epidural analgesia. *Can Anaesth Soc J* 22:79-83, 1975
127. Datta S, Ostheimer GW, Weiss JB, Brown WU, Jr., Alper MH: Neonatal effect of prolonged anesthetic induction for cesarean section. *Obstet Gynecol* 58:331-335, 1981
128. Davies SJ, Paech MJ, Welch H, Evans SF, Pavay TJJ: Maternal experience during epidural or combined spinal-epidural anesthesia for cesarean section: A prospective, randomized trial. *Anesth Analg* 85:607-613, 1997
129. de Leon Casasola OA, Lema MJ, Emrich L, Syed N: Continuous 2-chloroprocaine infusion versus intermittent bolus injections of bupivacaine or 2-chloroprocaine for epidural anesthesia in cesarean delivery. *Reg Anesth* 16:154-160, 1991
130. de Vries JI, Vellenga E, Aarnoudse JG: Plasma beta-thromboglobulin in normal pregnancy and pregnancy-induced hypertension. *Eur J Obstet Gynecol Reprod Biol* 14:209-216, 1983
131. Deckardt R, Fembacher PM, Schneider KT, Graeff H: Maternal arterial oxygen saturation during labor and delivery: pain-dependent alterations and effects on the newborn. *Obstet Gynecol* 70:21-25, 1987
132. Department of Health, Welsh Office, Scottish Office Department of Health and Social Services, Northern Ireland: Report on confidential enquiries into maternal deaths in the United Kingdom 1985-1987. Chapter 9, 1992
133. Department of Health, Welsh Office, Scottish Office Department of Health and Social Services, Northern Ireland: Report on confidential enquiries into maternal deaths in the United Kingdom 1988-1990. Chapter 9, 1995
134. Department of Health, Welsh Office, Scottish Office Department of Health and Social Services, Northern Ireland: Report on confidential enquiries into maternal deaths in the United Kingdom 1991-1993. Chapter 9, 1998
135. DeSimone CA, Leighton BL, Norris MC: Spinal anesthesia for cesarean delivery. A comparison of two doses of hyperbaric bupivacaine [see comments]. *Reg Anesth* 20:90-94, 1995
136. DeSimone CA, Norris MC, Leighton BL: Intravenous nitroglycerin aids manual extraction of a retained placenta [letter]. *Anesthesiology* 73:787, 1990
137. Desprats R, Dumas JC, Giroux M, Campistron G, Faure F, Teixeira MG, Grandjean H, Houin G, Pontonnier G: Maternal and umbilical cord concentrations of fentanyl after epidural analgesia for cesarean section. *Eur J Obstet Gynecol Reprod Biol* 42:1991
138. Desprats R, Mandry J, Grandjean H, Amar B, Pontonnier G, Lareng L: [Peridural analgesia during labor: comparative study of a fentanyl-marcaine combination and marcaine alone]. *J Gynecol Obstet Biol Reprod Paris* 12:901-905, 1983
139. Devcic A, Sprung J, Patel S, Kettler R, Maitra DA: PDPH in obstetric anesthesia: comparison of 24-gauge Sprotte and 25-gauge Quincke needles and effect of subarachnoid administration of fentanyl [see comments]. *Reg Anesth* 18:222-225, 1993
140. Dick W, Traub E, Kraus H, Tollner U, Burghard R, Muck J: General anaesthesia versus epidural anaesthesia for primary caesarean section--a comparative study. *Eur J Anaesthesiol* 9:15-21, 1992

141. Dobb G: Laryngeal oedema complicating obstetric anaesthesia [letter]. *Anaesthesia* 33:839-840, 1978
142. Donnai P, Nicholas AD: Epidural analgesia, fetal monitoring and the condition of the baby at birth with breech presentation. *Br J Obstet Gynaecol* 82:360-365, 1975
143. Downing JW, Buley RJ, Brock Utne JG, Houlton PC: Etomidate for induction of anaesthesia at caesarean section: comparison with thiopentone. *Br J Anaesth* 51:135-140, 1979
144. Downing JW, Coleman AJ: Anaesthesia for emergency caesarean section: propanidid versus thiopentone. *S Afr Med J* 47:1369-1372, 1973
145. Downing JW, Houlton PC, Barclay A: Extradural analgesia for caesarean section: a comparison with general anaesthesia. *Br J Anaesth* 51:367-374, 1979
146. Downing JW, Johnson HV, Gonzalez HF, Arney TL, Herman NL, Johnson RF: The pharmacokinetics of epidural lidocaine and bupivacaine during cesarean section. *Anesth Analg* 84:527-532, 1997
147. Downing JW, Mahomed MC, Coleman AJ, Mahomed YH, Jeal DE: Anaesthetic induction for Caesarean section. Althesin versus thiopentone. *Anaesthesia* 29:689-695, 1974
148. Downing JW, Mahomed MC, Jeal DE, Allen PJ: Anaesthesia for Caesarean section with ketamine. *Anaesthesia* 31:883-892, 1976
149. Driver I, Popham P, Glazebrook C, Palmer C: Epidural bupivacaine/fentanyl infusions vs. intermittent top-ups: a retrospective study of the effects on mode of delivery in primiparous women. *Eur J Anaesthesiol* 13:515-520, 1996
150. Druzin ML, Stier E: Maternal platelet count at delivery in patients with idiopathic thrombocytopenic purpura, not related to perioperative complications. *J Am Coll Surg* 179:264-266, 1994
151. Duvaldestin P, Demetriou M, Henzel D, Desmonts JM: The placental transfer of pancuronium and its pharmacokinetics during caesarian section. *Acta Anaesthesiol Scand* 22:327-333, 1978
152. Dwyer R, Fee JP, Moore J: Uptake of halothane and isoflurane by mother and baby during caesarean section. *Br J Anaesth* 74:379-383, 1995
153. Edwards ND, Hartley M, Clyburn P, Harmer M: Epidural pethidine and bupivacaine in labour. *Anaesthesia* 47:435-437, 1992
154. Edwards RD, Hansel NK, Pruessner HT, Barton B: Intrathecal morphine as analgesia for labor pain. *J Am Board Fam Pract* 1:245-250, 1988
155. Edwards RD, Hansel NK, Pruessner HT, Barton B: Intrathecal morphine sulfate for labor pain. *Tex Med* 81:46-48, 1985
156. el Hairy A, Karim A, Dormeval MH: [The "HELLP" syndrome: a severe sign in toxemia. Three new cases]. *J Gynecol Obstet Biol Reprod Paris* 21:227-229, 1992
157. Elliott RD: Continuous infusion epidural analgesia for obstetrics: bupivacaine versus bupivacaine-fentanyl mixture. *Can J Anaesth* 38:303-310, 1991
158. Endler GC, Mariona FG, Sokol RJ, Stevenson LB: Anesthesia-related maternal mortality in Michigan, 1972 to 1984. *Am J Obstet Gynecol* 159:187-193, 1988
159. Enever GR, Noble HA, Kolditz D, Valentine S, Thomas TA: Epidural infusion of diamorphine with bupivacaine in labour. A comparison with fentanyl and bupivacaine. *Anaesthesia* 46:169-173, 1991
160. Epstein H, Waxman A, Gleicher N, Lauersen NH: Meperidine-induced sinusoidal fetal heart rate pattern and reversal with naloxone. *Obstet Gynecol* 59:22S, 1982
161. Epstein RH, Kaplan S, Leighton BL, Norris MC, DeSimone CA: Evaluation of a continuous noninvasive blood pressure monitor in obstetric patients undergoing spinal anesthesia. *J Clin Monit* 5:157-163, 1989
162. Ewen A, McLeod DD, MacLeod DM, Campbell A, Tunstall ME: Continuous infusion epidural analgesia in obstetrics. A comparison of 0.08% and 0.25% bupivacaine. *Anaesthesia* 41:143-147, 1986
163. Fedorkow DM, Stewart TJ, Parboosinh J: Fetal heart rate changes associated with general anesthesia. *Am J Perinatol* 6:287-288, 1989
164. Ferouz F, Norris MC, Arkoosh VA, Leighton BL, Boxer LM, Corba RJ: Baricity, needle direction, and intrathecal sufentanil labor analgesia. *Anesthesiology* 86:592-598, 1997
165. Finfer SR: Management of labour and delivery in patients with intracranial neoplasms. *Br J Anaesth* 67:784-787, 1991
166. Fitzgerald MP, Floro C, Siegel J, Hernandez E: Laboratory findings in hypertensive disorders of pregnancy. *J Natl Med Assoc* 88:794-798, 1996
167. Flamm BL, Dunnett C, Fischermann E, Quilligan EJ: Vaginal delivery following cesarean section: use of oxytocin augmentation and epidural anesthesia with internal tocodynamometer and internal fetal monitoring. *Am J Obstet Gynecol* 148:759-763, 1984
168. Flamm BL, Lim OW, Jones C, Fallon D, Newman LA, Mantis JK: Vaginal birth after cesarean section: results of a multicenter study. *Am J Obstet Gynecol* 158:1079-1084, 1988
169. Flynn RJ, McMurray TJ, Dwyer R, Moore J: Comparison of plasma bupivacaine concentrations during continuous extradural infusion for labour. *Br J Anaesth* 61:382-384, 1988

170. Fogel ST, Shyken JM, Mormol J, Leighton BD, Smeltzer JS: Epidural anesthesia does not cause cesarean delivery. *Anesthesiology* 87(3A):A911, 1997
171. Fong J, Gurewitsch ED, Press RA, Gomillion MC, Volpe L: Prevention of maternal hypotension by epidural administration of ephedrine sulfate during lumbar epidural anesthesia for cesarean section. *Am J Obstet Gynecol* 175:985-990, 1996
172. Fox GS, Houle GL: Transmission of lidocaine hydrochloride across the placenta during Caesarian section. *Can Anaesth Soc J* 16:135-143, 1969
173. Fox GS, Smith JB, Namba Y, Johnson RC: Anesthesia for cesarean section: further studies. *Am J Obstet Gynecol* 133:15-19, 1979
174. Freedman MA, Wilds PL, George WM: Grotesque obesity: a serious complication of labor and delivery. *South Med J* 65:732-736, 1972
175. Gabert HA, Stenchever MA: Electronic fetal monitoring in association with paracervical blocks. *Am J Obstet Gynecol* 116:1143-1146, 1973
176. Gaffud MP, Bansal P, Lawton C, Velasquez N, Watson WA: Surgical analgesia for cesarean delivery with epidural bupivacaine and fentanyl. *Anesthesiology* 65:331-334, 1986
177. Gajraj NM, Wallace DH, Pace NA: Supraventricular tachycardia in a parturient under spinal anesthesia. *Reg Anesth* 18:261-263, 1993
178. Galbert MW, Gardner AE: Use of halothane in a balanced technic for cesarean section. *Anesth Analg* 51:701-704, 1972
179. Gale R, Slater PE, Zalkinder Luboshitz I: Neonatal advantage of epidural anesthesia in elective and emergency cesarean sections: a report of 531 cases. *Eur J Obstet Gynecol Reprod Biol* 23:1986
180. Gambling DR, Howell P, Huber C, Kozak S: Epidural butorphanol does not reduce side effects from epidural morphine after cesarean birth [see comments]. *Anesth Analg* 78:1099-1104, 1994
181. Gataure PS, Hughes JA: The laryngeal mask airway in obstetrical anaesthesia. *Can J Anaesth* 42:130-133, 1995
182. Gepts E, Heytens L, Camu F: Pharmacokinetics and placental transfer of intravenous and epidural alfentanil in parturient women. *Anesth Analg* 65:1155-1160, 1986
183. Gershon RY: Surgical anaesthesia for Caesarean section with a subdural catheter. *Can J Anaesth* 43:1068-1071, 1996
184. Ghosh AK, Tipton RH: Early postpartum tubal ligation under epidural analgesia. *Br J Obstet Gynaecol* 83:731-732, 1976
185. Giannina G, Guzman ER, Yu-Ling L, Lake MF, Cernadas M, Vintzileos M: Comparison of the effects of meperidine and nalbuphine on intrapartum fetal heart rate tracings. *Obstet Gynecol* 86:441-445, 1995
186. Gilstrap LC, III, Hauth JC, Hankins GD, Patterson AR: Effect of type of anesthesia on blood loss at cesarean section. *Obstet Gynecol* 69:t 1328-32, 1987
187. Gin T, Gregory MA, Chan K, Buckley T, Oh TE: Pharmacokinetics of propofol in women undergoing elective caesarean section. *Br J Anaesth* 64:148-153, 1990
188. Gin T, Gregory MA, Chan K, Oh TE: Maternal and fetal levels of propofol at caesarean section. *Anaesth Intensive Care* 18:180-184, 1990
189. Gin T, O'Meara ME, Kan AF, Leung RK, Tan P, Yau G: Plasma catecholamines and neonatal condition after induction of anaesthesia with propofol or thiopentone at caesarean section. *Br J Anaesth* 70:311-316, 1993
190. Gin T, Yau G, Chan K, Gregory MA, Oh TE: Disposition of propofol infusions for caesarean section. *Can J Anaesth* 38:31-36, 1991
191. Gin T, Yau G, Jong W, Tan P, Leung RK, Chan K: Disposition of propofol at caesarean section and in the postpartum period. *Br J Anaesth* 67:49-53, 1991
192. Giuffrida JG, Bizzarri DV, Masi R, Bondoc R: Continuous procaine spinal anesthesia for cesarean section. *Anesth Analg* 51:117-124, 1972
193. Glosten B, Dailey PA, Preston PG, Shnider SM, Ross BK, Rosen MA, Hughes SC: pH-adjusted 2-chloroprocaine for epidural anesthesia in patients undergoing postpartum tubal ligation. *Anesthesiology* 68:948-950, 1988
194. Godley M, Reddy AR: Use of LMA for awake intubation for caesarean section. *Can J Anaesth* 43:299-302, 1996
195. Goodlin RC: Intrapartum fetal heart rate responses and plethysmographic pulse. *Am J Obstet Gynecol* 110:210-226, 1971
196. Goodman EJ, Dumas SD, Eisenmann UB: The rate of successful reactivation of labor epidural catheters for postpartum tubal ligation surgery. *Anesthesiology* 85(A895):3A, 1996
197. Gordon H: Fetal bradycardia after paracervical block. *N Eng J Med.* 279:910, 1968
198. Gozal D, Gozal Y, Beilin B: Removal of knotted epidural catheters. *Reg Anesth* 21:71-73, 1996
199. Graham C, Goldstein A: Epidural analgesia and cardiac output in severe pre-eclamptics. *Anaesthesia* 35:709-712, 1980
200. Graham CW, Sutton L, Cozen HJ: Directional spinals in obstetric analgesia. *Anaesthesia* 33:192-195, 1978
201. Gran L, Maltau JM: Propanidid (epontol for induction of general anaesthesia in Caesarean section. *Anaesthetist* 20:247-249, 1971

202. Gray C, Richmond MN, Duncan SL: Management of Caesarean delivery in the presence of a large pelvic vascular malformation. *Anaesthesia* 51:961-964, 1996
203. Gray JH, Cudmore DW, Luther ER, Martin TR, Gardner AJ: Sinusoidal fetal heart rate pattern associated with alphaprodine administration. *Obstet Gynecol* 52:678-681, 1978
204. Green SL, Sarubbi FA, Jr. Risk factors associated with post cesarean section febrile morbidity. *Obstet Gynecol* 49:686-690, 1977
205. Greenhalgh CA: Respiratory arrest in a parturient following intrathecal injection of sufentanil and bupivacaine. *Anaesthesia* 51:173-175, 1996
206. Grieco WM, Norris MC, Leighton BL, Arkoosh VA, Huffnagle HJ, Honet JE, Costello D: Intrathecal sufentanil labor analgesia: the effects of adding morphine or epinephrine. *Anesth Analg* 77:1149-1154, 1993
207. Griffin RP, Reynolds F: Extradural anaesthesia for caesarean section: a double-blind comparison of 0.5% ropivacaine with 0.5% bupivacaine. *Br J Anaesth* 74:512-516, 1995
208. Griffin RP, Reynolds F: Maternal hypoxaemia during labour and delivery: the influence of analgesia and effect on neonatal outcome. *Anaesthesia* 50:151-156, 1995
209. Groenendijk R, Trimbos MJ, Wallenburg HCS: Hemodynamic measurements in preeclampsia: preliminary observations. *Am J Obstet Gynecol* 150:232-236, 1984
210. Grundy EM, Zamora AM, Winnie AP: Comparison of spread of epidural anesthesia in pregnant and nonpregnant women. *Anesth Analg* 57:544-546, 1978
211. Gullestad S, Sagen N: Epidural block in twin labour and delivery. *Acta Anaesthesiol Scand* 21:504-508, 1977
212. Gupta R, Singh B, Sachdev L, Gupta S: Neonatal side effects of intrathecal morphine for painless labor. *Indian Pediatr* 28:1321-1323, 1991
213. Hadzic A, Vloka J, Patel N, Birnbach D: Hypertensive crisis after a successful placement of an epidural anesthetic in a hypertensive parturient. Case report. *Reg Anesth* 20:156-158, 1995
214. Hall MH, Halliwell R, Carr Hill R: Concomitant and repeated happenings of complications of the third stage of labour. *Br J Obstet Gynaecol* 92:732-738, 1985
215. Hamilton CL, Cohen SE: High sensory block after intrathecal sufentanil for labor analgesia. *Anesthesiology* 83:1118-1121, 1995
216. Hankins GD, Wendel GD, Jr., Cunningham FG, Leveno KJ: Longitudinal evaluation of hemodynamic changes in eclampsia. *Am J Obstet Gynecol* 150:1506-1512, 1984
217. Haram K, Bakke OM: Diazepam as an induction agent for cesarean section: a clinical and pharmacokinetic study of fetal drug exposure. *Br J Obstet Gynecol* 87:506-512, 1980
218. Harding SA, Collis RE, Morgan BM: Meningitis after combined spinal-extradural anaesthesia in obstetrics [see comments]. *Br J Anaesth* 73:545-547, 1994
219. Harsten A, Gillberg L, Hakansson L, Olsson M: Intrathecal sufentanil compared with epidural bupivacaine analgesia in labour. *Eur J Anaesth* 14:642-645, 1997
220. Haspels AA, Smorenberg Schoorl ME: Fall in blood pressure during spinal anaesthesia for caesarian section. *Trop Geogr Med* 19:347-351, 1967
221. Hatjis CG, Meis PJ: Sinusoidal fetal heart rate pattern associated with butorphanol administration. *Obstet Gynecol* 67:377-380, 1986
222. Hauch MA, Gaiser RR, Hartwell BL, Datta S: Maternal and fetal colloid osmotic pressure following fluid expansion during cesarean section. *Crit Care Med* 23:510-514, 1995
223. Hauch MA, Hartwell BL, Hunt CO, Datta S: Comparative effects of subarachnoid hyperbaric bupivacaine and tetracaine-procaine for cesarean delivery. *Reg Anesth* 15:81-85, 1990
224. Hawkins JL, Hess KR, Kubicek MA, Joyce TH3, Morrow DH: A reevaluation of the association between instrument delivery and epidural analgesia. *Reg Anesth* 20:50-56, 1995
225. Hawkins JL, Johnson TD, Kubicek MA, Skjonsby BS, Morrow DH, Joyce TH, III: Vecuronium for rapid-sequence intubation for cesarean section. *Anesth Analg* 71:185-190, 1990
226. Hawkins JL, Koonin LM, Palmer SK, Gibbs CP: Anesthesia-related deaths during obstetric delivery in the United States, 1979-1990. *Anesthesiology* 86:277-284, 1997
227. Healy TE: Suxamethonium and intra-uterine pressure. *Br J Anaesth* 43:1156-1158, 1971
228. H'ee P, Stampe Sorensen S, Bock JE, Fernandes A, Steentoft A: Intrathecal administration of morphine for the relief of pains in labour and estimation of maternal and fetal plasma concentration of morphine. *Eur J Obstet Gynecol Reprod Biol* 25:1987
229. Heese HdV, Davey DA, Rorke M, Molteno C: Effect of maternal anaesthesia on oxygenation and acid-base status of the newborn infant. *S Afr Med J* 47:1991-1999, 1973
230. Hehre FW, Hook R, Hon EH: Continuous lumbar peridural anesthesia in obstetrics. VI. The fetal effects of transplacental passage of local anesthetic agents. *Anesth Analg* 48:909-913, 1969

231. Helbo Hansen S, Bang U, Garcia RS, Olesen AS, Kjeldsen L: Subarachnoid versus epidural bupivacaine 0.5% for caesarean section. *Acta Anaesthesiol Scand* 32:473-476, 1988
232. Herpolsheimer A, Schretenthaler J: The use of intrapartum intrathecal narcotic analgesia in a community-based hospital. *Obstet Gynecol* 84:931-936, 1994
233. Hingson R, Gould JB, Morelock S, Kaye H, Heeren T, Alpert JJ, Zuckerman B, Day N: Maternal cigarette smoking, psychoactive substance use, and infant Apgar scores. *Am J Obstet Gynecol* 144:959-966, 1982
234. Hjertberg R, Belfrage P, Hagenvik K: Hemodynamic measurements with Swan-Ganz catheter in women with severe proteinuric gestational hypertension (pre-eclampsia). *Acta Obstet Gynecol Scand* 70:193-198, 1991
235. Hodgkinson R, Bhatt M, Kim SS, Grewal G, Marx GF: Neonatal neurobehavioral tests following cesarean section under general and spinal anesthesia. *Am J Obstet Gynecol* 132:670-674, 1978
236. Hodgkinson R, Husain FJ: Caesarean section associated with gross obesity. *Br J Anaesth* 52:919-923, 1980
237. Hoffman CT III, Guzman ER, Richardson MJ, Vintzileos A, Houlihan C, Benito C: Effects of narcotic and non-narcotic continuous epidural anesthesia on intrapartum fetal heart rate tracings as measured by computer analysis. *J Maternal-Fetal Med* 6:200-205, 1997
238. Holdcroft A, Morgan M, Gordon H, Whitwam JG, White Y: Althesin as an induction agent for Caesarean section. *Br J Anaesth* 47:1213-1217, 1975
239. Holdcroft A, Robinson MJ, Gordon H, Whitwam JG: Comparison of effect of two induction doses of methohexitone on infants delivered by elective caesarean section. *Br Med J* 2:472-475, 1974
240. Hollm'en A, Jagerhorn M, Pystynen P: Foetal and maternal acid-base balance during labour and at birth. The effects of general anaesthesia and Caesarean section on the "physiological foetal acidosis" of normal delivery. *Acta Anaesthesiol Scand* 11:341-352, 1967
241. Hollm'en A, Jagerhorn M: The effects of epidural anaesthesia and Caesarean section on foetal and maternal acid-base balance at birth. *Acta Anaesthesiol Scand* 12:114-123, 1968
242. Hollm'en A, Karvonen P, Pihlajaniemi R: Comparison of thiopentone versus althesin for caesarean section. *Ann Chir Gynaecol* 65:392-397, 1976
243. Hollm'en A: Lumbar epidural analgesia as an obstetric analgesic method. *Acta Anaesthesiol Scand Suppl* 37:251-255, 1970
244. Hollm'en AI, Eskelinen P, Tolonen U, Arranto J, Sulg I, Jouppila R, Jouppila P: Effects of anaesthesia for caesarean section on the computerized EEG of the neonate. *Eur J Anaesthesiol* 2:39-51, 1985
245. Hollm'en AI, Jouppila R, Koivisto M, Maatta L, Pihlajaniemi R, Puukka M, Rantakyla P: Neurologic activity of infants following anesthesia for cesarean section. *Anesthesiology* 48:350-356, 1978
246. Honet JE, Arkoosh VA, Norris MC, Huffnagle HJ, Silverman NS, Leighton BL: Comparison among intrathecal fentanyl, meperidine, and sufentanil for labor analgesia. *Anesth Analg* 75:734-739, 1992
247. Hood DD, Dewan DM: Anesthetic and obstetric outcome in morbidly obese parturients. *Anesthesiology* 79:1210-1218, 1993
248. Horta ML, Horta BL: Inhibition of epidural morphine-induced pruritus by intravenous droperidol [see comments]. *Reg Anesth* 18:118-120, 1993
249. Horta ML, Ramos L, Goncalves ZdR, de Oliveira MA, Tonello D, Teixeira JP, de Melo PR: Inhibition of epidural morphine-induced pruritus by intravenous droperidol. The effect of increasing the doses of morphine and of droperidol. *Reg Anesth* 21:312-317, 1996
250. Houlton PC, Downing JW, Buley RJ, Brock Utne JG: Anaesthetic induction of caesarean section with thiopentone, methohexitone and ketamine. *S Afr Med J* 54:818-820, 1978
251. Houlton PJ, Downing JW, Buley RJ, Brock Utne JG: Anaesthetic induction for caesarean section with etomidate compared with thiopentone. *S Afr Med J* 54:773-775, 1978
252. Huffnagle SL, Norris MC, Leighton BL, Arkoosh VA, Elgart RL, Huffnagle J: Do patient variables influence the subarachnoid spread of hyperbaric lidocaine in the postpartum patient? *Reg Anesth* 19(5):330-334, 1994
253. Hunt CO, Naulty JS, Bader AM, Hauch MA, Vartikar JV, Datta S, Hertwig LM, Ostheimer GW: Perioperative analgesia with subarachnoid fentanyl-bupivacaine for cesarean delivery. *Anesthesiology* 71:535-540, 1989
254. Hunt CO, Naulty JS, Malinow AM, Datta S, Ostheimer GW: Epidural butorphanol-bupivacaine for analgesia during labor and delivery. *Anesth Analg* 68:323-327, 1989
255. Hutt R, Oggunyi SO, Sullivan MH, Elder MG: Increased platelet volume and aggregation precede the onset of preeclampsia. *Obstet Gynecol* 83:146-149, 1994
256. Hwa HL, Chen RJ, Chen YC, Wang TR, Huang SC, Chow SN: Maternal and fetal outcome of pregnant women with idiopathic thrombocytopenic purpura: retrospective analysis of 25 pregnancies. *J Formos Med Assoc* 92:957-961, 1993
257. Irestedt L, Lagercrantz H, Hjemdahl P, Hagenvik K, Belfrage P: Fetal and maternal plasma catecholamine levels at elective cesarean section under general or epidural anesthesia versus vaginal delivery. *Am J Obstet Gynecol* 142:1004-1010, 1982

258. James FM, III, Crawford JS, Hopkinson R, Davies P, Naiem H: A comparison of general anesthesia and lumbar epidural analgesia for elective cesarean section. *Anesth Analg* 56:228-235, 1977
259. Johnell HE: Caesarean section. A ten-year study. *Acta Obstet Gynecol Scand* 51:231-236, 1972
260. Johnson C, Oriol N, Flood K: Trial of labor: a study of 110 patients. *J Clin Anesth* 3:216-8; 1991
261. Johnson C, Ransil BJ, Oriol N: Comparison of onset time between 0.5% bupivacaine and 3% 2-chloroprocaine with and without 75 micrograms fentanyl. *Reg Anesth* 16:228-231, 1991
262. Johnson MD, Hurley RJ, Gilbertson LI, Datta S: Continuous microcatheter spinal anesthesia with subarachnoid meperidine for labor and delivery. *Anesth Analg* 70:658-661, 1990
263. Johnstone M, Breen PJ: Halothane in obstetrics: elective caesarean section. *Br J Anaesth* 38:386-393, 1966
264. Jones G, Paul DL, Elton RA, McClure JH: Comparison of bupivacaine and bupivacaine with fentanyl in continuous extradural analgesia during labour. *Br J Anaesth* 63:254-259, 1989
265. Jouppila P, Jouppila R, Kaar K, Merila M: Fetal heart rate patterns and uterine activity after segmental epidural analgesia. *Br J Obstet Gynaecol* 84:481-486, 1977
266. Jouppila P, Kuikka J, Jouppila R, Hollm'en A: Effect of induction of general anesthesia for cesarean section on intervillous blood flow. *Acta Obstet Gynecol Scand* 58:249-253, 1979
267. Jouppila R, Hollm'en A, Jouppila P, Karki N: Segmental epidural analgesia and urinary excretion of catecholamines during labour. *Acta Anaesthesiol Scand* 21:50-54, 1977
268. Jouppila R, Hollm'en A, Jouppila P, Karppanen H: Cyclic AMP and segmental epidural analgesia during labour. *Acta Anaesthesiol Scand* 21:95-99, 1977
269. Jouppila R, Hollm'en A: The effect of segmental epidural analgesia on maternal and foetal acid-base balance, lactate, serum potassium and creatine phosphokinase during labour. *Acta Anaesthesiol Scand* 20:259-268, 1976
270. Jouppila R, Jouppila P, Karinen JM, Hollm'en A: Segmental epidural analgesia in labour: related to the progress of labour, fetal malposition and instrumental delivery. *Acta Obstet Gynecol Scand* 58:135-139, 1979
271. Jouppila R, Jouppila P, Koivisto M, et al.: Maternal, foetal and neonatal blood creatine-phosphokinase-activities and creatine-phosphokinase-isoenzymes after labour....*Acta Anaesthesiol Scand* 22:491-496, 1978
272. Jouppila R, Jouppila P, Kuikka J, Hollm'en A: Placental blood flow during caesarean section under lumbar extradural analgesia. *Br J Anaesth* 50:275-279, 1978
273. Juhani TP, Hannele H: Complications during spinal anesthesia for cesarean delivery: a clinical report of one year's experience. *Reg Anesth* 18:128-131, 1993
274. Juneja MM, Ackerman WE, Kaczorowski DM, Sollo DG, Gunzenhauser LF: Continuous epidural lidocaine infusion in the parturient with paroxysmal ventricular tachycardia. *Anesthesiology* 71:305-308, 1989
275. Juul J, Lie B, Friberg Nielsen S: Epidural analgesia vs. general anesthesia for cesarean section. *Acta Obstet Gynecol Scand Suppl* 67:203-206, 1988
276. Kaidomar M, Raucoules M, Ben Miled M, Carlon M, Grimaud D: [Epidural anesthesia and prevention of autonomic hyperreflexia in a paraplegic parturient]. *Ann Fr Anesth Reanim* 12:493-496, 1993
277. Kain ZN, Mayes LC, Ferris CA, Pakes J, Schottenfeld R: Cocaine-abusing parturients undergoing cesarean section: A cohort study. *Anesthesiology* 85:1028-1035, 1996
278. Kalappa R, Ueland K, Hansen JM, Eng M, Parer JT: Maternal acid-base status during cesarean section under thiopental, N201 and succinylcholine anesthesia. *Am J Obstet Gynecol* 109:411-417, 1971
279. Kalla GN, Fening E, Obiaya MO: Anaesthetic management of achondroplasia. *Br J Anaesth* 58:117-119, 1986
280. Kamat SK, Shah MV, Chaudhary LS, Pandya S, Bhatt MM: Effect of induction-delivery and uterine-delivery on apgar scoring of the newborn. *J Postgrad Med* 37:125-127, 1991
281. Kaminski HM, Stafl A, Aiman J: The effect of epidural analgesia on the frequency of instrumental obstetric delivery. *Obstet Gynecol* 69:770-773, 1987
282. Kangas L, Erkkola R, Kanto J, Mansikka M: Halothane anaesthesia in caesarean section. *Acta Anaesthesiol Scand* 20:189-194, 1976
283. Kantor G, Rolbin SH: Acute intermittent porphyria and caesarean delivery. *Can J Anaesth* 39:282-285, 1992
284. Karinen J, Makarainen L, Alahuhta S, Jouppila R, Jouppila P: Single bolus compared with a fractionated dose injection technique of bupivacaine for extradural Caesarean section: Effect on uteroplacental and fetal haemodynamic state. *Br J Anesth* 77:140-144, 1996
285. Kartawiadi L, Vercauteren MP, van Steenberge AL, Adriaensen HA: Spinal analgesia during labor with low-dose bupivacaine, sufentanil, and epinephrine. A comparison with epidural analgesia. *Reg Anesth* 21:191-196, 1996
286. Kelly MC, Fitzpatrick KTJ, Hill DA: Respiratory effects of spinal anaesthesia for Caesarean section. *Anaesthesia* 51:1120-1122, 1996
287. Kivalo I, Saarikoski S: Placental transfer of 14C-dimethyltubocurarine during Caesarean section. *Br J Anaesth* 48:239-242, 1976

288. Kjellmer I, Magno R, Karlsson K: Anesthesia for Cesarean section. I. Effects on the respiratory adaptation of the newborn in elective Cesarean section. *Acta Anaesthesiol Scand* 18:48-57, 1974
289. Knottnerus JA, Delgado LR, Knipschild PG, Essed GG, Smits F: Haematologic parameters and pregnancy outcome. A prospective cohort study in the third trimester. *J Clin Epidemiol* 43:461-466, 1990
290. Kolliker K: A comparison between propanididum and thiomebumal sodium used for the induction of anaesthesia for Caesarean section. *Acta Anaesthesiol Scand Suppl* 37:220-224, 1969
291. Kreitzer JM, Reed AP, Dauro AT, Brodman ML, Bronster DJ: Ascending back pain and headache during attempted epidural placement. *J Clin Anesth* 3:414-417, 1991
292. Krissel J, Dick WF, Leyser KH, Gervais H, Brockerhoff P, Schranz D: Thiopentone, thiopentone/ketamine, and ketamine for induction of anaesthesia in caesarean section [see comments]. *Eur J Anaesthesiol* 11:115-122, 1994
293. Kristoffersen MB: Cyclopropane and Caesarean section. *Br J Anaesth* 51:227-232, 1979
294. Kuhnert BR, Harrison MJ, Linn PL, Kuhnert PM: Effects of maternal epidural anesthesia on neonatal behavior. *Anesth Analg* 63:301-308, 1984
295. Kuhnert BR, Kennard MJ, Linn PL: Neonatal neurobehavior after epidural anesthesia for cesarean section: a comparison of bupivacaine and chlorprocaine. *Anesth Analg* 67:64-68, 1988
296. Kuhnert BR, Zuspan KJ, Kuhnert PM, Syracuse CD, Brown DE: Bupivacaine disposition in mother, fetus, and neonate after spinal anesthesia for cesarean section. *Anesth Analg* 66:407-412, 1987
297. Landry DP, Bennett FM, Oriol NE: Analysis of heart rate dynamics as a measure of autonomic tone in obstetrical patients undergoing epidural or spinal anesthesia. *Reg Anesth* 19:189-195, 1994
298. Lao TT, Halpern SH, MacDonald D, Huh C: Spinal subdural hematoma in a parturient after attempted epidural anaesthesia [see comments]. *Can J Anaesth* 40:340-345, 1993
299. Larue F, Benhamou D, Jullien P, Labaille T, Champagne C: Temperature monitoring during epidural anesthesia for cesarean delivery. *Reg Anesth* 16:322-324, 1991
300. Latto IP, Wainwright AC: Anaesthesia for Caesarean section. Analysis of blood concentrations of methoxyflurane using 0.1 per cent methoxyflurane and 40 per cent oxygen. *Br J Anaesth* 44:1050-1056, 1972
301. Lawes EG, Newman B, Campbell MJ, et al.: Maternal inspired oxygen concentration and neonatal status for caesarean section under general anaesthesia. Comparison of effects of 33% or 50% oxygen in nitrous oxide. *Br J Anaesth* 61:250-254, 1988
302. Leduc L, Kirshon B, Diaz SF, Cotton DB: Intrathecal morphine analgesia and low-dose dopamine for oliguria in severe maternal pulmonary hypertension. A case report. *J Reprod Med* 35:727-729, 1990
303. Leduc L, Wheeler JM, Kirshon B, Mitchell P, Cotton DB: Coagulation profile in severe preeclampsia. *Obstet & Gynecol* 79(1):14-18, 1992
304. Lee PF: Anaesthesia for Caesarean section with methohexitol. *Acta Anaesthesiol Scand Suppl* 23P138-43:138-143, 1966
305. Lee PF: Methohexitol anaesthesia in caesarean section. *Acta Obstet Gynecol Scand* 44:458-466, 1965
306. LeFevre ML: Fetal heart rate pattern and postparacervical fetal bradycardia. *Obstet Gynecol* 64:343-346, 1984
307. Lehmann KA, Stern S, Breuker KH: [Obstetrical peridural anesthesia with bupivacaine and buprenorphine. A randomized double-blind study in comparison with untreated controls]. *Anaesthetist* 41:414-422, 1992
308. Leicht CH, Ebans DE, Durkan WJ: Intrathecal sufentanil for labor analgesia: Results of a pilot study. *Anesthesiology* 73(3A):A980, 1990
309. Leighton BL, Arkoosh VA, Huffnagle S, Huffnagle HJ, Kinsella SM, Norris MC: The dermatomal spread of epidural bupivacaine with and without prior intrathecal sufentanil. *Anesth Analg* 83:526-529, 1996
310. Leighton BL, DeSimone CA, Norris MC, Ben David B: Intrathecal narcotics for labor revisited: the combination of fentanyl and morphine intrathecally provides rapid onset of profound, prolonged analgesia. *Anesth Analg* 69:122-125, 1989
311. Leng CO: Decamethonium in anaesthesia for caesarean section. *Singapore Med J* 11:173-175, 1970
312. Lew TW, Tay DH, Thomas E: Venous air embolism during cesarean section: more common than previously thought. *Anesth Analg* 77:448-452, 1993
313. Lewis M, Thomas P, Wilkes RG: Hypotension during epidural analgesia for Caesarean section. Arterial and central venous pressure changes after acute intravenous loading with two litres of Hartmann's solution. *Anaesthesia* 38:250-253, 1983
314. Lieberman BA, Rosenblatt DB, Belsey E, Packer M, Redshaw M, Mills M, Caldwell J, Notarianni L, Smith RL, Williams M, Beard RW: The effects of maternally administered pethidine or epidural bupivacaine on the fetus and newborn. *Br J Obstet Gynaecol* 86:598-606, 1979
315. Lieberman E, Lang JM, Cohen A, D'Agostino R, Jr., Datta S, Frigoletto FD, Jr. Association of epidural analgesia with cesarean delivery in nulliparas. *Obstet Gynecol* 88:993-1000, 1996
316. Lindblad A, Bernow J, Mars'al K: Fetal blood flow during intrathecal anaesthesia for elective caesarean section. *Br J Anaesth* 61:376-381, 1988

317. Lindblad A, Bernow J, Mars' al K: Obstetric analgesia and fetal aortic blood flow during labour. *Br J Obstet Gynaecol* 94:306-311, 1987
318. Lindblad A, Bernow J, Vernersson E, Mars' al K: Effects of extradural anaesthesia on human fetal blood flow in utero. Comparison of three local anaesthetic solutions. *Br J Anaesth* 59:1265-1272, 1987
319. Lipton E, Sennott FW, Batt B: Mepivacaine for spinal anesthesia in vaginal delivery. An objective comparison with tetracaine. *Am J Obstet Gynecol* 96:333-336, 1966
320. Lirzin JD, Jacquinot P, Dailland P, Jorrot JC, Jasson J, Talafre ML, Conseiller C: Controlled trial of extradural bupivacaine with fentanyl, morphine or placebo for pain relief in labour. *Br J Anaesth* 62:641-644, 1989
321. Loftus JR, Hill H, Cohen SE: Placental transfer and neonatal effects of epidural sufentanil and fentanyl administered with bupivacaine during labor. *Anesthesiology* 83:300-308, 1995
322. Loftus JR, Holbrook RH, Cohen SE: Fetal heart rate after epidural lidocaine and bupivacaine for elective cesarean section. *Anesthesiology* 75:406-412, 1991
323. Long MG, Price M, Spencer JA: Uteroplacental perfusion after epidural analgesia for elective caesarean section. *Br J Obstet Gynaecol* 95:1081-1082, 1988
324. Loughran PG, Moore J, Dundee JW: Maternal stress response associated with caesarean delivery under general and epidural anaesthesia. *Br J Obstet Gynaecol* 93:943-949, 1986
325. Low JA, Pancham SR, Worthington D, Boston RW: The incidence of fetal asphyxia in six hundred high-risk monitored pregnancies. *Am J Obstet Gynecol* 121:456-459, 1975
326. Lowenwirt IP, Zauk RM, Handwerker SM: Safety of intravenous glyceryl trinitrate in management of retained placenta. *Aust N Z J Obstet Gynaecol* 37:20-24, 1997
327. Lowson SM, Eggers KA, Warwick JP, Moore WJ, Thomas TA: Epidural infusions of bupivacaine and diamorphine in labour. *Anesthesia* 50:420-422, 1995
328. Lu JK, Manullang TR, Staples MH, Kern SE, Bailey PL: Maternal respiratory arrest, severe hypotension, and fetal distress after administration of intrathecal sufentanil and bupivacaine after intravenous fentanyl. *Anesthesiology* 87:170-172, 1997
329. Lurie S, Priscu V, Hagay ZJ: Pulse oxygen saturation in parturients receiving epidural versus opioid analgesia for labor. *J Clin Anesth* 9:87, 1997
330. Lysak SZ, Eisenach JC, Dobson CE2: Patient-controlled epidural analgesia during labor: a comparison of three solutions with a continuous infusion control [see comments]. *Anesthesiology* 72:44-49, 1990
331. Mabie WC, Anderson GD, Addington MB, Reed CM, Jr., Peeden PZ, Sibai BM: The benefit of cesarean section in acute myocardial infarction complicated by premature labor. *Obstet Gynecol* 71:2503-2506, 1988
332. MacDonald D, Grant A, Sheridan Pereira M, Boylan P, Chalmers I: The Dublin randomized controlled trial of intrapartum fetal heart rate monitoring. *Am J Obstet Gynecol* 152:524-539, 1985
333. Magno R, Kjellmer I, Karlsson K: Anesthesia for cesarean section III: effects of epidural analgesia on the respiratory adaptation of the newborn in elective cesarean section. *Acta Anaesthesiol Scand* 20:73-82, 1976
334. Magno R, Selstam U, Karlsson K: Anesthesia for cesarean section II: effects of the induction-delivery interval on the respiratory adaptation of the newborn in elective cesarean section. *Acta Anaesthesiol Scand* 19:250-259, 1975
335. Mahomed MC, Downing JW, Jeal DE, Allen PJ: Ketamine for anaesthetic induction at Caesarean section. *S Afr Med J* 50:846-848, 1976
336. Mahomed MC, Downing JW, Jeal DE, Coleman AJ: Anaesthetic induction for Caesarean section with propanidid. *Anesthesia* 31:205-211, 1976
337. Mahomed MC, Downing JW, Jeal DE, Coleman AJ: Propanidid for anaesthetic induction at Caesarean section. *S Afr Med J* 49:1358-1360, 1975
338. Mahomed MC, Downing JW, Mahomed YH: Alfathesin for anaesthetic induction at caesarean section. *S Afr Med J* 49:1095-1096, 1975
339. Mahomed YH, Downing JW, Coleman AJ, Mahomed MC: Ketamine and the obstetric patient. *S Afr Med J* 48:734-736, 1974
340. Maitra AM, Palmer SK, Bachhuber SR, Abram SE: Continuous epidural analgesia for cesarean section in a patient with morbid obesity. *Anesth Analg* 58:348-349, 1979
341. Malan TP, Jr., Johnson MD: The difficult airway in obstetric anesthesia: techniques for airway management and the role of regional anesthesia [see comments]. *J Clin Anesth* 1:104-111, 1988
342. Malinow AM, Mokriski BLK, Nomura MK, Kaufman MA, Snell JA, Sharp GD, Howard RA: Effect of epinephrine on intrathecal fentanyl analgesia in patients undergoing postpartum tubal ligation. *Anesthesiology* 73:381-385, 1990
343. Mandell GL, Jamnback L, Ramanathan S: Hemodynamic effects of subarachnoid fentanyl in laboring parturients. *Reg Anesth* 21:103-111, 1996
344. Marcoux S, Brisson J, Fabia J: The effect of cigarette smoking on the risk of preeclampsia and gestational hypertension. *Am J Epidemiol* 130:950-957, 1989

345. Margarson MP: Delayed amniotic fluid embolism following caesarean section under spinal anaesthesia. *Anaesthesia* 50:804-806, 1995
346. Marquette GP, Mechas T, Charest J, Rey E: Epidural anaesthesia for elective caesarean section does not influence fetal umbilical artery blood flow indices. *Can J Anaesth* 41:1053-1056, 1994
347. Martin JN, Jr., Harris BA, Jr., Huddleston JF, Morrison JC, Propst MG, Wiser WL, Perlis HW, Davidson JT: Vaginal delivery following previous cesarean birth. *Am J Obstet Gynecol* 146:255-263, 1983
348. Marx GF, Luykx WM, Cohen S: Fetal-neonatal status following caesarean section for fetal distress. *Br J Anaesth* 56:1009-1013, 1984
349. Matouskova A, Hanson B, Elm'en H: Continuous mini-infusion of bupivacaine into the epidural space during labor. Part III: A clinical study of 225 patients. *Acta Obstet Gynecol Scand Suppl* 43-52, 1979
350. May WJ, Greiss FC: Maternal mortality in North Carolina: a forty-year experience. *Am J Obstet Gynecol* 161:555-561, 1989
351. Mayer DC, Weeks SK: Antepartum uterine relaxation with nitroglycerin at caesarean delivery. *Can J Anaesth* 39:166-169, 1992
352. McDonald JS, Mateo CV, Reed EC: Modified nitrous oxide or ketamine hydrochloride for cesarean section. *Anesth Analg* 51:975-985, 1972
353. McGuinness GA, Merkow AJ, Kennedy RL, Erenberg A: Epidural anesthesia with bupivacaine for Cesarean section: neonatal blood levels and neurobehavioral responses. *Anesthesiology* 49:270-273, 1978
354. Meehan FP, Burke G, Kehoe JT: Update on delivery following prior cesarean section: a fifteen year review 1972-1987. *Int J Gynecol Obstet* 30:205-212, 1989
355. Meer FM, Downing JW, Coleman AJ: An intravenous method of anaesthesia for Caesarean section. II. Ketamine. *Br J Anaesth* 45:191-196, 1973
356. Meyer Breiting P, Leuwer M: [Use of fentanyl for anesthesia induction in cesarean section. Pharmacokinetics and pharmacodynamics in mother and child]. *Anaesthetist* 39:144-150, 1990
357. Mikhael MS: Emergency caesarean section in a poor risk patient in Nepal. *Br J Anaesth* 58:681-682, 1986
358. Milon D, Lavenac G, Noury D, Allain H, Van den Driessche J, Saint Marc C: [Epidural anesthesia during labor: comparison of 3 combinations of fentanyl-bupivacaine and bupivacaine alone]. *Ann Fr Anesth Reanim* 5:18-23, 1986
359. Milsom I, Forssman L, Biber B, Dottori O, Rydgren B, Sivertsson R: Maternal haemodynamic changes during caesarean section: a comparison of epidural and general anaesthesia. *Acta Anaesthesiol Scand* 29:161-167, 1985
360. Modanlou HD, Freeman RK, Ortiz O, Hinkes P, Pilsbury G Jr. Sinusoidal fetal heart rate pattern and severe fetal anemia. *Obstet Gynecol* 44(5):537-541, 1977
361. Moir DD: Anaesthesia for Caesarean section. An evaluation of a method using low concentrations of halothane and 50 per cent of oxygen. *Br J Anaesth* 42:136-142, 1970
362. Montan S, Ingemarsson I: Intrapartum fetal heart rate patterns in pregnancies complicated by hypertension. A cohort study. *Am J Obstet Gynecol* 160:283-288, 1989
363. Montan S, Ingemarsson I: Intrapartum fetal heart rate patterns in pregnancies complicated by hypertension. A cohort study. *Am J Obstet Gynecol* 160:283-288, 1989
364. Moore CH, Wilhite A, Pan PH, Blass NH: The addition of epinephrine to subarachnoid administered hyperbaric bupivacaine with fentanyl for cesarean delivery: the effect on onset time. *Reg Anesth* 17:202-204, 1992
365. Moore J, McNabb TG, Dundee JW: Preliminary report on ketamine in obstetrics. *Br J Anaesth* 43:779-782, 1971
366. Moore TR, Key TC, Reisner LS, Resnik R: Evaluation of the use of continuous lumbar epidural anesthesia for hypertensive pregnant women in labor. *Am J Obstet Gynecol* 152:404-412, 1985
367. Morgan M: Anaesthetic contribution to maternal mortality. *Br J Anaesth* 59:842-855, 1987
368. Munson AK, Scott JR: Postpartum tubal ligation under local anesthesia. *Obstet Gynecol* 39(5):756-758, 1972
369. Murphy JD, Henderson K, Bowden MI, Lewis M, Cooper GM: Bupivacaine versus bupivacaine plus fentanyl for epidural analgesia: effect on maternal satisfaction [see comments]. *Br Med J (Clinical Research Ed* 302:564-567, 1991
370. Naef RW3, Chauhan SP, Chevalier SP, Roberts WE, Meydreich EF, Morrison JC: Prediction of hemorrhage at cesarean delivery. *Obstet Gynecol* 83:923-926, 1994
371. Nageotte MP, Larson D, Rumney PJ, Sidhu M, Hollenbach K: Epidural analgesia compared with combined spinal-epidural analgesia during labor in nulliparous women. *New Eng J Med* 337(24):1715-1719, 1997
372. Nasrat H, Fageeh W, Abalkhail B, Yamani T, Ardawi MSM: Determinants of pregnancy outcome in patients with gestational diabetes. *Int J Gynaecol Obstet* 53:117-123, 1996
373. Neeld JB, Jr., Seabrook PD, Jr., Chastain GM, Frederickson EL: A clinical comparison of pancuronium and tubocurarine for cesarean section anesthesia. *Anesth Analg* 53:7-11, 1974
374. Nelson DM, Main E, Crafford W, Ahumada GG: Peripartum heart failure due to primary pulmonary hypertension. *Obstet Gynecol* 62:58s-63s, 1983

375. Newman LM, Patel RV, Krolick T, Ivankovich AD: Muscular spasm in the lower limbs of laboring patients after intrathecal administration of epinephrine and sufentanil [see comments]. *Anesthesiology* 80:468-471, 1994
376. Newsome LR, Bramwell RS, Curling PE: Severe preeclampsia: hemodynamic effects of lumbar epidural anesthesia. *Anesth Analg* 65:31-36, 1986
377. Newton ER, Schroeder BC, Knape KG, Bennett BL: Epidural analgesia and uterine function [see comments]. *Obstet Gynecol* 85(5):749-755, 1995
378. Ng PC, Wong MY, Nelson EA: Paediatrician attendance at caesarean section. *Eur J Pediatr* 154:672-675, 1995
379. Nguyen Thi TV, Orliaguet G, Ngu TH, Bonnet F: Spinal anesthesia with meperidine as the sole agent for cesarean delivery. *Reg Anesth* 19:386-389, 1994
380. Nielsen PE, Erickson JR, Abouleish EI, Perriatt S, Sheppard C: Fetal heart rate changes after intrathecal sufentanil or epidural bupivacaine for labor analgesia: incidence and clinical significance. *Anesth Analg* 83:742-746, 1996
381. Nieminen K, Puolakka J: Effective obstetric paracervical block with reduced dose of bupivacaine - A prospective randomized double-blind study comparing 25 mg (0.25%) and 12.5 mg (0.125%) of bupivacaine. *Acta Obstet Gynecol Scand* 76:50-54, 1997
382. Niv D, Rudick V, Golan A, Chayen MS: Augmentation of bupivacaine analgesia in labor by epidural morphine. *Obstet Gynecol* 67:206-209, 1986
383. Noble HA, Enever GR, Thomas TA: Epidural bupivacaine dilution for labour. A comparison of three concentrations infused with a fixed dose of fentanyl. *Anaesthesia* 46:549-552, 1991
384. Nootens M, Rich S: Successful management of labor and delivery in primary pulmonary hypertension. *Am J Cardiol* 71:1124-1125, 1993
385. Norris MC, Grieco WM, Borkowski M, Leighton BL, Arkoosh VA, Huffnagle HJ, Huffnagle S: Complications of labor analgesia: epidural versus combined spinal epidural techniques [published erratum appears in *Anesth Analg* 79:529-37, 1994]. *Anesth Analg* 79:529-37, 1994
386. Norris MC, Honet JE, Leighton BL, Arkoosh VA: A comparison of meperidine and lidocaine for spinal anesthesia for postpartum tubal ligation [see comments]. *Reg Anesth* 21:84-88, 1996
387. Ohel G, Harats H: Epidural anesthesia in early compared with advanced labor. *Int J Gynaecol Obstet* 45:217-219, 1994
388. Ojo OA, Akinkugbe OO: Nontoxicemic hypertension in pregnancy in the African indigene. An analysis of 30 cases. *Am J Obstet Gynecol* 105:938-941, 1969
389. Olofsson CIJ, Ekblom AOA, EkmanOrdeberg GE, Irestedt LE: Post-partum urinary retention: A comparison between two methods of epidural analgesia. *Eur J Obstet Gynecol Reprod Biol* 71:31-34, 1997
390. Ong BY, Cohen MM, Palahniuk RJ: Anesthesia for cesarean section--effects on neonates. *Anesth Analg* 68:270-275, 1989
391. Owens WD, Zeitlin GL: Hypoventilation in a newborn following administration of succinylcholine to the mother: a case report. *Anesth Analg* 54:38-40, 1975
392. Paech MJ, Westmore MD, Speirs HM: A double-blind comparison of epidural bupivacaine and bupivacaine-fentanyl for caesarean section. *Anaesth Intensive Care* 18:22-30, 1990
393. Palahniuk RJ, Scatliff J, Biehl D, Wiebe H, Sankaran K: Maternal and neonatal effects of methoxyflurane, nitrous oxide and lumbar epidural anaesthesia for Caesarean section. *Can Anaesth Soc J* 24:586-596, 1977
394. Palmer CM, Voulgaropoulos D, Alves D: Subarachnoid fentanyl augments lidocaine spinal anesthesia for cesarean delivery. *Reg Anesth* 20:389-394, 1995
395. Parker J, Balis N, Chester S, Adey D: Cardiopulmonary arrest in pregnancy: successful resuscitation of mother and infant following immediate caesarean section in labour ward. *Aust N Z J Obstet Gynaecol* 36:207-210, 1996
396. Pash MP, Balaton J, Eagle C: Anaesthetic management of a parturient with severe muscular dystrophy, lumbar lordosis and a difficult airway. *Can J Anaesth* 43:959-963, 1996
397. Paul RH, Freeman RK. Fetal cardiac response to paracervical block anesthesia: part II. *Am J Obstet Gynacol* 113:592-597, 1972
398. Pearson JF, Davies P: The effect of continuous lumbar epidural analgesia on the acid-base status of maternal arterial blood during the first stage of labour. *J Obstet Gynaecol Br Commonw* 80:218-224, 1973
399. Pearson JF, Davies P: The effect of continuous lumbar epidural analgesia upon fetal acid-base status during the second stage of labour. *J Obstet Gynaecol Br Commonw* 81:975-979, 1974
400. Pedersen JE, Fernandes A, Christensen M: Halothane 2% for caesarean section. *Eur J Anaesthesiol* 9:319-324, 1992
401. Pello LC, Rosevear SK, Dawes GS, Moulden M, Redman CW: Computerized fetal heart rate analysis in labor. *Obstet Gynecol* 78:602-610, 1991
402. Peltz B, Sinclair DM: Induction agents for Caesarean section. A comparison of thiopentone and ketamine. *Anaesthesia* 28:37-42, 1973
403. Peng AT, Blancato LS, Motoyama EK: Effect of maternal hypcapnia v. eucapnia on the foetus during Caesarean section. *Br J Anaesth* 44:1173-1178, 1972

404. Peng AT, Gorman RS, Shulman SM, DeMarchis E, Nyunt K, Blancato LS: Intravenous nitroglycerin for uterine relaxation in the postpartum patient with retained placenta [letter]. *Anesthesiology* 71:172-173, 1989
405. Petrikovsky B, Cohen M, Fastman D, Tancer ML: Electronic fetal heart rate monitoring during cesarean section. *Int J Gynaecol Obstet* 26:203-207, 1988
406. Pham LH, Camann WR, Smith MP, Datta S, Bader AM: Hemodynamic effects of intrathecal sufentanil compared with epidural bupivacaine in laboring parturients. *J Clin Anesth* 8:497-501; 1996
407. Phelan JP, Yurth DA. Severe preeclampsia: I. Peripartum hemodynamic observations. *Am J Obstet Gynecol* 144:17-22, 1982
408. Philipsen T, Jensen NH: Epidural block or parenteral pethidine as analgesic in labour; a randomized study concerning progress in labour and instrumental deliveries. *Eur J Obstet Gynecol Reprod Biol* 30:27-33, 1989
409. Philipsen T, Jensen NH: Maternal opinion about analgesia in labour and delivery. A comparison of epidural blockade and intramuscular pethidine. *Eur J Obstet Gynecol Reprod Biol* 34:1990
410. Philipson EH, Kuhnert BR, Syracuse CD: Fetal acidosis, 2-chloroprocaine, and epidural anesthesia for cesarean section. *Am J Obstet Gynecol* 151:322-324, 1985
411. Phillips G: Continuous infusion epidural analgesia in labor: the effect of adding sufentanil to 0.125% bupivacaine. *Anesth Analg* 67:462-465, 1988
412. Phillips GH: Epidural sufentanil/bupivacaine combinations for analgesia during labor: effect of varying sufentanil doses. *Anesthesiology* 67:835-838, 1987
413. Pierce ET, Denson DD, Essell SK, Santos DJ, Edstrom HH: The effect of rate of infusion on continuous epidural analgesia for labor and delivery. *Reg Anesth* 14:31-34, 1989
414. Ploeckinger B, Ulm MR, Chalubinski K, Gruber W: Epidural anaesthesia in labour: influence on surgical delivery rates, intrapartum fever and blood loss. *Gynecol Obstet Invest* 39:24-27, 1995
415. Porter JS, Bonello E, Reynolds F: The effect of epidural opioids on maternal oxygenation during labour and delivery. *Anaesthesia* 51:899-903, 1996
416. Rafferty TD, Berkowitz RL: Hemodynamics in patients with severe toxemia during labor and delivery. *Am J Obstet Gynecol* 138:263-270, 1980
417. Ramanathan J, Bottorff M, Jeter JN, Khalil M, Sibai BM: The pharmacokinetics and maternal and neonatal effects of epidural lidocaine in preeclampsia. *Anesth Analg* 65:120-126, 1986
418. Ramanathan J, Sibai BM, Vu T, Chauhan D: Correlation between bleeding times and platelet counts in women with preeclampsia undergoing cesarean section. *Anesthesiology* 71:188-191, 1989
419. Ramin SM, Gambling DR, Lucas MJ, Sharma SK, Sidawi JE, Leveno KJ: Randomized trial of epidural versus intravenous analgesia during labor [see comments]. *Obstet Gynecol* 86:783-789, 1995
420. Ramin SM, Ramin KD, Cox K, Magness RR, Shearer VE, Gant NF: Comparison of prophylactic angiotensin II versus ephedrine infusion for prevention of maternal hypotension during spinal anesthesia [see comments]. *Am J Obstet Gynecol* 171:734-739, 1994
421. Ranta P, Jouppila P, Spalding M, Jouppila R: The effect of maternal obesity on labour and labour pain. *Anaesthesia* 50:322-326, 1995
422. Ratcliffe FM, Evans JM: Neonatal wellbeing after elective caesarean delivery with general, spinal, and epidural anaesthesia. *Eur J Anaesthesiol* 10:175-181, 1993
423. Rawal N, Schollin J, Wesstrom G: Epidural versus combined spinal epidural block for cesarean section. *Acta Anaesthesiol Scan* 32:61-66, 1988
424. Reddy BK, Pizer B, Bull PT: Neonatal serum cortisol suppression by etomidate compared with thiopentone, for elective caesarean section. *Eur J Anaesthesiol* 5:171-176, 1988
425. Redman CWG, Bonnar J, Beilin L: Early platelet consumption in pre-eclampsia. *Br Med J* 1:467-469, 1978
426. Reubinoff BE, Eldor A, Laufer N, Sadovsky E: Maternal hemorrhagic complications following prophylactic low-dose aspirin and dipyridamole therapy. *Gynecol Obstet Invest* 33:241-243, 1992
427. Reynolds F, O'Sullivan G: Epidural fentanyl and perineal pain in labour [see comments]. *Anaesthesia* 44:341-344, 1989
428. Riley ET, Flanagan B, Cohen SE, Chitkara U: Intravenous nitroglycerin: a potent uterine relaxant for emergency obstetric procedures. Review of literature and report of three cases. *Int J Obstet Anesth* 5:264-268, 1996
429. Riley ET, Ratner EF, Cohen SE: Intrathecal sufentanil for labor analgesia: Do sensory changes predict better analgesia and greater hypotension? *Anesth Analg* 84:346-351, 1997
430. Roberts SW, Leveno KJ, Sidawi JE, Lucas MJ, Kelly MA: Fetal acidemia associated with regional anesthesia for elective cesarean delivery [see comments]. *Obstet Gynecol* 85:79-83, 1995
431. Roberts WE, Perry KG, Jr., Woods JB, Files JC, Blake PG, Martin JN, Jr. The intrapartum platelet count in patients with HELLP (hemolysis, elevated liver enzymes, and low platelets) syndrome: is it predictive of later hemorrhagic complications? *Am J Obstet Gynecol* 171:799-804, 1994

432. Robertson A, Fothergill RJ, Bond RA: Effects of propanidid and thiopentone on the acid-base status of babies delivered by elective caesarean section. *S Afr Med J* 48:1019-1022, 1974
433. Robinson JO, Rosen M, Evans JM, Revill SI, David H, Rees GA: Maternal opinion about analgesia for labour. A controlled trial between epidural block and intramuscular pethidine combined with inhalation. *Anaesthesia* 35:1173-1181, 1980
434. Robson SC, Boys RJ, Rodeck C, Morgan B: Maternal and fetal haemodynamic effects of spinal and extradural anaesthesia for elective caesarean section [see comments]. *Br J Anaesth* 68:54-59, 1992
435. Robson SC, Dunlop W, Hunter S, Boys R, Bryson M: Haemodynamic changes associated with caesarean section under epidural anaesthesia. *Br J Obstet Gynaecol* 96:642-647, 1989
436. Robson SC, Samsoon G, Boys RJ, Rodeck C, Morgan B: Incremental spinal anaesthesia for elective caesarean section: maternal and fetal haemodynamic effects. *Br J Anaesth* 70:634-638, 1993
437. Rodriguez J, Abboud TK, Reyes A, Payne M, Zhu J, Steffens Z, Afrasiabi A: Continuous infusion epidural anesthesia during labor: a randomized, double-blind comparison of 0.0625% bupivacaine/0.002% butorphanol and 0.125% bupivacaine. *Reg Anesth* 15:300-3, 1990
438. Rolbin SH, Cohen MM, Levinton CM, Kelly EN, Farine D: The premature infant: anesthesia for cesarean delivery [see comments]. *Anesth Analg* 78:912-917, 1994
439. Romero R, Mazor M, Lockwood CJ, Emamian M, Belanger KP, Hobbins JC, Duffy T: Clinical significance, prevalence, and natural history of thrombocytopenia in pregnancy-induced hypertension. *Am J Perinatol* 6:32-38, 1989
440. Ron M, Menashe M, Scherer D, Palti Z: Fetal heart rate decelerations following the administration of meperidine-promethazine during labor. *Int J Gynaecol Obstet* 20:301-305, 1982
441. Ross RM, Baker T: Cardiac enzymes in patients undergoing caesarean section. *Can J Anaesth* 42:46-50, 1995
442. Rozkowski A, Smyczek D, Birnbach DJ: Continuous spinal anesthesia for cesarean delivery in a patient with arthrogryposis multiplex congenita: A clinical report. *Reg Anesth* 21:477-479, 1996
443. Rubin GL, Peterson HB, Rochat RW, McCarthy BJ, Terry JS: Maternal death after cesarean section in Georgia. *Am J Obstet Gynecol* 139:681-685, 1981
444. Russell IF: Spinal anesthesia for cesarean delivery with dilute solutions of plain bupivacaine: the relationship between infused volume and spread. *Reg Anesth* 16:130-136, 1991
445. Russell R, Reynolds F: Epidural infusion of low-dose bupivacaine and opioid in labour. Does reducing motor block increase the spontaneous delivery rate? *Anaesthesia* 51:266-273, 1996
446. Rust LA, Waring RW, Hall GL, Nelson EI: Intrathecal narcotics for obstetric analgesia in a community hospital. *Am J Obstet Gynecol* 170:1643-6; 1994
447. Ryhanen P, Jouppila R, Lanning M, Jouppila P, Hollm'en A, Kouvalainen K: Natural killer cell activity after elective cesarean section under general and epidural anesthesia in healthy parturients and their newborns. *Gynecol Obstet Invest* 19:139-142, 1985
448. Sachs BP, Oriol NE, Ostheimer GW, Weiss JB, Driscoll S, Acker D, Brown DA, Jewett JF: Anesthetic-related maternal mortality, 1954 to 1985. *J Clin Anesth* 1:333-338, 1989
449. Sakala EP, Kaye S, Murray RD, Munson LJ: Epidural analgesia: effect on the likelihood of a successful trial of labor after cesarean section. *J Reprod Med* 35(9):886-890, 1990
450. Schoenfeld A, Joel Cohen SJ, Duparc H, Levy E: Emergency obstetric anaesthesia and the use of beta2-sympathomimetic drugs. *Br J Anaesth* 50:969-971, 1978
451. Schwarcz RL, Belizan JM, Cifuentes JR, Cuadro JC, Marques MB, Caldeyro Barcia R: Fetal and maternal monitoring in spontaneous labors and in elective inductions. A comparative study. *Am J Obstet Gynecol* 120:356-362, 1974
452. Schwartz ML, Brenner W: Toxemia in a patient with none of the standard signs and symptoms of preeclampsia. *Obstet Gynecol* 66:19S-21S, 1985
453. Scott DB, Lees MM, Davie IT, Slawson KB, Kerr MG: Observations on cardiorespiratory function during caesarean section. *Br J Anaesth* 41:489-495, 1969
454. Scott PV, Bowen FE, Cartwright P, Mohan Rao BC, Deeley V, Wotherspoon HG, Sumarein IMA: Intrathecal morphine as sole analgesic during labour. *Br Med J* 281:351-353, 1980
455. Scull TJ, Carli F: Cardiac arrest after Caesarean section under subarachnoid block. *Br J Anaesth* 77:274-276, 1996
456. Sears DH, Leeman MI, Jassy LJ, O'Donnell LA, Allen SG, Reisner LS: The frequency of postdural puncture headache in obstetric patients: a prospective study comparing the 24-gauge versus the 22 gauge sprotte needle. *J Clin Anesth* 6:42-46, 1994
457. Sharma SK, Gajraj NM, Sidawi JE: Prevention of hypotension during spinal anesthesia: A comparison of intravascular administration of hetastarch versus lactated ringer's solution. *Anesth Analg* 84:111-114, 1997
458. Sharma SK, Gambling DR, Joshi GP, Sidawi JE, Herrera ER: Comparison of 26-gauge Atraucan and 25-gauge Whitacre needles: insertion characteristics and complications. *Can J Anaesth* 42(8):706-710, 1995

459. Sharma SK, Herrera ER, Sidawi JE, Leveno KJ: The pregnant patient with an intracranial arteriovenous malformation. Cesarean or vaginal delivery using regional or general anesthesia? *Reg Anesth* 20:455-458, 1995
460. Sharma SK, Sidawi JE, Ramin SM, Lucas MJ, Leveno KJ, Cunningham G: Cesarean delivery: A randomized trial of epidural versus patient-controlled meperidine analgesia during labor. *Anesthesiology* 87:487-595, 1997
461. Shehabi Y, Gatt S, Buckman T, Isert P: Effect of adrenaline, fentanyl and warming of injectate on shivering following extradural analgesia in labour. *Anaesth Intensive Care* 18:31-37, 1990
462. Shennan A, Cooke V, Lloyd Jones F, Morgan B, de Swiet M: Blood pressure changes during labour and whilst ambulating with combined spinal epidural analgesia. *Br J Obstet Gynaecol* 102:192-197, 1995
463. Shyken JM, Smeltzer JS, Baxi LV, Blakemore KJ, Ambrose SE, Petrie RH: A comparison of the effect of epidural, general, and no anesthesia on funic acid-base values by stage of labor and type of delivery [see comments]. *Am J Obstet Gynecol* 163:802-807, 1990
464. Simon L, Santi TM, Sacquin P, Hamza J: Pre-anaesthetic assessment of coagulation abnormalities in obstetric patients: usefulness, timing and clinical implications. *Br J Anaesth* 78:678-683, 1997
465. Sinatra RS, Goldstein R, Sevarino FB: The clinical effectiveness of epidural bupivacaine, bupivacaine with lidocaine, and bupivacaine with fentanyl for labor analgesia. *J Clin Anesth* 3:219-24; 1991
466. Skerman JH, Thompson BA, Goldstein MT, Jacobs MA, Gupta A, Blass NH: Combined continuous epidural fentanyl and bupivacaine in labor: A randomized study. *Anesthesiology* 63(3A):A450, 1986
467. Skovsted P, Misfeldt BB, Mogensen JV, Brockner J: The effects of anesthesia on blood loss at caesarean section. *Acta Anaesthesiol Scand* 17:153-158, 1973
468. Sparks CJ, Perndt H, Agiomea K, Fa'arondo J: Spinal anaesthesia for caesarean section in the Solomon Islands [see comments]. *Anaesth Intensive Care* 22:187-191, 1994
469. Speirs I, Sim AW: The placental transfer of pancuronium bromide. *Br J Anaesth* 44:370-373, 1972
470. Speirs I: The use of tacrine and suxamethonium in anaesthesia for caesarean section. *Br J Anaesth* 38:394-400, 1966
471. Spencer JA, Koutsoukis M, Lee A: Fetal heart rate and neonatal condition related to epidural analgesia in women reaching the second stage of labour. *Eur J Obstet Gynecol Reprod Biol* 41:1991
472. Spinapolice RX: The effects of continuous epidural anesthesia on fetal heart rate during labor. *J Am Osteopath Assoc* 76:695-698, 1977
473. Stacey RG, Watt S, Kadim MY, Morgan BM: Single space combined spinal-extradural technique for analgesia in labour [see comments]. *Br J Anaesth* 71:499-502, 1993
474. Stavrou C, Hofmeyr GJ, Boezaart AP: Prolonged fetal bradycardia during epidural analgesia. Incidence, timing and significance. *S Afr Med J* 77:66-68, 1990
475. Steinberg RB, Dunn SM, Dixon DE, Rehm KL, Pastides H, Hu X: Comparison of sufentanil, bupivacaine, and their combination for epidural analgesia in obstetrics. *Reg Anesth* 17:131-138, 1992
476. Stoddart AP, Nicholson KE, Popham PA: Low dose bupivacaine/fentanyl epidural infusions in labour and mode of delivery. *Anesthesia* 49:1087-1090, 1994
477. Stovall TG, Shaver DC, Solomon SK, Anderson GD: Trial of labor in previous cesarean section patients, excluding classical cesarean sections. *J Obstet Gynecol* 70(5):713-717, 1987
478. Stovner J, Vangen O: Diazepam compared to thiopentone as induction agent for caesarean sections. *Acta Anaesthesiol Scand* 18:264-269, 1974
479. Strauss RG, Keefer JR, Burke T, Civetta JM: Hemodynamic monitoring of cardiogenic pulmonary edema complicating toxemia of pregnancy. *Obstet Gynecol* 55:170-174, 1980
480. Strickland DM, Guzick DS, Cox K, Gant NF, Rosenfeld CR: The relationship between abortion in the first pregnancy and development of pregnancy-induced hypertension in the subsequent pregnancy. *Am J Obstet Gynecol* 154:146-148, 1986
481. Suppan P: Althesin anaesthesia and a long induction-delivery interval at Caesarean section. *Anaesthesia* 31:910-912, 1976
482. Sutcliffe N, Remington SA, Ramsay TM, Mason C: Severe tracheal stenosis and operative delivery. *Anaesthesia* 50:26-29, 1995
483. Swayze CR, Skerman JH, Walker EB, Sholte FG: Efficacy of subarachnoid meperidine for labor analgesia. *Reg Anesth* 16:309-313, 1991
484. Tay DHB, Tay SM, Thomas E: High-volume spinal anaesthesia: A dose-response study of bupivacaine 0.125%. *Anaesth Intens Care* 20:443-447, 1992
485. Taylor AB, Bryant MT, Ghosh Ray GC: Cesarean section with regional anesthesia using an extendable epidural block. *Int J Gynaecol Obstet* 16:385-387, 1979
486. Taylor AB: Fetal heart rate monitoring during caesarean section. *Br J Obstet Gynaecol* 84:281-284, 1977
487. Teviotdale BM: Vecuronium-thiopentone induction for emergency caesarean section under general anaesthesia [see comments]. *Anaesth Intensive Care* 21:288-291, 1993

488. Thaler I, Timor Tritsch IE, Blumenfeld Z: Effect of acute hypoxia on human fetal heart rate. The significance of increased heart rate variability. *Acta Obstet Gynecol Scand Suppl* 64:47-50, 1985
489. Thalme B, Belfrage P, Raabe N: Lumbar epidural analgesia in labour. I. Acid-base balance and clinical condition of mother, fetus and newborn child. *Acta Obstet Gynecol Scand* 53:27-35, 1974
490. Thor'en T, Holmstrom B, Rawal N, Schollin J, Lindeberg S, Skeppner G: Sequential combined spinal epidural block versus spinal block for cesarean section: effects on maternal hypotension and neurobehavioral function of the newborn [see comments]. *Anesth Analg* 78:1087-1092, 1994
491. Thorp JA, Hu DH, Albin RM, McNitt J, Meyer BA, Cohen GR, Yeast JD: The effect of intrapartum epidural analgesia on nulliparous labor: a randomized, controlled, prospective trial. *Am J Obstet Gynecol* 169:851-858, 1993
492. Thorp JA, Parisi VM, Boylan PC, Johnston DA: The effect of continuous epidural analgesia on cesarean section for dystocia in nulliparous women [see comments]. *Am J Obstet Gynecol* 161:670-675, 1989
493. Timor Tritsch I, Gergely Z, Abramovici H, Brandes JM: Electronic monitoring of the fetal heart rate and uterine contractions during cesarean section under balanced anesthesia. *Obstet Gynecol* 48:292-296, 1976
494. Toldy M, Scott DB: Blood loss during Caesarean section under general anaesthesia. *Br J Anaesth* 41:868-873, 1969
495. Trudinger BJ: Platelets and intrauterine growth retardation in pre-eclampsia. *Br J Obstet Gynecol* 83:284-286, 1976
496. Tunstall ME, Geddes C: "Failed intubation" in obstetric anesthesia. *Br J Anaesth* 56:659-661, 1984
497. Tunstall ME, Sheikh A: Comparison of 1.5% enflurane with 1.25% isoflurane in oxygen for caesarean section: avoidance of awareness without nitrous oxide. *Br J Anaesth* 62:138-143, 1989
498. Van Steenberge A, Debroux HC, Noorduin H: Extradural bupivacaine with sufentanil for vaginal delivery. A double-blind trial. *Br J Anaesth* 59:1518-1522, 1987
499. Vella LM, Willatts DG, Knott C, Lintin DJ, Justins DM, Reynolds F: Epidural fentanyl in labour. An evaluation of the systemic contribution to analgesia. *Anaesthesia* 40:741-747, 1985
500. Veren D, Boehm FH, Killam AP: The clinical significance of a sinusoidal fetal heart rate pattern associated with alphaprodine administration. *J Reprod Med* 27(7): 411-414, 1982
501. Vertommen JD, Van Aken H, Vandermeulen E, Vangerven M, Devlieger H, Van Assche AF, Shnider SM: Maternal and neonatal effects of adding epidural sufentanil to 0.5% bupivacaine for cesarean delivery. *J Clin Anesth* 3:371-376, 1991
502. Vertommen JD, Vandermeulen E, Van Aken H, Vaes L, Soetens M, Van Steenberge A, Mourisse P, Willaert J, Noorduin H, Devlieger H, et al: The effects of the addition of sufentanil to 0.125% bupivacaine on the quality of analgesia during labor and on the incidence of instrumental deliveries [see comments]. *Anesthesiology* 74:809-814, 1991
503. Vickers RJ, May AE: Long-term backache after extradural or general anaesthesia for manual removal of placenta: preliminary report [see comments]. *Br J Anaesth* 70:214-215, 1993
504. Vincent RD, Jr., Reid RW: Epidural anesthesia for postpartum tubal ligation using epidural catheters placed during labor. *J Clin Anesth* 5:289-291, 1993
505. Viscomi CM, Hood DD, Melone PJ, Eisenach JC: Fetal heart rate variability after epidural fentanyl during labor. *Anesth Analg* 71:679-683, 1990
506. Viscomi CM, Rathmell JP, Mason SB, Livermore M, Schapiro H: Analgesic efficacy and side effects of subarachnoid sufentanil-bupivacaine administered to women in advanced labor. *Reg Anesth* 21:424-429, 1996
507. Viscomi CM, Rathmell JP: Labor epidural catheter reactivation or spinal anesthesia for delayed postpartum tubal ligation: a cost comparison. *J Clin Anesth* 7:380-383, 1995
508. Visser W, van Pampus MG, Treffers PE, Wallenburg HC: Perinatal results of hemodynamic and conservative temporizing treatment in severe pre-eclampsia. *Eur J Obstet Gynecol Reprod Biol* 53:1994
509. Visser W, Wallenburg HC: Maternal and perinatal outcome of temporizing management in 254 consecutive patients with severe pre-eclampsia remote from term. *Eur J Obstet Gynecol Reprod Biol* 63:147-154, 1995
510. Visser W, Wallenburg HC: Temporising management of severe pre-eclampsia with and without the HELLP syndrome. *Br J Obstet Gynaecol* 102:111-117, 1995
511. Visser W, Wallenburg HCS: Central hemodynamic observations in untreated preeclamptic patients. *Hypertension* 17:1072-1077, 1991
512. Wallace DH, Leveno KJ, Cunningham FG, Giesecke AH, Shearer VE, Sidawi JE: Randomized comparison of general and regional anesthesia for cesarean delivery in pregnancies complicated by severe preeclampsia. *Obstet Gynecol* 86:193-199, 1995
513. Wallenburg HCS: Invasive hemodynamic monitoring in pregnancy. *Eur J Obstet Gynecol Reprod Biol* 42:S45-S51, 1991
514. Walmsley AJ, Giesecke AH, Lipton JM: Contribution of extradural temperature to shivering during extradural anaesthesia. *Br J Anaesth* 58:1130-1134, 1986
515. Warren TM, Datta S, Ostheimer GW, Naulty JS, Weiss JB, Morrison JA: Comparison of the maternal and neonatal effects of halothane, enflurane, and isoflurane for cesarean delivery. *Anesth Analg* 62:516-520, 1983

516. Weekes AR, Cheridjian VE, Mwanje DK: Lumbar epidural analgesia in labour in twin pregnancy. *Br Med J* 2:730-732, 1977
517. Wiener PC, Hogg MI, Rosen M: Neonatal respiration, feeding and neurobehavioural state. Effects of intrapartum bupivacaine, pethidine and pethidine reversed by naloxone. *Anaesthesia* 34:996-1004, 1979
518. Withington DE, Weeks SK: Repeat epidural analgesia and unilateral block. *Can J Anaesth* 41:568-571, 1994
519. Wolfe HM, Gross TL, Sokol RJ, Bottoms SF, Thompson KL: Determinants of morbidity in obese women delivered by cesarean. *Obstet Gynecol* 71:691-696, 1988
520. Yancey MK, Pierce B, Schweitzer D, Daniels D: Observations on labor epidural analgesia and operative delivery rates. (UNPUBLISHED PAPER)
521. Yau G, Gin T, Ewart MC, Kotur CF, Leung RK, Oh TE: Propofol for induction and maintenance of anaesthesia at caesarean section. A comparison with thiopentone/enflurane. *Anaesthesia* 46:20-23, 1991
522. Yau G, Gregory MA, Gin T, Bogod DG, Oh TE: The addition of fentanyl to epidural bupivacaine in first stage labour. *Anaesth Intensive Care* 18:532-535, 1990
523. Yau G, Gregory MA, Gin T, Oh TE: Obstetric epidural analgesia with mixtures of bupivacaine, adrenaline and fentanyl. *Anaesthesia* 45:1020-1023, 1990
524. Youngstrom P, Eastwood D, Patel H, Bhatia R, Cowan R, Sutheimer C: Epidural fentanyl and bupivacaine in labor: double-blind study. *Anesthesiology* 61(3A):A414, 1984
525. Yun E, Topulos GP, Body SC, Datta S, Bader AM: Pulmonary function changes during epidural anesthesia for cesarean delivery. *Anesth Analg* 82:750-753, 1996
526. Zaaijman JD, Slabber CF: Fetal heart rate changes associated with standard, segmental and modified segmental epidural block. *S Afr Med J* 54:481-482, 1978
527. Zagorzycki MT, Brinkman CR, III: The effect of general and epidural anesthesia upon neonatal Apgar scores in repeat cesarean section. *Surg Gynecol Obstet* 155:641-645, 1982
528. Zilianti M, Salazar JR, Aller J, Aguero O: Fetal heart rate and pH of fetal capillary blood during epidural analgesia in labor. *Obstet Gynecol* 36(6):881-886, 1970