

Practice Guidelines for Obstetric Anesthesia: An Updated Report
American Society of Anesthesiologists

Bibliography in Alphabetical Order

1. Abboud TK, Afrasiabi A, Zhu J, Mantilla M, Reyes A, D'Onofrio L, Khoo N, Mosaad P, Richardson M, Kalra M, Cheung M, Paul R: Epidural morphine or butorphanol augments bupivacaine analgesia during labor. *Reg Anesth* 1989; 14:115-120
2. Abboud TK, Khoo SS, Miller F, Doan T, Henriksen EH: Maternal, fetal, and neonatal responses after epidural anesthesia with bupivacaine, 2-chloroprocaine, or lidocaine. *Anesth Analg* 1982; 61:638-644
3. Abboud TK, Nagappala S, Murakawa K, David S, Haroutunian S, Zakarian M, Yanagi T, Sheikh-Ol-Eslam A: Comparison of the effects of general and regional anesthesia for cesarean section on neonatal neurologic and adaptive capacity scores. *Anesth Analg* 1985; 64:996-1000
4. Abboud TK, Zhu J, Afrasiabi A, Reyes A, Sherman G, Khan R, Vera Cruz R, Steffens Z: Epidural butorphanol augments lidocaine sensory anesthesia during labor. *Reg Anesth* 1991; 16:265-267
5. Abouleish E: Caudal analgesia for quadruplet delivery. *Anesth Analg* 1976; 55:61-66
6. Abouleish E: Foetal bradycardia during caudal analgesia: A discussion of possible causative factors. *Br J Anaesth* 1976; 48:481-484
7. Abouleish EI: Postpartum tubal ligation requires more bupivacaine for spinal anesthesia than does cesarean section. *Anesth Analg* 1986; 65:897-900
8. Adams AP, Morgan M, Jones BC, McCormick PW: A case of massive aspiration of gastric contents during obstetric anaesthesia. Treatment by tracheostomy and prolonged intermittent positive pressure ventilation. *Br J Anaesth* 1969; 41:176-183
9. Adams HA, Meyer P, Stoppa A, Muller-Goch A, Bayer P, Hecker H: Anaesthesia for caesarean section. Comparison of two general anaesthetic regimens and spinal anaesthesia. *Anaesthesia* 2003; 52:23-32
10. Ahmed SM, Khan RM, Bano S, Ajmani P, Kumar A: Is spinal anaesthesia safe in pre-eclamptic toxæmia patients? *J Indian Med Assoc* 1999; 97:165-168
11. Alahuhta S, Rasanen J, Jouppila P, Hollmen AI: Ephedrine and phenylephrine for avoiding maternal hypotension due to spinal anaesthesia for caesarean section. *Int J Obstet Anesth* 1992; 1:129-134
12. Alfrevic Z, Elbourne D, Pavord S, Bolte An, Van Geijn H, Mercier F, Ahonen J, Bremme K, Bodker B, Magnusdottir E, Salvesen K, Prendiville W, Truesdale A, Clemens F, Piercy D, Gyte G: Primary postpartum hemorrhage-the Northern European Registry 200-2004. *Obstet Gynecol* 2007; 110:1270-1278
13. Allen T, George R, White W, Muir H, Habib A: A double-blind, placebo-controlled trial of four fixed rate infusion regimens of phenylephrine for hemodynamic support during spinal anesthesia for cesarean delivery. *Anesth Analg* 2010; 111:1221-9
14. Allyn J, Guglielminotti J, Omnes S, Guezouli L, Egan M, Jondeau G, Longrois D, Montravers P: Marfan's syndrome during pregnancy: anesthetic management of delivery in 16 consecutive patients. *Anesth Analg* 2013; 116:392-398
15. Aneiros F, Vazquez M, Valino C, Taboada M, Sabate S, Otero P, Costa J, Carceller J, Vazquez R, Diaz-Vieito M, Rodriguez A, Alvarez J: Does epidural versus combined

- spinal-epidural analgesia prolong labor and increase the risk of instrumental and cesarean delivery in nulliparous women? *J Clin Anesth* 2009; 21:94-97
- 16. Anstey JT, Sheldon GW, Blythe JG: Infectious morbidity after primary cesarean sections in a private institution. *Am J Obstet Gynecol* 1980; 136:205-210
 - 17. Axemo P, Fu X, Lindberg B, Ulmsten U, Wessen A: Intravenous nitroglycerin for rapid uterine relaxation. *Acta Obstetricia et Gynecologica Scandinavica*. 1998;77:50–53
 - 18. Aya AG, Vialles N, Tanoubi I, Mangin R, Ferrer JM, Robert C, Ripart J, de La Coussaye JE: Spinal anesthesia-induced hypotension: a risk comparison between patients with severe preeclampsia and healthy women undergoing preterm cesarean delivery. *Anesth Analg* 2005; 101:869-875
 - 19. Ayorinde BT, Buczkowski P, Brown J, Shah J, Buggy DJ: Evaluation of pre-emptive intramuscular phenylephrine and ephedrine for reduction of spinal anaesthesia-induced hypotension during Caesarean section. *Br J Anaesth* 2001; 86:372-376
 - 20. Bader AM, Datta S, Arthur GR, Benvenuti E, Courtney M, Hauch M: Maternal and fetal catecholamines and uterine Incision-to-delivery interval during elective cesarean. *Obstet Gynecol* 1990; 75:600-603
 - 21. Baird EJ, Arkoosh VA: Hemodynamic effects of aortocaval compression and uterine contractions in a parturient with left ventricular outflow tract obstruction. *Anesthesiology* 2012; 117:879
 - 22. Bajorek J, Kornacki Z, Ronin Walknowska E: Influence of the method of anesthesia for cesarean section on the clinical state of the newborn. *Mater Med Pol* 1972; 4:28-31
 - 23. Bang E, Lee H, Kang Y, Cho K, et al. Onset of labor epidural analgesia with ropivacaine and a varying dose of fentanyl: a randomized controlled trial *Int J Obstet Anesth* 2012; 21:45-50.
 - 24. Barrilleaux PS, Adair D, Johnson G, Lewis DF: Splenic rupture associated with severe preeclampsia. A case report. *J Reprod Med* 1999; 44:899-901
 - 25. Bateman BT, Bansil P, Hernandez-Diaz S, Mhyre JM, Callaghan WM, Kuklina EV: Prevalence, trends, and outcomes of chronic hypertension: a nationwide sample of delivery admissions. *Am J Obstet Gynecol*. 2012; 206(2):134.e1-8
 - 26. Behforouz N, Dounas M, Benhamou D: Epidural anaesthesia for caesarean delivery in triple and quadruple pregnancies. *Acta Anaesthesiol Scand* 1998; 42:1088-1091
 - 27. Belfrage P, Irestedt L, Raabe N, Arner S: General anaesthesia or lumbar epidural block for caesarean section? Effects on the foetal heart rate. *Acta Anaesthesiol Scand* 21:67-70, 1977
 - 28. Ben Aissia N, Batar S, Sadfi A, Bouhaja B, Haddad Ben Ammar MS, Gara MF: The obstetrical advantages of epidural analgesia in a trial of labor. *Tunis Med* 2004 82:37-40
 - 29. Blanche R, Eisenach JC, Tuttle R, Dewan DM: Previous wet tap does not reduce success rate of labor epidural analgesia. *Anesth Analg* 1994; 79:291-294
 - 30. Bloom SL, Spong CY, Weiner SJ, Landon MB, Rouse DJ, Varner MW, Moawad AH, Caritis SN, Harper M, Wapner RJ, Sorokin Y, Miodovnik M, O'Sullivan MJ, Sibai B, Langer O, Gabbe SG: Complications of anesthesia for cesarean delivery. *Obstet Gynecol* 2005; 106:281-287
 - 31. Bobrowski RA, Bottoms SF: Underappreciated risks of the elderly multipara. *Am J Obstet Gynecol* 1995; 172:1764-1767
 - 32. Boehm FH, Woodruff LF, Jr., Growdon JH, Jr: The effect of lumbar epidural anesthesia on fetal heart rate baseline variability. *Anesth Analg* 1975; 54:779-782

33. Bofill JA, Vincent RD, Road EI, Martin RW, Norman PF, Werham CF, Morrison JC: Nulliparous active labor, epidural analgesia, and cesarean delivery for dystocia. *Am J Obstet Gyn* 1997; 177:1465-1470
34. Boselli E, Debon R, Cimino Y, Rimmele T, Allaouchiche B, Chassard D: Background infusion is not beneficial during labor patient-controlled analgesia with 0.1% ropivacaine plus 0.5 microg/ml sufentanil. *Anesthesiology* 2004; 100: 968-72
35. Boutros A, Blary S, Bronchard R, Bonnet F: Comparison of intermittent epidural bolus, continuous epidural infusion and patient controlled-epidural analgesia during labor. *Int J Obstet Anesth* 1999; 8:236-241
36. Boyle R: Caesarean section anaesthesia and the Apgar score. *Aust N Z J Obstet Gynaecol* 1993; 33:282-284
37. Boys JE: Failed intubation in obstetric anaesthesia. A case report. *Br J Anaesth* 1983; 55:187-188
38. Bremerich DH, Waibel HJ, Mierdl S, Meininger D, Byhahn C, Zwissler BC, Ackermann HH: Comparison of continuous background infusion plus demand dose and demand-only parturient-controlled epidural analgesia (PCEA) using ropivacaine combined with sufentanil for labor and delivery. *Int J Obstet Anesth* 2005; 14:114-120
39. Brogly N, Schiraldi R, Vazquez B, Perez J, Guasch E, Gilsanz F: A randomized control trial of patient-controlled epidural analgesia (PCEA) with and without a background infusion using levobupivacaine and fentanyl. *Minerva Anestesiol* 2011;77:1149-54
40. Bullarbo M, Bokström H, Lilja H, Almström E, Lassenius N, Hansson A, Ekerhovd E: Nitroglycerin for management of retained placenta: a multicenter study. *Obstet Gynecol Int.* 2012;2012:321207
41. Bullarbo M, Tjugum J, Ekerhovd E: Sublingual nitroglycerin for management of retained placenta. *Int J Gynaecol Obstet* 2005; 91:228-232
42. Bursac D, Kulas T, Persec J, Persec Z, Duić Z, Partl JZ, Glavić Z, Hrgović Z, Bojanić K: Pregnancy and vaginal delivery in epidural analgesia in woman with cerebrospinal fluid shunt. *Coll Antropol* 2013; 37:1343-1345
43. Caldwell LE, Rosen MA, Shnider SM: Subarachnoid morphine and fentanyl for labor analgesia. Efficacy and adverse effects. *Reg Anesth* 1994; 19:2-8
44. Camann WR, Denney RA, Holby ED, Datta S: A comparison of intrathecal, epidural, and intravenous sufentanil for labor analgesia. *Anesthesiology* 1992; 77:884-887
45. Camann WR, Hortvet LA, Hughes N, Bader AM, Datta S: Maternal temperature regulation during extradural analgesia for labour. *Br J Anaesth* 1991; 67:565-568
46. Campbell DC, Camann WR, Datta S: The addition of bupivacaine to intrathecal sufentanil for labor analgesia. *Anesth Analg* 1995; 81:305-309
47. Caritis SN, Abouleish E, Edelstone DI, Mueller Heubach E: Fetal acid-base state following spinal or epidural anesthesia for cesarean section. *Obstet Gynecol* 1980; 56:610-615
48. Carlsson C, Nybell-Lindahl G, Ingemarsson I: Extradural block in patients who have previously undergone caesarean section. *Br J Anaesth* 1980; 52:827-830
49. Carvalho B, Mercier FJ, Riley ET, Brummel C, Cohen SE: Hetastarch co-loading is as effective as pre-loading for the prevention of hypotension following spinal anesthesia for cesarean delivery. *Int J Obstet Anesth* 2009; 18:150-155
50. Cassidy B, Cyna AM: Challenges that opioid-dependent women present to the obstetric anaesthetist. *Anaesth Intensive Care* 2004; 32 494-501

51. Celleno D, Capogna G: Epidural fentanyl plus bupivacaine 0.125 per cent for labour: analgesic effects. *Can J Anaesth* 1988; 35:375-378
52. Cesarini M, Torrielli R, Lahaye F, Mene JM, Cabiro C: Sprotte needle for intrathecal anaesthesia for caesarean section: incidence of postdural puncture headache. *Anaesthesia* 1990; 45:656-658
53. Chadwick LS, Vohra A: Anaesthesia for emergency Caesarean section using the Brain laryngeal mask airway. *Anaesthesia* 1989; 44:261-262
54. Chambers WA, Mowbray A, Wilson J: Extradural morphine for the relief of pain following caesarean section. *Br J Anaesth* 1983, 55:1201-1203
55. Chan AS, Ananthanarayan C, Rolbin SH: Alternating nitroglycerin and syntocinon to facilitate uterine exploration and removal of an adherent placenta. *Can J Anaesth* 1995; 42:335-337
56. Chedraui PA, Insuasti DF: Intravenous nitroglycerin in the management of retained placenta. *Gynecol Obstet Invest* 2003; 56:61-64
57. Chen Sheng-Huan, Liou Shiue-Chin, Hung Chao-Tsen, Shih Ming-Hung, Chen Chit, Tsai Shih-Chang, Tseng Chi-Hao, Wong Shu-Yam: Comparison of patient-controlled epidural analgesia and continuous epidural infusion for labor analgesia. *Chang Gung Med J* 2006; 29:576-82
58. Chestnut DH, McGrath JM, Vincent RD, Jr., Penning DH, Choi WW, Bates JN, McFarlane C: Does early administration of epidural analgesia affect obstetric outcome in nulliparous women who are in spontaneous labor? *Anesthesiology* 1994; 80:1201-1208
59. Chestnut DH, Owen CL, Bates JN, Ostman LG, Choi WW, Geiger MW: Continuous infusion epidural analgesia during labor: a randomized, double-blind comparison of 0.0625% bupivacaine/0.0002% fentanyl versus 0.125% bupivacaine. *Anesthesiology* 1988; 68:754-759
60. Chestnut DH, Vincent Jr RD, McGrath JM, Choi WW, Bates JN: Does early administration of epidural analgesia affect obstetric outcome in nulliparous women who are receiving intravenous oxytocin? *Anesthesiology* 1994; 80:1193-2000
61. Chetty V, Amanor-Boadu SD, Tora E, Andrada J, Goneyale S, Niumataiwalu K: Case report: difficult spinal anaesthesia for caesarian section in two obese pregnant patients. *West Afr J Med* 2001; 20:274-276
62. Choi DH, Ahn HJ, Kim JA. Combined low-dose spinal-epidural anesthesia versus single-shot spinal anesthesia for elective cesarean delivery. *Int J of Obstet Anesth* 2006; 15:13-17
63. Choi DH, Kim JA, Chung IS: Comparison of combined spinal epidural anesthesia and epidural anesthesia for cesarean section. *Acta Anaesthesiol Scand* 2000; 44:214-219
64. Choi DH, Park NK, Cho HS, Hahn TS, Chung IS: Effects of epidural injection on spinal block during combined spinal and epidural anesthesia for cesarean delivery. *Reg Anesth Pain Med* 2000; 25:591-595
65. Clark RB, Thompson DS, Thompson CH: Prevention of spinal hypotension associated with Cesarean section. *Anesthesiology* 1976; 45:670-674
66. Cohen S, Pantuck CB, Amar D, Burley E, Pantuck EJ: The primary action of epidural fentanyl after cesarean delivery is via a spinal mechanism. *Anesth Analg* 2002; 94:674-679

67. Cohen SE, Jasson J, Talafre ML, Chauvelot Moachon L, Barrier G: Does metoclopramide decrease the volume of gastric contents in patients undergoing cesarean section? *Anesthesiology* 61:604-607, 1984
68. Cohen SE, Tan S, White PF: Sufentanil analgesia following cesarean section: Epidural versus intravenous administration. *Anesthesiology* 1988; 68:129-134
69. Cohen SE, Woods WA: The role of epidural morphine in the postcesarean patient: Efficacy and effects on bonding. *Anesthesiology* 1983; 58:500-504
70. Collins KM, Bevan DR, Beard RW: Fluid loading to reduce abnormalities of fetal heart rate and maternal hypotension during epidural analgesia in labour. *Br Med J* 1978; 2:1460-1461
71. Collis RE, Plaat FS, Morgan BM: Comparison of midwife top-ups, continuous infusion and patient-controlled epidural analgesia for maintaining mobility after a low-dose combined spinal-epidural. *Br J Anaesth* 1999; 82:233-236
72. Combs CA, Laros RK, Jr: Prolonged third stage of labor: morbidity and risk factors. *Obstet Gynecol* 1991; 77:863-867
73. Combs CA, Murphy EL, Laros RK, Jr: Factors associated with hemorrhage in cesarean deliveries. *Obstet Gynecol* 1991; 77:77-82
74. Comparative Obstetric Mobile Epidural Trial (COMET) Study Group UK: Effects of low-dose mobile versus traditional epidural techniques on mode of delivery: a randomized controlled trial. *Lancet* 2001; 358:19-23
75. Comparative Obstetric Mobile Epidural Trial (COMET) Study Group UK: Randomized controlled trial comparing traditional with two "mobile" epidural techniques: anesthetic and analgesic efficacy. *Anesthesiology* 2002; 97:1567-1575
76. Cooke RD, Comyn DJ, Ball RW: Prevention of postoperative nausea and vomiting by domperidone: A double-blind randomized study using domperidone, metoclopramide and a placebo. *S Afr Med J* 1979; 56:827-829
77. Cooper D, Schofield L, Hynd R, Selvan D, Lloyd A, Meed T, Winnard J: Prospective evaluation of systolic arterial pressure control with a phenylephrine infusion regimen during spinal anaesthesia for caesarean section. *Int J Obstet Anesth* 2012; 21:245-252
78. Cooper D, Sharma S, Orakkan P, Gurung S: Retrospective study of association between choice of vasopressor given during spinal anaesthesia for high-risk caesarean delivery and fetal pH. *Int J Obstet Anesth* 2010; 19:44-49
79. Cooper DW, Carpenter M, Mowbray P, Desira WR, Ryall DM, Kokri MS: Fetal and maternal effects of phenylephrine and ephedrine during spinal anesthesia for cesarean delivery. *Anesthesiology* 2002; 97:1582-1590
80. Cooper DW, Jeyaraj L, Hynd R, Thompson R, Meek T, Ryall DM, Kokri MS: Evidence that intravenous vasopressors can affect rostral spread of spinal anesthesia in pregnancy. *Anesthesiology* 2004; 101:28-33
81. Cooper G, MacArthur C, Wilson M, Moore P, Shennan A (COMET Study Group): Satisfaction, control and pain relief: short-and long-term assessments in a randomised controlled trial of low-dose and traditional epidurals and a non-epidural comparison group. *Int J Obstet Anesth* 2010; 19:31-37
82. Cortes CAF, Sanchez CA, Oliveira AS, Sanchez FM: Labor Analgesia: a comparative study between combined spinal-epidural anesthesia versus continuous epidural anesthesia. *Rev Bras Anestesiol* 2007; 57:39-51

83. Cosmi EV, Marx GF: Acid-base status of the fetus and clinical condition of the newborn following cesarean section. *Am J Obstet Gynecol* 1968; 102:378-382
84. Cousins LM, Teplick FB, Poeltler DM: Pre-cesarean blood bank orders: a safe and less expensive approach. *Obstet Gynecol* 1996; 87:912-916
85. Crawford JS: A comparison of spinal analgesia and general anesthesia for elective cesarean section. Interim report. *Am J Obstet Gynecol* 1966; 94:858-860
86. Crites J, Ramanathan J: Acute hypoglycemia following combined spinal-epidural anesthesia (CSE) in a parturient with diabetes mellitus. *Anesthesiology* 2000; 93:591-592
87. Crosby ET: Obstetrical anaesthesia for patients with the syndrome of haemolysis, elevated liver enzymes and low platelets. *Can J Anaesth* 1991; 38:227-233
88. Curry PD, Pacsoo C, Heap DG: Patient-controlled epidural analgesia in obstetric anaesthetic practice. *Pain* 1994; 57:125-127
89. Daley MD, Sandler AN, Turner KE, Vosu H, Slavchenko P: A comparison of epidural and intramuscular morphine in patients following cesarean section. *Anesthesiology* 1990, 72:289-294
90. Danzer BI, Birnbach DJ, Stein DJ, Kuroda MM, Thys DM: Does metoclopramide supplement postoperative analgesia using patient-controlled analgesia with morphine in patients undergoing elective cesarean delivery? *Reg Anesth* 1997; 22:424-427
91. Datta S, Alper MH, Ostheimer GW, Weiss JB: Method of ephedrine administration and nausea and hypotension during spinal anesthesia for cesarean section. *Br J Anaesth* 1999; 83:475-477
92. Datta S, Brown WU, Jr. Acid-base status in diabetic mothers and their infants following general or spinal anesthesia for cesarean section. *Anesthesiology* 1977; 47:272-276
93. Datta S, Ostheimer GW, Weiss JB, Brown WU, Jr, Alper MH: Neonatal effect of prolonged anesthetic induction for cesarean section. *Obstet Gynecol* 1981; 58:331-335
94. Davies SJ, Paech MJ, Welch H, Evans SF, Pavay TJG: Maternal experience during epidural or combined spinal-epidural anesthesia for cesarean section: A prospective, randomized trial. *Anesth Analg* 1997; 85:607-613
95. de Mello WF, Kocan M: The laryngeal mask in failed intubation. *Anaesthesia* 1990; 41:689-690
96. de Vries JI, Vellenga E, Aarnoudse JG: Plasma beta-thromboglobulin in normal pregnancy and pregnancy-induced hypertension. *Eur J Obstet Gynecol Reprod Biol* 1983; 14:209-216
97. Dean LS, Rogers RP 3rd, Harley RA, Hood DD: Case scenario: amniotic fluid embolism. *Anesthesiology* 2012; 116:186-192
98. Desalu I, Kushimo OT: Is ephedrine infusion more effective at preventing hypotension than traditional prehydration during spinal anaesthesia for caesarean section in African parturients? *Int J Obstet Anesth* 2005; 14:294-299
99. Desprats R, Mandry J, Grandjean H, Amar B, Pontonnier G, Lareng L: Peridural analgesia during labor: comparative study of a fentanyl-marcaine combination and marcaine alone. *J Gynecol Obstet Biol Reprod Paris* 1983; 12:901-905
100. Devcic A, Sprung J, Patel S, Kettler R, Maitra DA: PDPH in obstetric anesthesia: comparison of 24-gauge Sprotte and 25-gauge Quincke needles and effect of subarachnoid administration of fentanyl. *Reg Anesth* 1993; 18:222-225
101. Dewan DM, Floyd HM, Thistlewood JM, Bogard TD, Spielman FJ: Sodium citrate pretreatment in elective cesarean section patients. *Anesth Analg* 1985; 64:34-37

102. Dewandre P, Decurninge V, Bonhommme V, Hans P, Brichant J: Side effects of the addition of clonidine 75 µg or sufentanil 5 µg to 0.2% ropivacaine for labour epidural analgesia. *Int J Obstet Anesth* 2010; 19:149-154
103. Dick W, Traub E, Kraus H, Tollner U, Burghard R, Muck J: General anaesthesia versus epidural anaesthesia for primary caesarean section--a comparative study. *Eur J Anaesthesiol* 1992; 9:15-21
104. Dobb G: Laryngeal oedema complicating obstetric anaesthesia. *Anaesthesia* 1978; 33:839-840
105. Downing JW, Houlton PC, Barclay A: Extradural analgesia for caesarean section: a comparison with general anaesthesia. *Br J Anaesth* 1979; 51:367-374
106. Dresner M, Bamber J, Calow C, Freeman J, Charlton P: Comparison of low-dose epidural with combined spinal-epidural analgesia for labour. *Br J Anaesth* 1999; 83:756-760
107. Dresner M, Pinder A: Anaesthesia for caesarean section in women with complex cardiac disease: 34 cases using the Braun Spinocath spinal catheter. *Int J Obstet Anesth* 2009; 18:131-136
108. Druzin ML, Stier E: Maternal platelet count at delivery in patients with idiopathic thrombocytopenic purpura, not related to perioperative complications. *J Am Coll Surg* 1994; 179:264-266
109. Dyer R, Piercy J, Reed A, Strathie G, Lombard, Anthony J, James M: Comparison between pulse waveform analysis and thermodilution cardiac output determination in patients with severe pre-eclampsia. *Br J Anaesth* 2011; 106:77-81
110. Dyer R, Reed An, vanDyk D, Arcache M, Hodges O, Lombard C, Greenwood J, James M: Hemodynamic effects of ephedrine, phenylephrine, and the coadministration of phenylephrine with oxytocin during spinal anesthesia for elective cesarean delivery. *Anesthesiology* 2009; 111:753-65
111. Dyer RA, Els I, Farbas J, Torr GJ, Schoeman LK, James MF: Prospective randomized trial comparing general with spinal anesthesia for cesarean delivery in preeclamptic patients with a nonreassuring fetal heart tract. *Anesthesiology* 2003; 99:561-569
112. Dyer RA, Farina Z, Joubert IA, DuToit P, Meyer M, Torr G, Wells K, James MFM: Crystalloid preload versus rapid crystalloid administration after induction of spinal anaesthesia (coload) for elective caesarean section. *Anaesth Intensive Care* 2004; 32:351-357
113. Edwards ND, Hartley M, Clyburn P, Harmer M: Epidural pethidine and bupivacaine in labour. *Anaesthesia* 1992; 47:435-437
114. Edwards RD, Hansel NK, Pruessner HT, Barton B: Intrathecal morphine as analgesia for labor pain. *J Am Board Fam Pract* 1988; 1:245-250
115. Edwards RD, Hansel NK, Pruessner HT, Barton B: Intrathecal morphine sulfate for labor pain. *Tex Med* 1985; 81:46-48
116. Eisenach JC, Grice SC, Dewan DM. Patient-controlled analgesia following cesarean section: a comparison with epidural and intramuscular narcotics. *Anesthesiology* 1988; 68:444-448
117. Ekerhovd E, Bullarbo M: Sublingual nitroglycerin seems to be effective in the management of retained placenta. *Acta Obstet Gynecol Scand* 2008; 87:222-225
118. El-Kerdawy H, Farouk A: Labor analgesia in preeclampsia: remifentanil patient controlled intravenous analgesia versus epidural analgesia. *MEJ Anesth* 2010; 20:539-545

119. Elliott RD: Continuous infusion epidural analgesia for obstetrics: bupivacaine versus bupivacaine-fentanyl mixture. *Can J Anaesth* 1991; 38:303-310
120. Endler GC, Mariona FG, Sokol RJ, Stevenson LB: Anesthesia-related maternal mortality in Michigan, 1972 to 1984. *Am J Obstet Gynecol* 1988; 159:187-193
121. Evans CM, Murphy JF, Gray OP, Rosen M: Epidural versus general anaesthesia for elective Caesarean section. Effect on Apgar score and acid-base status of the newborn. *Anaesthesia* 1989; 44:778-782
122. Fabris L, Maretoc A: Effects of general anaesthesia versus spinal anaesthesia for caesarean section on postoperative analgesic consumption and postoperative pain. *Period Biol* 2009; 111:251-255
123. Fanning N, Balki M, Sermer M, Colman J, Carvalho JC: Noninvasive cardiac output monitoring during general anesthesia for Cesarean delivery in a patient with severe aortic stenosis. *Can J Anaesth* 2011; 58:837-841
124. Ferguson E, Paech M, Veltman M: Hypertrophic cardiomyopathy and caesarean section: intraoperative use of transthoracic echocardiography. *Int J Obstet Anesth* 2006; 15:311-316
125. Ferouz F, Norris MC, Leighton BL: Risk of respiratory arrest after intrathecal sufentanil. *Anesth Analg* 1997; 85:1088-1090
126. Ferrante FM, Barber MJ, Segal M, Hughes NJ, Datta S: 0.0625% bupivacaine with 0.0002% fentanyl via patient-controlled epidural analgesia for pain of labor and delivery. *Clin J Pain* 1995; 11:121-126
127. Ferrante FM, Lu L, Jamison SB, Datta S: Patient-controlled epidural analgesia: demand dosing. *Anesth Analg* 1991; 73:547-552
128. Ferrante FM, Rosinia FA, Gordon C, Datta S: The role of continuous background infusions in patient-controlled epidural analgesia for labor and delivery. *Anesth Analg* 1994; 79:80-84
129. Finegold H, Darwich A, Romeo R, Vallejo M, Ramanathan S: Successful resuscitation after maternal cardiac arrest by immediate cesarean section in the labor room. *Anesthesiology* 2002; 96:1278
130. FitzGerald MP, Floro C, Siegel J, Hernandez E: Laboratory findings in hypertensive disorders of pregnancy. *J Natl Med Assoc* 1996; 88:794-798
131. Flamm BL, Dunnett C, Fischermann E, Quilligan EJ: Vaginal delivery following cesarean section: use of oxytocin augmentation and epidural anesthesia with internal tocodynamometer and internal fetal monitoring. *Am J Obstet Gynecol* 1984; 148:759-763
132. Flamm BL, Lim OW, Jones C, Fallon D, Newman LA, Mantis JK: Vaginal birth after cesarean section: results of a multicenter study. *Am J Obstet Gynecol* 1988; 158:1079-1084
133. Fox GS, Houle GL: Acid-base studies in elective Caesarean sections during epidural and general anaesthesia. *Can Anaesth Soc J* 1971; 18:60-71
134. Fox GS, Smith JB, Namba Y, Johnson RC: Anesthesia for cesarean section: further studies. *Am J Obstet Gynecol* 1979; 133:15-19
135. Freedman MA, Wilds PL, George WM: Grotesque obesity: a serious complication of labor and delivery. *South Med J* 1972; 65:732-736
136. Fujii Y, Tanaka H, Toyooka H: Prevention of nausea and vomiting with granisetron, droperidol and metoclopramide during and after spinal anaesthesia for caesarean section:

- a randomized, double-blind, placebo-controlled trial. *Acta Anaesthesiol Scand* 1998; 42:921-925
137. Gajraj NM, Wallace DH, Pace NA: Supraventricular tachycardia in a parturient under spinal anesthesia. *Reg Anesth* 1993; 18:261-263
 138. Gale R, Slater PE, Zalkinder Luboshitz I: Neonatal advantage of epidural anesthesia in elective and emergency cesarean sections: a report of 531 cases. *Eur J Obstet Gynecol Reprod Biol* 1986; 23:369-377
 139. Gambling D, Berkowitz J, Farrell T, Pue A, Shay D: A randomized controlled comparison of epidural analgesia and combined spinal-epidural analgesia in a private practice setting: pain scores during first and second stages of labor and delivery. *Anesth Anal* 2013; 116:636-43
 140. Gambling DR, Huber CJ, Berkowitz J, Howell P, Swenerton JE, Ross PL, Crochetiere CT, Pavay TJ: Patient-controlled epidural analgesia in labour: varying bolus dose and lockout interval. *Can J Anaesth* 1993; 40:211-217
 141. Gambling DR, Yu P, Cole C, McMorland GH, Palmer L: A comparative study of patient controlled epidural analgesia (PCEA) and continuous infusion epidural analgesia (CIEA) during labour. *Can J Anaesth* 1988; 35:249-254
 142. Ghosh AK, Tipton RH: Early postpartum tubal ligation under epidural analgesia. *Br J Obstet Gynaecol* 1976; 83:731-732
 143. Gilbert WM, Towner DR, Field NT, Anthony J: The safety and utility of pulmonary artery catheterization in severe preeclampsia and eclampsia. *Am J Obstet Gynecol* 2000; 182:1397-1403
 144. Gilstrap LC, III, Hauth JC, Hankins GD, Patterson AR: Effect of type of anesthesia on blood loss at cesarean section. *Obstet Gynecol* 1987; 69:1328-1332
 145. Godley M, Reddy AR: Use of LMA for awake intubation for caesarean section. *Can J Anaesth* 1996; 43:299-302
 146. Gomez P, Echevarria M, Calderon J, Caba F, Martinez A, Rodriguez R: The efficacy and safety of continuous epidural analgesia versus intradural-epidural analgesia during labour. *Revista Espanola de Anestesiologia y Reanimacion* 2001; 48:217-222
 147. Goodall PT, Ahn JT, Chapa JB, Hibbard JU: Obesity as a risk factor for failed trial of labor in patients with previous cesarean delivery. *Am J Obstet Gynecol* 2005; 192:1423-1426
 148. Greenhalgh CA: Respiratory arrest in a parturient following intrathecal injection of sufentanil and bupivacaine. *Anesthesia* 1996; 51:173-175
 149. Greer IA, Lowe GD, Walker JJ, Forbes CD: Haemorrhagic problems in obstetrics and gynaecology in patients with congenital coagulopathies. *Br J Obstet Gynaecol* 1991; 98:909-918
 150. Grotegut CA1, Kuklina EV, Anstrom KJ, Heine RP, Callaghan WM, Myers ER, James AH: Factors associated with the change in prevalence of cardiomyopathy at delivery in the period 2000-2009: a population-based prevalence study. *BJOG* 2014; 121:1386-1394
 151. Gutsche BB: Prophylactic ephedrine preceding spinal analgesia for cesarean section. *Anesthesiology* 1976; 45:462-465
 152. Hall PA, Bennett A, Wilkes MP, Lewis M: Spinal anaesthesia for caesarean section: comparison of infusions of phenylephrine and ephedrine. *Br J Anaesth* 1994; 73:471-474
 153. Harrison DM, Sinatra R, Morgese L, Chung JH: Epidural narcotic and patient-controlled analgesia for post-cesarean section pain relief. *Anesthesiology* 1988; 68:454-457

154. Hasham FM, Andrews PJD, Juneja MM, Ackermann III WE: The laryngeal mask airway facilitates intubation at cesarean section: a case report of difficult intubation. *Int J Obstet Anesth* 1993; 2:181-182
155. Hawkins JL, Hess KR, Kubicek MA, Joyce TH, III, Morrow DH: A reevaluation of the association between instrument delivery and epidural analgesia. *Reg Anesth* 1995; 20:50-56
156. Hawksworth CR, Purdie J: Failed combined spinal epidural then failed intubation at an elective caesarean section. *Hosp Med* 1998; 59:173
157. Haydon ML, Larson D, Reed E, Shrivastava V, Preslicka C, Nageotte M: Obstetric outcomes and maternal satisfaction in nulliparous women using patient-controlled epidural analgesia. *Am J Obstet Gynecol* 2011; 205:271
158. Helbo Hansen S, Bang U, Garcia RS, Olesen AS, Kjeldsen L: Subarachnoid versus epidural bupivacaine 0.5% for caesarean section. *Acta Anaesthesiol Scand* 1988; 32:473-476
159. Henderson SK, Matthew E, Cohen H, Avram MJ: Epidural hydromorphone: A double blind comparison with intramuscular hydromorphone for postcesarean section analgesia. *Anesthesiology* 1987; 66:825-830
160. Hepner DL, Gaiser RR, Cheek TG, Gutsche BB: Comparison of combined spinal-epidural and low dose epidural for labour analgesia. *Can J Anaesth* 2000; 47:232-236
161. Hepner DL, Tsen LC: Severe thrombocytopenia, type 2B von Willebrand disease and pregnancy. *Anesthesiology* 2004; 101:1465-1467
162. Herpolsheimer A, Schretenthaler J: The use of intrapartum intrathecal narcotic analgesia in a community-based hospital. *Obstet Gynecol* 1994; 84:931-936
163. Hill JB, Alexander JM, Sharma SK, McIntire DD, Leveno KJ: A comparison of the effects of epidural and meperidine analgesia during labor on fetal heart rate. *Obstet Gynecol* 2003; 102:333-337
164. Hilton G, Mihm F, Butwick A: Anesthetic management of a parturient with VACTERL association undergoing Cesarean delivery. *Can J Anaesth* 2013; 60:570-576
165. Hinchliffe D, Norris A: Management of failed intubation in a septic parturient. *Br J Anaesth* 2002; 89:328-330
166. Hingson R, Gould JB, Morelock S, Kayne H, Heeren T, Alpert JJ, Zuckerman B, Day N: Maternal cigarette smoking, psychoactive substance use, and infant Apgar scores. *Am J Obstet Gynecol* 1982; 144:959-966
167. Hjertberg R, Belfrage P, Hagnvik K: Hemodynamic measurements with Swan-Ganz catheter in women with severe proteinuric gestational hypertension (pre-eclampsia). *Acta Obstet Gynecol Scand* 1991; 70:193-198
168. Hodgkinson R, Bhatt M, Kim SS, Grewal G, Marx GF: Neonatal neurobehavioral tests following cesarean section under general and spinal anesthesia. *Am J Obstet Gynecol* 1978; 132:670-674
169. Hodgkinson R, Husain FJ: Caesarean section associated with gross obesity. *Br J Anaesth* 1980; 52:919-923
170. Hodgson CA, Wauchob TD: A comparison of spinal and general anaesthesia for elective Caesarean section: effect on neonatal condition at birth. *Int J Obstet Anesth* 1994; 3:25-30

171. Hoffman SL, Zaphiratos V, Girard MA, Boucher M, Crochetière C: Failed epidural analgesia in a parturient with advanced ankylosing spondylitis: a novel explanation. *Can J Anaesth* 2012; 59:871-874
172. Hollm'en AI, Eskelinen P, Tolonen U, Arranto J, Sulg I, Jouppila R, Jouppila P: Effects of anaesthesia for caesarean section on the computerized EEG of the neonate. *Eur J Anaesthesiol* 1985; 2:39-51
173. Hollm'en AI, Jouppila R, Koivisto M, Maatta L, Pihlajaniemi R, Puukka M, Rantakyla P: Neurologic activity of infants following anesthesia for cesarean section. *Anesthesiology* 1978; 48:350-356
174. Hong JY, Jee YS, Yoon HJ, Kim SM: Comparison of general and epidural anesthesia in elective cesarean section for placenta previa totalis: maternal hemodynamics, blood loss and neonatal outcome. *Int J Obstet Anesth* 2003; 12:12-16
175. Hood DD, Curry R: Spinal versus epidural anaesthesia for Caesarean section in severely pre-eclamptic patients. A retrospective study. *Anesthesiology* 1999; 90:1276-1282
176. Hood DD, Dewan DM: Anesthetic and obstetric outcome in morbidly obese parturients. *Anesthesiology* 1993; 79:1210-1218
177. Huang C, Fan Y, Tsai P: Differential impacts of modes of anaesthesia on the risk of stroke among preeclamptic women who undergo caesarean delivery: a population-based study. *Br J Anaesth* 2010; 105:818-26
178. Hunt CO, Naulty JS, Malinow AM, Datta S, Ostheimer GW: Epidural butorphanol-bupivacaine for analgesia during labor and delivery. *Anesth Analg* 1989; 68:323-327
179. Husaini SW, Russell IF: Volume preload: lack of effect in the prevention of spinal-induced hypotension at caesarean section. *Int J Obstet Anesth* 1998; 7:76-81
180. Hussain S, Khan RA, Iqbal M, Shafiq M, Khan FA: A comparison of the effects of erythromycin and metoclopramide on gastric fluid volume and pH patients undergoing elective caesarean section. *Anaesth Pain Intensive Care* 2011; 15:148-152
181. Hwa HL, Chen RJ, Chen YC, Wang TR, Huang SC, Chow SN: Maternal and fetal outcome of pregnant women with idiopathic thrombocytopenic purpura: retrospective analysis of 25 pregnancies. *J Formos Med Assoc* 1993; 92:957-961
182. Hwang JJ, Ho ST, Wang JJ, Liu HS: Post dural puncture headache in cesarean section: comparison of 25-gauge Whitacre with 25- and 26-gauge Quincke needles. *Acta Anaesthesiol Sin* 1997; 35:33-37
183. Irestedt L, Lagercrantz H, Hjemdahl P, Hagenvik K, Belfrage P: Fetal and maternal plasma catecholamine levels at elective cesarean section under general or epidural anesthesia versus vaginal delivery. *Am J Obstet Gynecol* 1982; 142:1004-1010
184. Ishiyama T, Oguchi T, Iijima T, Matsukawa T, Kashimoto S, Kumazawa T: Combined spinal and epidural anesthesia for cesarean section in a patient with hypertrophic obstructive cardiomyopathy. *Anesth Analg* 2003; 96:629-630
185. Ismail M, Hassanin M: Neuraxial analgesia versus intravenous remifentanil for pain relief in early labor in nulliparous women. *Arch Gynecol Obstet* 2012; 286:1375-1381
186. Ismail S, Huda A: An observational study of anaesthesia and surgical time in elective caesarean section: spinal compared with general anaesthesia. *Int J Obstet Anesth* 2009; 18:352-355
187. Jacob JJ WA, Verghese M, Afzal L: Crystalloid preload versus crystalloid coload for parturients undergoing cesarean section under spinal anaesthesia. *J Obstet Anaesth Crit Care* 2012; 2:10-15

188. James FM, III, Crawford JS, Hopkinson R, Davies P, Naiem H: A comparison of general anesthesia and lumbar epidural analgesia for elective cesarean section. *Anesth Analg* 1977; 56:228-235
189. James KS, McGrady E, Quasim I, Patrick A: Comparison of epidural bolus administration of 0.25% bupivacaine and 0.1% bupivacaine with 0.0002% fentanyl for analgesia during labour. *Br J Anaesth* 1998; 81:507-510
190. Jasson J, Lefevre G, Tallet F, Talafre ML, Legagneux F, Conseiller C: Oral sodium citrate before general anesthesia for elective cesarean section. Effects on pH and volume of gastric content. *Ann Fr Anesth Reanim* 1989; 8:12-18
191. Jayaram A, Bowen MP, Deshpande S, Carp HM: Ultrasound examination of the stomach contents of women in the postpartum period. *Anesth Analg* 1997; 84:522-526
192. Johnson C, Oriol N, Flood K: Trial of labor: a study of 110 patients. *J Clin Anesth* 1991; 3:216-218
193. Jouppila P, Jouppila R, Kaar K, Merila M: Fetal heart rate patterns and uterine activity after segmental epidural analgesia. *Br J Obstet Gynaecol* 1977; 84:481-486
194. Juneja MM, Ackerman WE, Kaczorowski DM, Sollo DG, Gunzenhauser LF: Continuous epidural lidocaine infusion in the parturient with paroxysmal ventricular tachycardia. *Anesthesiology* 1989; 71:305-308
195. Juul J, Lie B, Friberg Nielsen S: Epidural analgesia vs. general anesthesia for cesarean section. *Acta Obstet Gynecol Scand Suppl* 1988; 67:203-206
196. Kain ZN, Mayes LC, Ferris CA, Pakes J, Schottenfeld R: Cocaine-abusing parturients undergoing cesarean section: A cohort study. *Anesthesiology* 1996; 85:1028-1035
197. Kamat SK, Shah MV, Chaudhary LS, Pandya S, Bhatt MM: Effect of induction-delivery and uterine-delivery on apgar scoring of the newborn. *J Postgrad Med* 1991; 37:125-127
198. Kamenik M, Paver-Erzen V: The effects of lactated Ringer's solution infusion on cardiac output changes after spinal anesthesia. *Anesth Analg* 2001; 92:710-714
199. Karaman S, Akercan F, Akarsu T, Firat V, Ozcan O, Karadadas N: Comparison of the maternal and neonatal effects of epidural block and of combined spinal-epidural block for cesarean section. *Eur J Obstet Gynecol Reprod Biol* 2005; 121:18-23
200. Kartawiadi L, Vercauteren MP, van Steenberge AL, Adriaensen HA: Spinal analgesia during labor with low-dose bupivacaine, sufentanil, and epinephrine. A comparison with epidural analgesia. *Reg Anesth* 1996; 21:191-196
201. Kato R, Shimamoto H, Terui K, Yokota K, Miyao H: Delayed respiratory depression associated with 0.15mg intrathecal morphine for cesarean section: a review of 1915 cases. *J Anesth* 2008; 22:112-116
202. Kato R, Terui K, Yokota K, Nakagawa C, Uchida J, Miyao H. Anesthetic management for cesarean section in moyamoya disease: a report of five consecutive cases and a mini-review. *Int J Obstet Anesth* 2006; 15:152-158
203. Katsiris S, Williams S, Leighton BL, Halpern S: Respiratory arrest following intrathecal injection of sufentanil and bupivacaine in a parturient. *Can J Anaesth* 1998; 45:880-883
204. Kavak ZN, Basgul A, Ceyhan N: Short-term outcome of newborn infants: Spinal versus general anesthesia for elective cesarean section: a prospective randomized study. *Eur J Obstet Reprod Biol* 2001; 100:50-54
205. Kehl F, Erfkamp S, Roewer N: Respiratory arrest during caesarean section after intrathecal administration of sufentanil in combination with 0.1% bupivacaine 10 ml. *Anaesth Intensive Care* 2002; 30:698-699

206. Keller C, Brimacombe J, Lirk P, Puhringer F: Failed obstetric tracheal intubation and postoperative respiratory support with the ProSeal laryngeal mask airway. *Anesth Analg* 2004; 98:1467-1470
207. Kessous R, Weintraub AY, Wiznitzer A, Zlotnik A, Pariente G, Polachek H, Press F, Aricha-Tamir B, Leizerovich A, Sheiner E: Spinal versus general anesthesia in cesarean sections: the effects on postoperative pain perception. *Arch Gynecol Obstet* 2012; 286:75-79
208. Khan M, ul-Nisai W, Farooqi A, Ahmad N, Qaz S: Crystalloid co-load: a better option than crystalloid pre-load for prevention Of postspinal hypotension in elective caesarean section. *Internet J Anesthesiol*. 2013; 32: <https://ispub.com/IJA/32/1/1503#>. Accessed July 8, 2015.
209. King M, Wrench I, Galimberti A, Spray R: Introduction of cell salvage to a large obstetric unit: the first six months. *Int J Obstet Anesth* 2009; 18:111-117
210. King SW, Rosen MA: Prophylactic ephedrine and hypotension associated with spinal anesthesia for cesarean delivery. *Int J Obstet Anesth* 1998; 7:18-22
211. Kinney MA, Rose CH, Traynor KD, Deutsch E, Memon HU, Tanouye S, Arendt KW, Hebl JR: Emergency bedside cesarean delivery: lessons learned in teamwork and patient safety. *BMC Res Notes* 2012; 5:412
212. Kluger MT: Ephedrine may predispose to arrhythmias in obstetric anaesthesia. *Anaesth Intensive Care* 2000; 28:336
213. Knottnerus JA, Delgado LR, Knipschild PG, Essed GG, Smits F: Haematologic parameters and pregnancy outcome. A prospective cohort study in the third trimester. *J Clin Epidemiol* 1990; 43:461-466
214. Kolatat T, Somboonnanonda A, Lertakyamanee J, Chinachot T, Tritrakarn T, Muangkasem J: Effects of general and regional anesthesia on the neonate (a prospective, randomized trial). *J Med Assoc Thai* 1999; 82:40-45
215. Lambert DH, Hurley RJ, Hertwig L, Datta S: Role of needle gauge and tip configuration in the production of lumbar puncture headache. *Reg Anesth* 1997; 22:66-72
216. Landry DP, Bennett FM, Oriol NE: Analysis of heart rate dynamics as a measure of autonomic tone in obstetrical patients undergoing epidural or spinal anesthesia. *Reg Anesth* 1994; 19:189-195
217. Langesaeter E, Dragsund M, Rosseland L: Regional anaesthesia for a caesarean section in women with cardiac disease: a prospective study. *Acta Anesthesia Scand* 2010; 54:46-54
218. Langesaeter E, Rosseland L, Stubhaug A: Continuous invasive blood pressure and cardiac output monitoring during cesarean delivery. *Anesthesiology* 2008; 109:856-63
219. LaPorta RF, Arthur GR, Datta S: Phenylephrine in treating maternal hypotension due to spinal anaesthesia for caesarean delivery: effects on neonatal catecholamine concentrations, acid base status and Apgar scores. *Acta Anaesth Scand* 1995; 39:901-905
220. Larsen R, Titlestad K, Lillevang ST, Thomsen SG, Kidholm K, Georgsen J: Cesarean section: is pretransfusion testing for alloantibodies necessary? *Acta Obstet Gynecol Scand* 2005; 84:448-455
221. Ledin Eriksson S, Gentile C, Olofsson CH: PCEA compared to continuous epidural infusion in an ultra-low-dose regimen for labor pain relief: a randomized study. *Acta Anaesthesiol Scand* 2003; 47:1085-1090
222. Leduc L, Wheeler JM, Kirshon B, Mitchell P, Cotton DB: Coagulation profile in severe preeclampsia. *Obstet Gynecol* 1992; 79:14-18

223. Lee BB, Ngan Kee WD, Ng FF, Lau TK, Wong EL: Epidural infusions for labor analgesia: a comparison of 0.2% ropivacaine, 0.1% ropivacaine, and 0.1% ropivacaine with fentanyl. *Reg Anesth Pain Med* 2002; 27:31-36
224. Lee SY, Choi DH, Park HW: The effect of colloid co-hydration on the use of phenylephrine and hemodynamics during low-dose combined spinal-epidural anesthesia for cesarean delivery. *Korean J Anesthesiol* 2008; 55:685-690
225. Leffert LR1, Clancy CR, Bateman BT, Bryant AS, Kuklina EV: Hypertensive disorders and pregnancy-related stroke: frequency, trends, risk factors, and outcomes. *Obstet Gynecol* 2015; 125:124-131
226. Lehmann KA, Stern S, Breuker KH: Obstetrical peridural anesthesia with bupivacaine and buprenorphine. A randomized double-blind study in comparison with untreated controls. *Anaesthetist* 1992; 41:414-422
227. Lertakyamanee J, Chinachoti T, Tritrakarn T, Muangkasem J, Somboonnanonda A, Kolatat T: Comparison of general and regional anesthesia for cesarean section: success rate, blood loss and satisfaction from a randomized trial. *J Med Assoc Thai* 1999; 82: 672-680
228. Liang CC, Chang SD, Wong SY, Chang YL, Cheng PJ: Effects of postoperative analgesia on postpartum urinary retention in women undergoing cesarean delivery. *J Obstet Gynaecol Res* 2010; 36:991-995
229. Lieberman E, Lang JM, Cohen A, D'Agostino R, Jr., Datta S, Frigoletto FD, Jr: Association of epidural analgesia with cesarean delivery in nulliparas. *Obstet Gynecol* 1996; 88:993-1000
230. Lilker S, Meyer, R, Downey K, Macarthur A: Anesthetic considerations for placenta accreta. *Int J Obstet Anesth* 2011; 20:288-292
231. Lim SK, Elegbe EO: The use of single dose of sodium citrate as a prophylaxis against acid aspiration syndrome in obstetric patients undergoing caesarean section. *Med J Malaysia* 1991; 46:349-355
232. Lim Y, Sia AT, Ocampo CE: Comparison of computer integrated patient controlled epidural analgesia vs. conventional patient controlled epidural analgesia for pain relief in labour. *Anaesthesia* 2006; 61:339-344
233. Lin CJ, Huang CL, Hsu HW, Chen TL: Prophylaxis against acid aspiration in regional anesthesia for elective cesarean section: a comparison between oral single-dose ranitidine, famotidine and omeprazole assessed with fiberoptic gastric aspiration. *Acta Anaesthesiol Sin* 1996; 34:179-184
234. Lindblad A, Bernow J, Mars'al K: Obstetric analgesia and fetal aortic blood flow during labour. *Br J Obstet Gynaecol* 1987; 94:306-311
235. Lirzin JD, Jacquinot P, Dailland P, Jorrot JC, Jasson J, Talafre ML, Conseiller C: Controlled trial of extradural bupivacaine with fentanyl, morphine or placebo for pain relief in labour. *Br J Anaesth* 1989; 62:641-644
236. Loughnan BA, Carli F, Romney M, Dore CJ, Gordon H: Epidural analgesia and backache: a randomized controlled comparison with intramuscular meperidine for analgesia during labour. *Br J Anaesth* 2002; 89:466-472
237. Loughnan BA, Carli F, Romney M, Doré CJ, Gordon H: Randomized controlled comparison of epidural bupivacaine versus pethidine for analgesia in labour. *Br J Anaesth* 2000; 84:715-719
238. Loughran PG, Moore J, Dundee JW: Maternal stress response associated with caesarean delivery under general and epidural anaesthesia. *Br J Obstet Gynaecol* 1986; 93:943-949

239. Loughrey JP, Walsh F, Gardiner J: Prophylactic intravenous bolus ephedrine for elective Caesarean section under spinal anaesthesia. *Eur J Anaesthesiol* 2002; 19:63-68
240. Low JA, Pancham SR, Worthington D, Boston RW: The incidence of fetal asphyxia in six hundred high-risk monitored pregnancies. *Am J Obstet Gynecol* 1975; 121:456-459
241. Lowenwirt IP, Zauk RM, Handwerker SM: Safety of intravenous glyceryl trinitrate in management of retained placenta. *Aust N Z J Obstet Gynaecol* 1997; 37:20-24
242. Lucas MJ, Sharma SK, McIntire DD, Wiley J, Sidawi JE, Ramin SM, Leveno KJ, Cunningham EG: A randomized trial of labor analgesia in women with pregnancy-induced hypertension. *Am J Obstet Gynecol* 2001; 185:970-975
243. Lussos SA, Bader AM, Thornhill ML, Datta S: The antiemetic efficacy and safety of prophylactic metoclopramide for elective cesarean delivery during spinal anesthesia. *Reg Anesth* 1992; 17:126-130
244. Luxman E, Wolman I, Groutz A, Cohen JR, Lottan M, Pauzner D, David MP: The effect of early epidural block administration on the progression and outcome of labor. *Int J Obstet Anesth* 1998; 7:161-164
245. Lyra TG, Pinto VA, Ivo FA, Nascimento Jdos S: Osteogenesis imperfecta in pregnancy. Case report. *Rev Bras Anestesiol* 2010; 60:321-324
246. Lysak SZ, Eisenach JC, Dobson CE II: Patient-controlled epidural analgesia during labor: a comparison of three solutions with a continuous infusion control. *Anesthesiology* 1990; 72:44-49
247. Macrae DJ, Munishankrappa S, Burrow LM, Milne MK, Grant IS: Double-blind comparison of the efficacy of extradural diamorphine, extradural phenoperidine and i.m. diamorphine following caesarean section. *Br J Anaesth* 1987, 59:354-359
248. Magann EF, Evans S, Hutchinson M, Collins R, Howard BC, Morrison JC: Postpartum hemorrhage after vaginal birth: An analysis of risk factors. *South Med J* 2005; 98:419-422
249. Mancuso A, De Vivo A, Giacobbe A, Priola V, Savasta M, Guzzo M, DeVivo D, Mancuso A: General versus spinal anaesthesia for elective caesarean sections: effects on neonatal short-term outcome. A prospective randomised study. *J Matern Fetal Neonatal Med* 2010; 23:1114-1118
250. Mandhyan R, Tiwari A, Cherian G: Congenital factor VII deficiency. *Br J Anaesth* 2010; 104:267-268
251. Marcoux S, Brisson J, Fabia J: The effect of cigarette smoking on the risk of preeclampsia and gestational hypertension. *Am J Epidemiol* 1989; 130:950-957
252. Margarson MP: Delayed amniotic fluid embolism following caesarean section under spinal anaesthesia. *Anaesthesia* 1995; 50:804-806
253. Martin JN, Jr., Harris BA, Jr., Huddleston JF, Morrison JC, Propst MG, Wiser WL, Perlis HW, Davidson JT: Vaginal delivery following previous cesarean birth. *Am J Obstet Gynecol* 1983; 146:255-263
254. Maruotti GM, Anfora R, Scanni E, Rispoli M, Mazzarelli LL, Napolitano R, Morlando M, Sarno L, Milanes GM, Simioli S, Migliucci A, Martinelli P, Mastronardi P: Anesthetic management of a parturient with spinal muscular atrophy type II. *J Clin Anesth* 2012; 24:573-577
255. Marx GF, Luykx WM, Cohen S: Fetal-neonatal status following caesarean section for fetal distress. *Br J Anaesth* 1984; 56:1009-1013

256. Matouskova A, Hanson B, Elm'en H: Continuous mini-infusion of bupivacaine into the epidural space during labor. Part III: A clinical study of 225 patients. *Acta Obstet Gynecol Scand Suppl* 1979; 83:43-52
257. Mayer DC, Quance D, Weeks SK: Headache after spinal anesthesia for cesarean section: a comparison of the 27-gauge Quincke and 24-gauge Sprotte needles. *Anesth Analg* 1992; 75:377-380
258. McAndrew CR, Harms P: Paraesthesiae during needle-through-needle combined spinal epidural versus single-shot spinal for elective caesarean section. *Anaesth Intensive Care* 2003; 31:514-517
259. McCaughey W, Howe JP, Moore J, Dundee JW: Cimetidine in elective Caesarean section. Effect on gastric acidity. *Anaesthesia* 1981; 36:167-172
260. McClune S, Regan M, Moore J: Laryngeal mask airway for caesarean section. *Anaesthesia* 1990; 45:227-228
261. McFarlane C: Failed intubation in an obese obstetric patient and the laryngeal mask. *Int J Obstet Anesth* 1993; 2:183-184
262. McGuinness GA, Merkow AJ, Kennedy RL, Erenberg A: Epidural anesthesia with bupivacaine for Cesarean section: neonatal blood levels and neurobehavioral responses. *Anesthesiology* 1978; 49:270-273
263. McNeilly G, Nicholl A, Broadway J, Rao S: Dural ectasia in Marfan's syndrome: magnetic resonance imaging appearances and anaesthetic experience of three deliveries. *Int J Obstet Anesth* 2013; 22:337-339
264. Meehan FP, Burke G, Kehoe JT: Update on delivery following prior cesarean section: a fifteen year review 1972-1987. *Int J Gynecol Obstet* 1989; 30:205-212
265. Mercier FJ, Riley ET, Frederickson WL, Roger-Christoph S, Benhamou D, Cohen SE: Phenylephrine added to prophylactic ephedrine infusion during spinal anesthesia for elective cesarean section. *Anesthesiology* 2001; 95:668-674
266. Mhyre JM, Bateman BT, Leffert LR: Influence of patient comorbidities on the risk of near-miss maternal morbidity or mortality. *Anesthesiology* 2011; 115:963-972
267. Milon D, Lavenac G, Noury D, Allain H, Van den Driessche J, Saint Marc C: Epidural anesthesia during labor: comparison of 3 combinations of fentanyl-bupivacaine and bupivacaine alone. *Ann Fr Anesth Reanim* 1986; 5:18-23
268. Milsom I, Forssman L, Biber B, Dottori O, Rydgren B, Sivertsson R: Maternal haemodynamic changes during caesarean section: a comparison of epidural and general anaesthesia. *Acta Anaesthesiol Scand* 1985; 29:161-167
269. Miro M, Guasch E, Gilsanz F: Comparison of epidural analgesia with combined spinal-epidural analgesia for labor: a retrospective study of 6497 cases. *Int J Obstet Anesth* 2008; 17:15-19
270. Missant C, Teunkenst A, Vandermeersch E, Van de Velde M: Patient-controlled epidural analgesia following combined spinal-epidural analgesia in labour: the effects of adding a continuous epidural infusion. *Anaesth Intensive Care* 2005; 33:452-456
271. Modanlou HD, Freeman RK, Ortiz O, Hinkes P, Pilsbury G Jr: Sinusoidal fetal heart rate pattern and severe fetal anemia. *Obstet Gynecol* 1977; 44:537-541
272. Módolo NS, de Azevedo VL, Santos PS, Rosa ML, Corvino DR, Alves LJ: Anesthetic strategy for cesarean section in a patient with factor XI deficiency. Case report. *Rev Bras Anestesiol* 2010; 60:176-180

273. Mojica JL, Melendez HJ, Bautista LE: The timing of intravenous crystalloid administration and incidence of cardiovascular side effects during spinal anesthesia: the results from a randomized controlled trial. *Anesth Analg* 2002; 94:432-437
274. Montan S, Ingemarsson I: Intrapartum fetal heart rate patterns in pregnancies complicated by hypertension. A cohort study. *Am J Obstet Gynecol* 1989; 160:283-288
275. Month RC, Vaida SJ: A combined spinal-epidural technique for labor analgesia and symptomatic relief in two parturients with idiopathic intracranial hypertension. *Int J Obstet Anesth* 2012; 21:192-194
276. Moodley J, Jjuuko G, Rout C: Epidural compared with general anaesthesia for caesarean delivery in conscious women with eclampsia. *Br J Obstet Gynecol* 2001; 108:378-382
277. Moore TR, Key TC, Reisner LS, Resnik R: Evaluation of the use of continuous lumbar epidural anesthesia for hypertensive pregnant women in labor. *Am J Obstet Gynecol* 1985; 152:404-412
278. Moran DH, Perillo M, LaPorta RF, Bader AM, Datta S: Phenylephrine in the prevention of hypotension following spinal anesthesia for cesarean delivery. *J Clin Anesth* 1991; 3:301-305
279. Morgan D, Philip J, Sharma S, Gottumukkala V, Perez B, Wiley J: A neonatal outcome with ephedrine infusions with or without preloading during spinal anesthesia for cesarean section. *Anesthesiology* 2000; A5
280. Morgan PJ, Halpern S, Lam-McCulloch J: Comparison of maternal satisfaction between epidural and spinal anesthesia for elective Cesarean section. *Can J Anaesth* 2000; 47:956-961
281. Moslemi F, Rasooli S: Comparison of spinal versus general anesthesia for cesarean delivery in patients with severe preeclampsia. *J Med Sci* 2007; 7:1044-1048
282. Mueller MD, Bruhwiler H, Schupfer GK, Luscher KP: Higher rate of fetal acidemia after regional anesthesia for elective cesarean delivery. *Obstet Gynecol* 1997; 90:131-134
283. Murphy DF, Nally B, Gardiner J, Unwin A: Effect of metoclopramide on gastric emptying before elective and emergency caesarean section. *Br J Anaesth* 1984; 56:1113-1116
284. Naef RW III, Chauhan SP, Chevalier SP, Roberts WE, Meydreich EF, Morrison JC: Prediction of hemorrhage at cesarean delivery. *Obstet Gynecol* 1994; 83:923-926
285. Nageotte MP, Larson D, Rumney PJ, Sidhu M, Hollenbach K: Epidural analgesia compared with combined spinal-epidural analgeisa during labor in nullaparous women. *New Eng J Med* 1997; 337:1715-1719
286. Nagy C, Wheeler A, Archer T: Acute normovolemic hemodilution, intraoperative cell salvage and pulse CO hemodynamic monitoring in a Jehovah's Witness with placenta percreta. *Int J Obstet Anesth* 2008; 17:159-163
287. Nasrat H, Fageeh W, Abalkhail B, Yamani T, Ardawi MSM: Determinants of pregnancy outcome in patients with gestational diabetes. *Int J Gynaecol Obstet* 1996; 53:117-123
288. Newton ER, Schroeder BC, Knape KG, Bennett BL: Epidural analgesia and uterine function. *Obstet Gynecol* 1995; 85:749-755
289. Ng PC, Wong MY, Nelson EA: Paediatrician attendance at caesarean section. *Eur J Pediatr* 1995; 154:672-675
290. Ngan Kee W, Khaw K, Lau T, Ng F, Chui K, Ng K: Randomised double-blinded comparison of phenylephrine vs ephedrine for maintaining blood pressure during spinal anesthesia for non-elective caesarean section. *Anaesthesia* 2008; 63:1319-1326

291. Ngan Kee WD, Khaw KS, Lee BB, Lau TK, Gin T: A dose-response study of prophylactic intravenous ephedrine for the prevention of hypotension during spinal anesthesia for cesarean delivery. *Anesth Analg* 2000; 90:1390-1395
292. Ngan Kee WD, Khaw KS, Lee BB, Ng FF, Wong MMS: Randomized controlled study of colloid preload before spinal anaesthesia for Caesarean section. *Br J Anaesth* 2001; 87:772-774
293. Ngan Kee WD, Khaw KS, Lee BB, Wong MM, Ng FF: Metaraminol infusion for maintenance of arterial blood pressure during spinal anesthesia for cesarean delivery: the effect of a crystalloid bolus. *Anesth Analg* 2001; 93:703-708
294. Ngan Kee WD, Khaw KS, Tan PE, Ng FF, Karmakar MK: Placental transfer and fetal metabolic effects of phenylephrine and ephedrine during spinal anesthesia for cesarean delivery. *Anesthesiology* 2009; 111: 506-512
295. Ngan Kee WD, Lee A, Khaw KS, Ng FF, Karmakar MK, Gin T: A randomized double-blinded comparison of phenylephrine and ephedrine combinations given by infusion to maintain blood pressure during spinal anesthesia for cesarean delivery: effects on fetal acid-base status and hemodynamic control. *Anesth Analg* 2008; 107:1295– 1302
296. Nickells JS, Vaughan DJ, Lillywhite NK, Loughnan B, Hasan M, Robinson PN: Speed of onset of regional analgesia in labour: a comparison of the epidural and spinal routes. *Anaesthesia* 2000; 55:17-20
297. Nielsen PE, Erickson JR, Abouleish EI, Perriatt S, Sheppard C: Fetal heart rate changes after intrathecal sufentanil or epidural bupivacaine for labor analgesia: incidence and clinical significance. *Anesth Analg* 1996; 83:742-746
298. Nishikawa K, Yokoyama N, Saito S, Goto F: Comparison of effects of rapid colloid loading before and after spinal anesthesia on maternal hemodynamics and neonatal outcomes in cesarean section. *J Clin Monit Comput* 2007; 21:125-129
299. Niv D, Rudick V, Golan A, Chayen MS: Augmentation of bupivacaine analgesia in labor by epidural morphine. *Obstet Gynecol* 1986; 67:206-209
300. Norris MC, Fogel ST, Conway-Long C: Combined spinal-epidural versus epidural labor analgesia. *Anesthesiology* 2001; 95:913-920
301. Norris MC, Grieco WM, Borkowski M, Leighton BL, Arkoosh VA, Huffnagle HJ, Huffnagle S: Complications of labor analgesia: epidural versus combined spinal epidural techniques. *Anesth Analg* 1994; 79:529-537
302. Numazaki M, Fujii Y: Reduction of emetic symptoms during cesarean delivery with antiemetics: propofol at subhypnotic dose versus traditional antiemetics. *J Clin Anesth* 2003; 15:423-427
303. Nuthalapaty FS, Rouse DJ, Owen J: The association of maternal weight with cesarean risk, labor duration, and cervical dilation rate during labor induction. *Obstet Gynecol* 2004; 103 452-456
304. Oh AY, Hwang JW, Song IA, Kim MH, Ryu JH, Park HP, Jeon YT, Do SH: Influence of the timing of administration of crystalloid on maternal hypotension during spinal anesthesia for cesarean delivery: preload versus coload. *BMC Anesthesiol* 2014; 14:36
305. Ohel G, Gonon R, Vaida S, Barak S, Gaitini L: Early versus late initiation of epidural analgesia in labor: Does it increase the risk of cesarean section? A randomized trial. *Am J Obstet Gynecol* 2006; 194:600-605
306. Ohel G, Harats H: Epidural anesthesia in early compared with advanced labor. *Int J Gynaecol Obstet* 1994; 45:217-219

307. Ojo OA, Akinkugbe OO: Nontoxemic hypertension in pregnancy in the African indigene. An analysis of 30 cases. *Am J Obstet Gynecol* 1969; 105:938-941
308. Olofsson C, Ekblom A, Skoldefors E, Waglund B, Irestedt L: Anesthetic quality during cesarean section following subarachnoid or epidural administration of bupivacaine with or without fentanyl. *Acta Anaesthesiol Scand* 1997; 41:332-338
309. Ong BY, Cohen MM, Palahniuk RJ: Anesthesia for cesarean section--effects on neonates. *Anesth Analg* 1989; 68:270-275
310. Ormezzano X, Francois TP, Viaud JY, Bukowski JG, Bourgeonneau MC, Cottron D, Ganansia MF, Gregoire FM, Grinand MR, Wessel PE: Aspiration pneumonitis prophylaxis in obstetric anaesthesia: comparison of effervescent cimetidine-sodium citrate mixture and sodium citrate. *Br J Anaesth* 1990; 64:503-506
311. Osinaike B, Ogah J: Anaesthesia for emergency Caesarean section in a patient with peripartum cardiomyopathy. *Cardiovasc J Afr* 2011; 22:337-340
312. O'Sullivan GM, Sear JW, Bullingham RES, Carrie LES: The effect of magnesium trisilicate mixture, metoclopramide and ranitidine on gastric pH, volume and serum gastrin. *Anaesthesia* 1985; 40:246-253
313. O'Sullivan GM, Smith M, Morgan B, Brighouse D, Reynolds F: H₂ antagonists and bupivacaine clearance. *Anaesthesia* 1988; 43:93-95
314. Ou CH, Tsou MY, Ting CK, Chiou CS, Chan KH, Tsai SK: Occurrence of the Bezold-Jarisch reflex during Cesarean section under spinal anesthesia--a case report. *Acta Anaesthesiol Taiwan* 2004; 42:175-178
315. Ozcakir HT, Lacin S, Baytur YB, Luleci N, Inceboz US: Different anesthesiologic strategies have no effect on neonatal jaundice. *Arch Gynecol Obstet* 2004; 270:179-181
316. Ozgen SZ, Ozgen S, Celiker R, Toraman F, Baykan N: Back pain after labour under epidural analgesia. *Middle East J Anesthesiol* 2014; 22:429-432
317. Paech MJ: Patient-controlled epidural analgesia in labour--is a continuous infusion of benefit? *Anaesth Intensive Care* 1992; 20:15-20
318. Palahniuk RJ, Scatliff J, Biehl D, Wiebe H, Sankaran K: Maternal and neonatal effects of methoxyflurane, nitrous oxide and lumbar epidural anaesthesia for Caesarean section. *Can Anaesth Soc J* 1977; 24:586-596
319. Palmer AW, Waugaman WR, Conklin KA, Kotelko DM: Does the administration of oral Bicitra before elective cesarean section affect the incidence of nausea and vomiting in the parturient? *Nurse Anesth* 1991; 2:126-133
320. Pan PH, Moore CH, Ross VH: Severe maternal bradycardia and asystole after combined spinal-epidural labor analgesia in a morbidly obese parturient. *J Clin Anesth* 2004; 16: 461-4
321. Pan PH, Moore CH: Comparing the efficacy of prophylactic metoclopramide, ondansetron, and placebo in cesarean section patients given epidural anesthesia. *J Clin Anesth* 2001; 13:430-435
322. Parameswara G, Kshama K, Murthy HK, Jalaja K, Venkat S: Early epidural labour analgesia: Does it increase the chances of operative delivery? *Br J Anaesth* 2012; 108 (Suppl 2):ii213-ii214
323. Parekh N, Husaini SW, Russell IF: Caesarean section for placenta praevia: a retrospective study of anaesthetic management. *Br J Anaesth* 2000; 84:725-730

324. Parker J, Balis N, Chester S, Adey D: Cardiopulmonary arrest in pregnancy: successful resuscitation of mother and infant following immediate caesarean section in labour ward. *Aust N Z J Obstet Gynaecol* 1996; 36:207-210
325. Parker RK, White PF: Epidural patient-controlled analgesia: an alternative to intravenous patient-controlled analgesia for pain relief after cesarean delivery. *Anesth Analg* 1992, 75:245-251
326. Pascual-Ramirez J, Haya J, Perez-Lopez F, Gil-Trujillo S, Garrido-Esteban R, Berna G: Effect of combined spinal-epidural analgesia versus epidural analgesia on labor and delivery duration. *Int J Gynaecol Obstet* 2011; 114:246-250
327. Patel N, Armstrong S, Fernando R, Columb M, Bray J, Sodhi V, Lyons G: Combined spinal epidural vs epidural labour analgesia: does initial intrathecal analgesia reduce the subsequent minimum local analgesic concentration of epidural bupivacaine? *Anaesthesia* 2012; 67:584-593.
328. Patel NP, El-Wahab N, Fernando R, Wilson S, Robson SC, Columb MO, Lyons GR: Fetal effects of combined spinal-epidural vs epidural labour analgesia: a prospective, randomised double-blind study. *Anaesthesia* 2014; 69:458-467
329. Paternoster D, Pascoli I, Parotta M, Cester M, Nanhorngue K, Merati E, Arrigoni S, Garofalo G, Bertolino M, Micaglio M, Ripa C: Pulmonary hypertension during pregnancy: management of two cases. *Arch Gynecol Obstet* 2010; 281:431-434
330. Pearson JF, Davies P: The effect of continuous lumbar epidural analgesia on the acid-base status of maternal arterial blood during the first stage of labour. *J Obstet Gynaecol Br Commonw* 1973; 80:218-224
331. Pearson JF, Davies P: The effect of continuous lumbar epidural analgesia upon fetal acid-base status during the second stage of labour. *J Obstet Gynaecol Br Commonw* 1974; 81:975-979
332. Perriss BW, Latham BV, Wilson IH: Analgesia following extradural and i.m. pethidine in post-caesarean section patients. *Br J Anaesth* 1990, 64:355-357
333. Petropoulos G, Siristatidis C, Salamalekis E, Creatsas G: Spinal and epidural versus general anesthesia for elective cesarean section at term: effect on the acid-base status of the mother and newborn. *J Matern Fetal Neonatal Med* 2003;13:260-266
334. Petry J, Vercauteren M, Van Mol I, Van Houwe P, Adriaensen HA: Epidural PCA with bupivacaine 0.125%, sufentanil 0.75 microgram and epinephrine 1/800.000 for labor analgesia: is a background infusion beneficial? *Acta Anaesthesiol Belg* 2000; 51:163-166
335. Phillips GH: Epidural sufentanil/bupivacaine combinations for analgesia during labor: effect of varying sufentanil doses. *Anesthesiology* 1987; 67:835-838
336. Pierce ET, Carr DB, Datta S: Effects of ephedrine and phenylephrine on maternal and fetal atrial natriuretic peptide levels during elective cesarean section. *Acta Anaesth Scand* 1994; 38:48-51
337. Polley LS, Columb MO, Lyons G, Nair SA: The effect of epidural fentanyl on the minimum local analgesic concentration of epidural chlorprocaine in labor. *Anesth Analg* 1996; 83:987-990
338. Popat MT, Chippa JH, Russell R: Awake fibreoptic intubation following failed regional anaesthesia for caesarean section in a parturient with Still's disease. *Eur J Anaesthesiol* 2000; 17:211-214

339. Porreco RP, Boehm FH, Dildy GA, Miller HS, Wickstrom EA, Garite TJ, Swedlow D: Dystocia in nulliparous patients monitored with fetal pulse oximetry. *Am J Obstet Gynecol* 2004; 190: 113-7
340. Porter JS, Bonello E, Reynolds F: The effect of epidural opioids on maternal oxygenation during labour and delivery. *Anaesthesia* 1996; 51:899-903
341. Potter PS, Waters JH, Burger GA, Mraovic B: Application of cell-salvage during Cesarean section. *Anesthesiology* 1999; 90:619-621
342. Prakash S, Pramanik V, Chellani H, Salhan S, Gogia A: Maternal and neonatal effects of bolus administration of ephedrine and phenylephrine during spinal anaesthesia for caesarean delivery: a randomised study. *Int J Obstet Anesth* 2010; 19:24-30
343. Price C, Lafreniere L, Brosnan C, Findlay I: Regional analgesia in early active labour: combined spinal-epidural vs epidural. *Anaesthesia* 1998; 53:951-955
344. Priscu V, Priscu L, Soroker D: Laryngeal mask for failed intubation in emergency Caesarean section. *Can J Anaesth* 1992; 39:893
345. Pruefer C, Bewlay A: Respiratory arrest with remifentanil patient-controlled analgesia--another case. *Anaesthesia* 2012; 67:1044-1045
346. Purdie J, Reid J, Thorburn J, Asbury AJ: Continuous extradural analgesia: comparison of midwife top-ups, continuous infusions and patient controlled administration. *Br J Anaesth* 1992; 68:580-584, 1992
347. Qublan HS, Merhej A, Dabbas MA, Hindawi IM: Spinal versus general anesthesia for elective cesarean delivery: a prospective comparative study. *Clin Exp Obstet Gynecol* 2001; 28:246-248
348. Qvist N, Storm K, Holmskov A: Cimetidine as pre-anesthetic agent for cesarean section: perinatal effects on the infant, the placental transfer of cimetidine and its elimination in the infants. *J Perinat Med* 1985; 13:179-183
349. Qvist N, Storm K: Cimetidine pre-anesthetic. A prophylactic method against Mendelson's syndrome in cesarean section. *Acta Obstet Gynecol Scand* 1983; 62:157-159
350. Rafferty TD, Berkowitz RL: Hemodynamics in patients with severe toxemia during labor and delivery. *Am J Obstet Gynecol* 1980; 138:263-270
351. Ramanathan J, Sibai BM, Vu T, Chauhan D: Correlation between bleeding times and platelet counts in women with preeclampsia undergoing cesarean section. *Anesthesiology* 1989; 71:188-191
352. Ramin SM, Gambling DR, Lucas MJ, Sharma SK, Sidawi JE, Leveno KJ: Randomized trial of epidural versus intravenous analgesia during labor. *Obstet Gynecol* 1995; 86:783-789
353. Ramin SM, Ramin KD, Cox K, Magness RR, Shearer VE, Gant NF: Comparison of prophylactic angiotensin II versus ephedrine infusion for prevention of maternal hypotension during spinal anesthesia. *Am J Obstet Gynecol* 1994; 171:734-739
354. Ransom SB, Fundaro G, Dombrowski MP: Cost-effectiveness of routine blood type and screen testing for cesarean section. *J Reprod Med* 1999; 44:592-594
355. Ransom SB, Fundaro G, Dombrowski MP: The cost effectiveness of routine type and screen admission testing from expected vaginal delivery. *Obstet Gynecol* 1998; 92:493-495
356. Ranta P, Jouppila P, Spalding M, Jouppila R: The effect of maternal obesity on labour and labour pain. *Anaesthesia* 1995; 50:322-326

357. Ratcliffe FM, Evans JM: Neonatal wellbeing after elective caesarean delivery with general, spinal, and epidural anaesthesia. *Eur J Anaesthesiol* 1993; 10:175-181
358. Rawal N, Schollin J, Wesstrom G: Epidural versus combined spinal epidural block for cesarean section. *Acta Anaesthesiol Scan* 1988; 32:61-66
359. Redman CWG, Bonnar J, Beilin L. Early platelet consumption in pre-eclampsia. *Br Med J* 1978; 1:467-469
360. Reynolds F, O'Sullivan G: Epidural fentanyl and perineal pain in labour. *Anaesthesia* 1989; 44:341-344
361. Reynolds F, Russell R, Porter J, Smeeton M: Does the use of low dose bupivacaine/opioid epidural infusion increase the normal delivery rate? *Int J Obstet Anesth* 2003; 12:156-163
362. Riley ET, Cohen SE, Macario A, Desai JB, Ratner EF: Spinal versus epidural anesthesia for cesarean section: a comparison of time efficiency, costs, charges, and complications. *Anesth Analg* 1995; 80:709-712
363. Riley ET, Flanagan B, Cohen SE, Chitkara U. Intravenous nitroglycerin: a potent uterine relaxant for emergency obstetric procedures. Review of literature and report of three cases. *Int J Obstet Anesth* 1996; 5:264-268
364. Roberts SW, Leveno KJ, Sidawi JE, Lucas MJ, Kelly MA: Fetal acidemia associated with regional anesthesia for elective cesarean delivery. *Obstet Gynecol* 1995; 85:79-83
365. Roberts WE, Perry KG, Jr., Woods JB, Files JC, Blake PG, Martin JN, Jr.: The intrapartum platelet count in patients with HELLP (hemolysis, elevated liver enzymes, and low platelets) syndrome: is it predictive of later hemorrhagic complications? *Am J Obstet Gynecol* 1994; 171:799-804
366. Robinson HE, O'Connell CM, Joseph KS, McLeod NL: Maternal outcomes in pregnancies complicated by obesity. *Obstet Gynecol* 2005; 106:1357-1364
367. Robson SC, Boys RJ, Rodeck C, Morgan B: Maternal and fetal haemodynamic effects of spinal and extradural anaesthesia for elective caesarean section. *Br J Anaesth* 1992; 68:54-59
368. Rodriguez J, Abboud TK, Reyes A, Payne M, Zhu J, Steffens Z, Afrasiabi A: Continuous infusion epidural anesthesia during labor: a randomized, double-blind comparison of 0.0625% bupivacaine/0.002% butorphanol and 0.125% bupivacaine. *Reg Anesth* 1990; 15:300-303
369. Rogers R, Gilson G, Hammerer-Doak D: Epidural analgesia and active management of labor and mode of delivery. *Obstet Gynecol* 1999; 93:995-998
370. Rogers W, Wernimont S, Kumar G, Bennett E, Chestnut D: Acute hypotension associated with intraoperative cell salvage using a leukocyte depletion filter during management of obstetric hemorrhage due to amniotic fluid embolism. *Anesth Anag* 2013; 117:449-52
371. Rolbin SH, Cohen MM, Levinton CM, Kelly EN, Farine D: The premature infant: anesthesia for cesarean delivery. *Anesth Analg* 1994; 78:912-917
372. Romero R, Mazor M, Lockwood CJ, Emamian M, Belanger KP, Hobbins JC, Duffy T: Clinical significance, prevalence, and natural history of thrombocytopenia in pregnancy-induced hypertension. *Am J Perinatol* 1989; 6:32-38
373. Rosen MA, Hughes SC, Shnider SM, Abboud TK, Norton M, Dailey PA, Curtis JD: Epidural morphine for the relief of postoperative pain after cesarean delivery. *Anesth Analg* 1983; 62:666-672

374. Ross AW, Greenhalgh C, McGlade DP, Balson IG, Chester SC, Hutchinson RC, Ashley JE: The Sprotte needle and post dural puncture headache following caesarean section. *Anaesth Intensive Care* 1993; 21:280-283
375. Ross BK, Chadwick HS, Mancuso JJ, Benedetti C: Sprotte needle for obstetric anesthesia: decreased incidence of post dural puncture headache. *Reg Anesth* 1992; 17:29-33
376. Rout CC, Rocke DA, Levin J, Gouws E, Reddy D: A reevaluation of the role of crystalloid preload in the prevention of hypotension associated with spinal anesthesia for elective cesarean section. *Anesthesiology* 1993 79:262-269
377. Roux M, Wattrisse G, Tai RB, Dufossez F, Krivacic-Horber R: Obstetric analgesia: peridural analgesia versus combined spinal and peridural analgesia. *Ann Fr Anesth Reanim* 1999; 18:487-498
378. Rudick V, Niv D, Hetman Peri M, Geller E, Avni A, Golan A: Epidural analgesia for planned vaginal delivery following previous cesarean section. *Obstet Gynecol* 1984; 64:621-623
379. Russell R, Reynolds F: Epidural infusion of low-dose bupivacaine and opioid in labour. Does reducing motor block increase the spontaneous delivery rate? *Anaesthesia* 1996; 51:266-273
380. Ryhanen P, Jouppila R, Lanning M, Jouppila P, Hollm'en A, Kouvalainen K: Natural killer cell activity after elective cesarean section under general and epidural anesthesia in healthy parturients and their newborns. *Gynecol Obstet Invest* 1985; 19:139-142
381. Saito M, Okutomi T, Kanai Y, Mochizuki J, Tani A, Amano K, Hoka S: Patient-controlled epidural analgesia during labor using ropivacaine and fentanyl provides better maternal satisfaction with less local anesthetic requirement. *J Anesth* 2005; 19:208-212
382. Sakala EP, Kaye S, Murray RD, Munson LJ: Epidural analgesia: effect on the likelihood of a successful trial of labor after cesarean section. *J Reprod Med* 1990; 35:886-890
383. Salim R, Nachum Z, Moscovici R, Lavee M, Shalev E: Continuous compared with intermittent epidural infusion on progress of labor and patient satisfaction. *Obstet Gynecol* 2005;106:301-306
384. Salman C, Kayacan N, Ertugrul F, Bigat Z, Karsh B: Combined spinal-epidural anesthesia with epidural volume extension causes a higher level of block than single-shot spinal anesthesia. *Rev Bras Anestesiol* 2013; 63:267-272
385. Saravanan S, Kocarev M, Wilson RC, Watkins E, Columb MO, Lyons G: Equivalent dose of ephedrine and phenylephrine in the prevention of post-spinal hypotension in Caesarean section. *Br J Anaesth* 2006; 96:95-99
386. Sarvela J, Halonen P, Soikkeli A, Korttila K: A double-blinded, randomized comparison of intrathecal and epidural morphine for elective cesarean delivery. *Anesth Analg* 2002; 95:436-440
387. Schewe J, Komusin A, Zinserling J, Nadstawek J, Hoeft A, Hering R: Effects of spinal anaesthesia versus epidural anaesthesia for caesarean section on postoperative analgesic consumption and postoperative pain. *Eur J Anaestesiol* 2009; 26:52-59
388. Sendag F, Terek C, Oztekin K, Sagol S, Asena U: Comparison of epidural and general anaesthesia for elective caesarean delivery according to the effects of apgar scores and acid-base status. *Aust N Z J Obstet Gynaecol* 1999; 39:464-468
389. Sener EB, Guldogus F, Karakaya D, Baris S, Kocamanoglu S, Tur A: Comparison of neonatal effects of epidural and general anesthesia for cesarean section. *Gynecol Obstet Invest* 2003; 55:41-45

390. Sezer O, Gunaydin B: Efficacy of patient-controlled epidural analgesia after initiation with epidural or combined spinal-epidural analgesia. *Int J Obstet Anesth* 2007; 16:226-230
391. Shaban M, Ali N, Abd El-Razek A: Spinal versus general anesthesia in preeclamptic patients undergoing cesarean delivery. *El-Minia Med Bull* 2005; 16:328-343
392. Shakil Y: Efficacy of crystalloid preload and co-load in prevention of hypotension in spinal anesthesia for elective caesarean section [abstract]. *Anaesth, Pain & Intensive Care* 2011; 15:151
393. Sharma R, Chatterjee J, Edmonds K: Horner's syndrome with epidural anaesthesia. *BMJ Case Rep* 2010 Aug 23;2010. pii: bcr0120102698. doi:
394. Sharma SK, Philip J: The effect of anesthetic techniques on blood coagulability in parturients as measured by thromboelastography. *Anesth Analg* 1997; 85:82-86
395. Shatken S, Greenough K, McPherson C: Epidural fever and its implications for mothers and neonates: taking the heat. *J Midwifery Womens Health* 2012; 57:82-85
396. Shehabi Y, Gatt S, Buckman T, Isert P: Effect of adrenaline, fentanyl and warming of injectate on shivering following extradural analgesia in labour. *Anaesth Intensive Care* 1990; 18:31-37
397. Sheiner E, Mazor M, Levy A, Wiznitzer A, Bashiri A: Pregnancy outcome of asthmatic patients: a population-based study. *J Matern Fetal Neonatal Med* 2005; 18:237-240
398. Shutt LE, Valentine SJ, Wee MYK, Page RJ, Prossner A, Thomas TA: Spinal anaesthesia for caesarean section: comparison of 22 gauge and 25 gauge Whitacre needles with 26 gauge Quincke needles. *Br J Anaesth* 1992; 69:589-594
399. Shyken JM, Smeltzer JS, Baxi LV, Blakemore KJ, Ambrose SE, Petrie RH: A comparison of the effect of epidural, general, and no anesthesia on funic acid-base values by stage of labor and type of delivery. *Am J Obstet Gynecol* 1990; 163:802-807
400. Sia A, Leo S, Ocampo C: A randomised comparison of variable-frequency automated mandatory boluses with a basal infusion for patient-controlled epidural analgesia during labour and delivery. *Anaesthesia* 2013; 68:267-275
401. Sia AT, Chong JL: Epidural 0.2% ropivacaine for labour analgesia: parturient-controlled or continuous infusion? *Anaesth Intensive Care* 1999; 27:154-158
402. Siddik-Sayyid S, Taha S, Kanazi E, Aouad: A randomized controlled trial of variable rate phenylephrine infusion with rescue phenylephrine boluses versus rescue boluses alone on physician interventions during spinal anesthesia for elective cesarean delivery. *Anesth Analg* 2014; 118:611-8
403. Siddik-Sayyid SM, Gellad PH, Aouad MT: Total spinal block after spinal anesthesia following ongoing epidural analgesia for cesarean delivery. *J Anesth* 2012; 26:312-313
404. Siddik-Sayyid SM¹, Nasr VG, Taha SK, Zbeide RA, Shehade JM, Al Alami AA, Mokadem FH, Abdallah FW, Baraka AS, Aouad MT: A randomized trial comparing colloid preload to colloid during spinal anesthesia for elective cesarean delivery. *Anesth Analg* 2009; 109:1219-1224
405. Simon L, Provenchere S, de Saint Blanquat L, Boulay G, Hamza J: Dose of prophylactic intravenous ephedrine during spinal anesthesia for cesarean section. *J Clin Anesth* 2001; 13:366-369
406. Simon L, Santi TM, Sacquin P, Hamza J: Pre-anaesthetic assessment of coagulation abnormalities in obstetric patients: usefulness, timing and clinical implications. *Br J Anaesth* 1997; 78:678-683

407. Sinatra RS, Goldstein R, Sevarino FB: The clinical effectiveness of epidural bupivacaine, bupivacaine with lidocaine, and bupivacaine with fentanyl for labor analgesia. *J Clin Anesth* 1991; 3:219-224
408. Skret-Magierło J, Barnaś E, Sek-Kłebukowska B, Nicpoń J, Kloc G: Intracranial hematoma as the cause of headache after subarachnoid anesthesia for cesarean section--a case report. *Ginekol Pol* 2014; 85:58-61
409. Skupski D, Abramovitz S, Samuels J, Pressimone V, Kjaer K: Adverse effects of combined spinal-epidural versus traditional epidural analgesia during labor. *Int J Obstet Anesth* 2009; 106:242-245
410. Smedvig JP, Soreide E, Gjessing L: Ropivacaine 1 mg/ml, plus fentanyl 2 microg/ml for epidural analgesia during labour. Is mode of administration important? *Acta Anaesth Scand* 2001; 45:595-599
411. Smith ID, Klubien KE, Wood MLB, Macrae DJ, Carli F: Diamorphine analgesia after caesarean section: comparison of intramuscular and epidural administration of four dose regimens. *Anaesthesia* 1991, 46:973-976
412. Spencer JA, Koutsoukis M, Lee A: Fetal heart rate and neonatal condition related to epidural analgesia in women reaching the second stage of labour. *Eur J Obstet Gynecol Reprod Biol* 1991; 41:173-178
413. Spinapolic RX: The effects of continuous epidural anesthesia on fetal heart rate during labor. *J Am Osteopath Assoc* 1977; 76:695-698
414. Stavrou C, Hofmeyr GJ, Boenzaart AP: Prolonged fetal bradycardia during epidural analgesia. Incidence, timing and significance. *S Afr Med J* 1990; 77:66-68
415. Stein DJ, Birnbach DJ, Danzer BI, Kuroda MM, Grunbaum A, Thys DM: Acupressure versus intravenous metoclopramide to prevent nausea and vomiting during spinal anesthesia for cesarean section. *Anesth Analg* 1997; 84:342-345
416. Storey J: The laryngeal mask for failed intubation at caesarean section. *Anaesth Intensive Care* 1992; 20:118-119
417. Stovall TG, Shaver DC, Solomon SK, Anderson GD: Trial of labor in previous cesarean section patients, excluding classical cesarean sections. *Obstet Gynecol* 1987; 70:713-717
418. Strickland DM, Guzick DS, Cox K, Gant NF, Rosenfeld CR: The relationship between abortion in the first pregnancy and development of pregnancy-induced hypertension in the subsequent pregnancy. *Am J Obstet Gynecol* 1986; 154:146-148
419. Suelto MD, Vincent RD Jr, Larmon JE, Norman PF, Werhan CF: Spinal anesthesia for postpartum tubal ligation after pregnancy complicated by preeclampsia or gestational hypertension. *Reg Anesth Pain Med* 2000; 25:170-173
420. Sutcliffe N, Remington SA, Ramsay TM, Mason C: Severe tracheal stenosis and operative delivery. *Anaesthesia* 1995; 50:26-29
421. Swayze CR, Skerman JH, Walker EB, Sholte FG: Efficacy of subarachnoid meperidine for labor analgesia. *Reg Anesth* 1991; 16:309-313
422. Tan S, Reid J, Thorburn J: Extradural analgesia in labour: complications of three techniques of administration. *Br J Anaesth* 1994; 73:619-623
423. Tawfik MM¹, Hayes SM², Jacoub FY², Badran BA², Gohar FM², Shabana AM³, Abdelkhalek M², Emara MM: Comparison between colloid preload and crystalloid co-load in cesarean section under spinal anesthesia: a randomized controlled trial. *Int J Obstet Anesth.* 2014; 23:317-323

424. Teoh WH, Sia AT: Colloid preload versus colloid for spinal anaesthesia for cesarean delivery: the effects on maternal cardiac output. *Anesth Analg* 2009; 108:1592-1598
425. Thomas DG, Robson SC, Redfern N, Hughes D, Boys RJ: Randomized trial of bolus phenylephrine or ephedrine for maintenance of arterial pressure during spinal anaesthesia for Caesarean section. *Br J Anaesth* 1996; 76:61-65
426. Thoren T, Holmstrom B, Rawal N, Schollin J, Lindeberg S, Skeppner G: Sequential combined spinal epidural block versus spinal block for cesarean section: effects on maternal hypotension and neurobehavioral function of the newborn. *Anesth Analg* 1994; 78:1087-1092
427. Thorp JA, Hu DH, Albin RM, McNitt J, Meyer BA, Cohen GR, Yeast JD: The effect of intrapartum epidural analgesia on nulliparous labor: a randomized, controlled, prospective trial. *Am J Obstet Gynecol* 1993; 169:851-858
428. Toldy M, Scott DB: Blood loss during Caesarean section under general anaesthesia. *Br J Anaesth* 1969; 41:868-873
429. Trudinger BJ: Platelets and intrauterine growth retardation in pre-eclampsia. *Br J Obstet Gynecol* 1976; 83:284-286
430. Tsai PS, Hsu CS, Fan YC, Huang CJ: General anaesthesia is associated with increased risk of surgical site infection after Caesarean delivery compared with neuraxial anaesthesia: a population-based study. *Br J Anaesth* 2011; 107:757-761
431. Tsen LC, Cherayil G: Sickle cell-induced peripheral neuropathy following spinal anesthesia for cesarean delivery. *Anesthesiology* 2001; 95:1298-1299
432. Tsen LC, Thue B, Datta S, Segal S: Is combined spinal-epidural analgesia associated with more rapid cervical dilation in nulliparous patients when compared with conventional epidural analgesia? *Anesthesiology* 1999; 91:920-925
433. Tunstall ME, Geddes C: "Failed intubation" in obstetric anesthesia. *Br J Anaesth* 1984; 56:659-661
434. Turkoz A, Togal T, Gokdeniz R, Toprak HI, Ersoy O: Effectiveness of intravenous ephedrine infusion during spinal anaesthesia for caesarean section based on maternal hypotension, neonatal acid-base status and lactate levels. *Anaesth Intensive Care* 2002; 30:316-320
435. Tympa A, Grigoriadis C, Petropoulos G, Hassiakos D: Horner syndrome following bolus low dose epidural analgesia for labor. *J Anesth* 2013; 27:799-800
436. Ulbrich F, Engelstädtter H, Wittau N, Steinmann D: Anaesthetic management of emergency caesarean section in a parturient with systemic mastocytosis. *Int J Obstet Anesth* 2013; 22:243-246
437. Usha Kiran TS, Hemmadi S, Bethel J, Evans J: Outcome of pregnancy in a woman with an increased body mass index. *BJOG* 2005; 112:768-772
438. Vahrtian A, Zhang J, Hasling J, Troendle JF, Klebanoff MA, Thorp JM, Jr: The effect of early epidural versus early intravenous analgesia use on labor progression: a natural experiment. *Am J Obstet Gynecol* 2004; 191:259-265
439. Vallejo M, Ramesh V, Phelps A, Sah N: Patient-controlled epidural analgesia with background infusion versus without a background infusion. *J Pain* 2007; 8:970-975
440. Vallejo MC, Mandell GL, Sabo DP, Ramanathan S: Postdural puncture headache: a randomized comparison of five spinal needles in obstetric patients. *Anesth Analg* 2000; 91:916-920

441. Van Bogaert LJ: Spinal block caesarean section in parturients with pregnancy-induced hypertension. *East Afr Med J* 1998; 75:227-231
442. Van de Velde M, Berends N, Spitz B, Teunkens A, Vandermeersch E: Low-dose combined spinal-epidural anaesthesia vs. conventional epidural anaesthesia for Caesarean section in pre-eclampsia: a retrospective analysis. *Eur J Anaesthesiol* 2004; 21:454-459
443. Van de Velde M, Mignolet K, Vandermeersch E, Van Assche A: Prospective, randomized comparison of epidural and combined spinal epidural analgesia during labor. *Acta Anaesthesiol Belg* 1999; 50:129-136
444. Van de Velde M, Vercauteren M, Vandermeersch E: Fetal heart rate abnormalities after regional analgesia for labor pain: the effect of intrathecal opioids. *Reg Anesth Pain Med* 2001; 26:257-262
445. Van den Berg A, Ghatge S, Armendariz G, Cornelius D, Wang S: Responses to dural puncture during institution of combined spinal-epidural analgesia: a comparison of 27 gauge pencil-point and 27 gauge cutting-edge needles. *Anaesth Intensive Care* 2011; 39:247-251
446. Van Steenberge A, Debroux HC, Noorduin H: Extradural bupivacaine with sufentanil for vaginal delivery. A double-blind trial. *Br J Anaesth* 1987; 59:1518-1522
447. Varshney R, Jain G: Comparison of colloid preload versus colloid under low dose spinal anesthesia for cesarean delivery. *Anesth Essays Res* 2013; 7:376-380
448. Vercauteren MP, Coppejans HC, Hoffmann VH, Mertens E, Adriaensen HA: Prevention of hypotension by a single 5-mg dose of ephedrine during small-dose spinal anesthesia in prehydrated cesarean delivery patients. *Anesth Analg* 2000; 90:324-327
449. Vernis L, Duale C, Storme B, Mission JP, Rol B, Schoeffler P: Perispinal analgesia for labour followed by patient-controlled infusion with bupivacaine and sufentanil: combined spinal-epidural vs. epidural analgesia alone. *Eur J Anaesthesiol* 2004; 21:186-192
450. Vertommen JD, Vandermeulen E, Van Aken H, Vaes L, Soetens M, Van Steenberge A, Mourisse P, Willaert J, Noorduin H, Devlieger H, Van Assche AF: The effects of the addition of sufentanil to 0.125% bupivacaine on the quality of analgesia during labor and on the incidence of instrumental deliveries. *Anesthesiology* 1991; 74:809-814
451. Vickers RJ, May AE: Long-term backache after extradural or general anaesthesia for manual removal of placenta: preliminary report. *Br J Anaesth* 1993; 70:214-215
452. Vigil-De Gracia P, Silva S, Montufar C, Carroll I, De Los Rios S: Anesthesia in pregnant women with HELLP syndrome. *Int J Gynaecol Obstet* 2001; 74:23-27
453. Vincent RD Jr, Martin RW: Postpartum tubal ligation after pregnancy complicated by preeclampsia or gestational hypertension. *Obstet Gynecol* 1996; 88:119-122
454. Visalyaputra S, Prechapanich J, Suwanvichai S, Yimyam S, Permpolprasert L, Suksoppee P: Intravenous nitroglycerin for controlled cord traction in the management of retained placenta. *Int J Gynaecol Obstet* 2011; 112:103-106
455. Visalyaputra S, Rodanant O, Somboonviboon W, Tantivitayatan K, Thienthong S, Saengchote W: Spinal versus epidural anesthesia for cesarean delivery in severe preeclampsia: a prospective randomized, multicenter study. *Anesth Analg* 2005; 101:862-868
456. Viscomi C, Eisenach JC: Patient-controlled epidural analgesia during labor. *Obstet Gynecol* 1991; 77:348-351
457. Viscomi CM, Hood DD, Melone PJ, Eisenach JC: Fetal heart rate variability after epidural fentanyl during labor. *Anesth Analg* 1990; 71:679-683

458. Visser W, Wallenburg HC: Temporising management of severe pre-eclampsia with and without the HELLP syndrome. *Br J Obstet Gynaecol* 1995; 102:111-117
459. von Ungern-Sternberg BS, Regli A, Bucher E, Reber A, Schneider MC: Impact of spinal anaesthesia and obesity on maternal respiratory function during elective Caesarean section. *Anaesthesia* 2004; 59:743-749
460. Wallace DH, Leveno KJ, Cunningham FG, Giesecke AH, Shearer VE, Sidawi JE: Randomized comparison of general and regional anesthesia for cesarean delivery in pregnancies complicated by severe preeclampsia. *Obstet Gynecol* 1995; 86:193-199
461. Wang F, Shen X, Guo X, Peng Y, Gu X (The Labor Analgesia Examining Group): Epidural analgesia in the latent phase of labor and the risk of cesarean delivery. *Anesthesiology* 2009; 111:871-80
462. Wang LZ, Chang XY, Hu XX, Tang BL, Xia F: The effect on maternal temperature of delaying initiation of the epidural component of combined spinal-epidural analgesia for labor: A pilot study. *Ing J Obstet Anesth* 2011; 20:312-317
463. Waterstone M, Bewley S, Wolfe C: Incidence of severe obstetric morbidity: case-control study. *BMJ* 2001; 322:1089-1093
464. Webb AA, Shipton EA: Re-evaluation of i.m. ephedrine as prophylaxis against hypotension associated with spinal anaesthesia for Caesarean section. *Can J Anaesth* 1998; 45:367-369
465. Weiner MM, Vahl TP, Kahn RA: Case scenario: Cesarean section complicated by rheumatic mitral stenosis. *Anesthesiology* 2011; 114:949-957
466. Weissman A, Torkhov O, Weissman A, Drugan A: The effects of meperidine and epidural analgesia in labor on maternal heart rate variability. *Int J Obstet Anesth* 2009; 18:118-124
467. Wig J, Biswas GC, Malhotra SK, Gupta AN: Comparison of sodium citrate with magnesium trisilicate as pre-anaesthetic antacid in emergency caesarean sections. *Indian J Med Res* 1987; 85:306-310
468. Withington DE, Weeks SK: Repeat epidural analgesia and unilateral block. *Can J Anaesth* 1994; 41:568-571
469. Wong CA, McCarthy RJ, Sullivan JT, Scavone BM, Gerber SE, Yaghmour EA: Early compared with late neuraxial analgesia in nulliparous labor induction: a randomized controlled trial. *Obstet Gynecol* 2009; 113:1066-1074
470. Wong CA, Scavone BM, Peaceman AM, McCarthy RJ, Sullivan JT, Diaz NT, Yaghmour E, Marcus RJ, Sherwani SS, Sproviero MT, Yilmaz M, Patel R, Robles C, Grouper S: The risk of cesarean delivery with neuraxial analgesia given early versus late in labor. *N Engl J Med* 2005; 352:655-665
471. Wright PMC, Allen RW, Moore J, Donnelly JP: Gastric emptying during lumbar extradural analgesia in labour: effect of fentanyl supplementation. *Br J Anaesth* 1992; 68:248-251
472. Yang YW, Chen WT, Chen JY, Lee SC, Chang Y, Wen YR: Bacterial infection in deep paraspinal muscles in a parturient following epidural analgesia. *Acta Anaesthesiol Taiwan* 2011; 49:75-78
473. Yau G, Gregory MA, Gin T, Bogod DG, Oh TE: The addition of fentanyl to epidural bupivacaine in first stage labour. *Anaesth Intensive Care* 1990; 18:532-535
474. Yau G, Gregory MA, Gin T, Oh TE: Obstetric epidural analgesia with mixtures of bupivacaine, adrenaline and fentanyl. *Anaesthesia* 1990; 45:1020-1023

475. Yentur E, Topcu, Ekici Z, Ozturk T, Keles G, Civi M: The effect of epidural and general anesthesia on newborn rectal temperature at elective cesarean section. *Braz J Med Biol Res* 2009; 42:863-867
476. Zagorzycki MT, Brinkman CR, III: The effect of general and epidural anesthesia upon neonatal Apgar scores in repeat cesarean section. *Surg Gynecol Obstet* 1982; 155:641-645
477. Zilanti M, Salazar JR, Aller J, Aguero O: Fetal heart rate and pH of fetal capillary blood during epidural analgesia in labor. *Obstet Gynecol* 1970; 36(6):881-888