

Appendix 1. Postpartum Hemorrhage After Nulliparous, Term, Singleton Vertex Vaginal Deliveries in Women With Normal Platelet Count or Mild Thrombocytopenia but Without Gestational Hypertension or Preeclampsia

Outcome measures		Platelet count ≥150 k/μl N=1879	Platelet count 100-149 k/μl N=192	P
Postpartum hemorrhage (primary outcome)		141 (7.5)	27 (14.1)	0.002
Methergine administered after delivery		116 (6.2)	21 (10.9)	0.01
Hemabate administered after delivery		17 (0.9)	11 (5.7)	<0.001
Methergine and hemabate administered after delivery		16 (0.9)	8 (4.2)	<0.001
Estimated blood loss ≥ 1000 ml		7 (0.4)	1 (0.5)	0.54
Blood transfusion		19 (1.0)	0 (0)	0.25

Data are n (percent) with P-value by χ^2 test or Fisher's exact test (cell count ≤5).

Govindappagari S, Moyle K, Burwick RM. Mild thrombocytopenia and postpartum hemorrhage in nulliparous, term, singleton vertex deliveries. *Obstet Gynecol* 2020;135.

The authors provided this information as a supplement to their article.

©2020 American College of Obstetricians and Gynecologists.