
## **Appendix 1. Primary Outcome and Prespecified Secondary Outcomes**

Outcome (Per Protocol)	Outpatient DS (n=164)	Inpatient DS (n=171)	RR or MD (95% CI)
Primary outcome Hospital stay > 48 hours	87 (53.0)	152 (88.9)	0.6 (0.51-0.7)
Secondary outcomes Total hospital length of stay (hours:mins)	54:03	62:19	-08:15 (-11:4704:43)
Data are presented as n (%), median [Interquartile range], or mean + SD unless otherwise indicated.			


Saad AF, Gavara R, Beavers R, Goncharov AD, Berry M, Eid J, et al. Outpatient compared with inpatient preinduction cervical ripening using a synthetic osmotic dilator: a randomized clinical trial. Obstet Gynecol 2022;140.

Appendix 2. Kaplan-Meier curves with censoring for cesarean delivery comparing the time from admission to reaching active labor between outpatient and inpatient groups.



Saad AF, Gavara R, Beavers R, Goncharov AD, Berry M, Eid J, et al. Outpatient compared with inpatient preinduction cervical ripening using a synthetic osmotic dilator: a randomized clinical trial. Obstet Gynecol 2022;140.

## Appendix 3. Patient satisfaction survey.



Saad AF, Gavara R, Beavers R, Goncharov AD, Berry M, Eid J, et al. Outpatient compared with inpatient preinduction cervical ripening using a synthetic osmotic dilator: a randomized clinical trial. Obstet Gynecol 2022;140.