

Supplemental Table: Prediction of clinical watchers alone and clinical watchers plus $IDO_2 < 5$. CI=confidence interval

	Clinical Watcher Only 12-Hour Time Periods, N=4,017		Clinical Watchers & $IDO_2 < 5$ 12 Hour Time Periods, N=2958	
	Clinical Deterioration Events	CPR Events	Clinical Deterioration Events	CPR Events
N	93	23	77	23
Prevalence	0.9%	0.2%	0.9%	0.2%
Positive Likelihood Ratio (95% CI)	2.6 (2.2-3.0)	2.9 (2.2-4.0)	2.9 (2.4-3.5)	3.8 (2.9-5.2)
Negative Likelihood Ratio (95% CI)	0.7 (0.6-0.8)	0.6 (0.5-0.8)	0.7 (0.7-0.8)	0.6 (0.5-0.8)
Positive Predictive Value (95% CI)	2.3% (2.0-2.7%)	0.6% (0.4-0.8%)	2.6% (2.2-3.1%)	0.8% (0.6-1.0%)
Negative Predictive Value (95% CI)	99.4% (99.3-99.4%)	99.9% (99.8-99.9%)	99.3% (99.3-99.4%)	99.9% (99.8-99.9%)
Number Needed to Alert (95% CI)	43 (37-50)	167 (125-250)	38 (32-42)	125 (100-167)