

Supplementary Table. Multivariable Cox model (continuous variables) on ESRD and death

	ESRD HR (95% CI)	P	Death HR (95% CI)	P
Age, 5 yrs	0.94 (0.90-0.98)	0.006	1.56 (1.47-1.67)	<0.001
Male gender	1.21 (0.97-1.53)	0.09	1.21 (0.94-1.56)	0.14
Body Mass Index, kg/m^2	0.97 (0.95-0.99)	0.01	1.00 (0.97-1.02)	0.78
Diabetes Mellitus	1.14 (0.88-1.46)	0.32	1.29 (1.00-1.67)	<0.05
Previous CV event	1.26 (0.99-1.61)	0.06	1.35 (1.06-1.72)	0.02
Smoking	1.02 (0.72-1.45)	0.90	1.18 (0.80-1.74)	0.39
Systolic BP, 5 mmHg	1.00 (0.97-1.03)	0.99	0.97 (0.93-1.00)	0.06
Hemoglobin, g/dL	0.92 (0.86-0.99)	0.04	0.92 (0.85-0.99)	0.03
Phosphate, mg/dL	1.17 (1.03-1.34)	0.02	1.01 (0.86-1.19)	0.88
Cholesterol, mg/dL	1.00 (1.00-1.00)	0.41	1.00 (1.00-1.00)	0.59
Uric Acid, mg/dL	0.95 (0.89-1.01)	0.09	1.09 (1.02-1.16)	0.01
ESRD	-	-	1.50 (1.07-2.12)	0.02
Proteinuria, g/24h ^				0.02
0.0 vs 0.5	-	-	0.66 (0.46-0.95)	
1.0 vs 0.5	-	-	1.12 (0.96-1.31)	
3.0 vs 0.5	-	-	1.19 (0.82-1.73)	
Proteinuria – Stage 3, g/24h ^		<0.001		
0.0 vs 0.5	0.56(0.23-1.39)			
1.0 vs 0.5	1.56(1.14-2.13)			
3.0 vs 0.5	3.03(1.39-6.59)			
Proteinuria – Stage 4, g/24h ^		<0.001		
0.0 vs 0.5	0.57 (0.31-1.06)			
1.0 vs 0.5	1.25 (1.04-1.50)			
3.0 vs 0.5	1.83 (1.16-2.88)			
Proteinuria – Stage 5, g/24h ^		0.96		
0.0 vs 0.5	1.09 (0.55-2.16)			
1.0 vs 0.5	1.05 (0.86-1.28)			
3.0 vs 0.5	1.15 (0.71-1.85)			

Model is stratified for center and CKD stage.

[^]HRs for proteinuria are derived from restricted cubic splines and for specific values.