

Table 1. Linear regression analysis of correlates of suPAR in young patients

Variables	SuPAR			
	Univariable Model		Multivariable Model	
	Regression Coefficient	<i>P</i>	Regression Coefficient	<i>P</i>
Gender (male)	0.07	0.58		
Age (yr)	0.02	0.47		
eGFR (ml/min per 1.73 m ²)	<0.001	0.53		
24 h urine protein (g/24 h)	0.01	0.38		
Serum albumin (g/dl)	-0.003	0.53		
FSGS (yes)	0.23	<0.001	0.23	<0.001

Table 2. Linear regression analysis of correlates of suPAR in adult patients

Variables	SuPAR			
	Univariable Model		Multivariable Model	
	Regression Coefficient	<i>P</i>	Regression Coefficient	<i>P</i>
Gender (male)	-0.13	0.01		
Age (yr)	<0.001	0.87		
eGFR (ml/min per 1.73 m ²)	<0.001	0.92		
24 h urine protein (g/24 h)	0.02	0.04		
Serum albumin (g/dl)	<0.001	0.96		
FSGS (yes)	0.30	<0.001	0.30	<0.001

Table 3. SuPAR levels comparisons between steroid sensitive and non-sensitive patients in the tip group and the other variants group

	Steroid sensitive	Steroid non-sensitive	<i>P</i>
Tip variant	3114 (2189-5391)	2750 (2266-3500)	0.03
Other variants	4143(2966-5770)	2417 (1926-3292)	0.001