Supplemental material is neither peer-reviewed nor thoroughly edited by CJASN. The authors alone are responsible for the accuracy and presentation of the material.

Removal of protein-bound uremic toxins during hemodialysis using a binding competitor

Magdalena Madero, MD[1], Karla B. Cano, MD[1], Israel Campos, MD[2], Xia Tao, PhD[2],

Vaibhav Maheshwari, PhD[2], Jillian Brown, BS[3], Beatriz Cornejo, MD[1], Garry Handelman, PhD[3],

Stephan Thijssen, MD[2], Peter Kotanko, MD[2,4]

[1] Instituto Nacional de Cardiología Ignacio Chávez, Ciudad de México, México

[2] Renal Research Institute, New York, NY, USA

[3] University of Massachusetts, Lowell, MA, USA

[4] Icahn School of Medicine at Mount Sinai, New York, NY, USA

Supplemental material

Table of Contents

Figure 1. Indoxyl sulfate arterial serum concentration over time 3
Figure 2. p−Cresyl sulfate arterial serum concentration over time 4
Figure 3. Trytophan arterial serum concentration over time 5
Figure 4. Urea arterial serum concentration over time 6
Figure 5. Creatinine arterial serum concentration over time 7
Figure 6. Ibuprofen arterial serum concentration over time 8
Figure 7. Indoxyl sulfate venous serum concentration over time 9
Figure 8. p−Cresyl sulfate venous serum concentration over time 10
Figure 9. Trytophan venous serum concentration over time 11
Figure 10. Urea venous serum concentration over time 12
Figure 11. Creatinine venous serum concentration over time 13
Figure 12. Ibuprofen venous serum concentration over time 14
Figure 13. Indoxyl sulfate dialysate outlet concentration over time 15
Figure 14. p−Cresyl sulfate dialysate outlet concentration over time 16
Figure 15. Trytophan dialysate outlet concentration over time 17
Figure 16. Urea dialysate outlet concentration over time 18


Supplemental material is neither peer-reviewed nor thoroughly edited by CJASN. The authors alone are responsible for the accuracy and presentation of the material.

Figure 17. Creatinine dialysate outlet concentration over time 19

Figure 18. Ibuprofen dialysate outlet concentration over time 20

Supplemental material is neither peer-reviewed nor thoroughly edited by CJASN. The authors alone are responsible for the accuracy and presentation of the material.


Figure 1. Indoxyl sulfate arterial serum concentration over time



The shaded area between minute 21 and minute 40 indicates the ibuprofen infusion period. Blue bars represent the median over the corresponding period or time point.

Supplemental material is neither peer-reviewed nor thoroughly edited by CJASN. The authors alone are responsible for the accuracy and presentation of the material.


Figure 2. p–Cresyl sulfate arterial serum concentration over time



The shaded area between minute 21 and minute 40 indicates the ibuprofen infusion period. Blue bars represent the median over the corresponding period or time point.

Supplemental material is neither peer-reviewed nor thoroughly edited by CJASN. The authors alone are responsible for the accuracy and presentation of the material.


Figure 3. Trytophan arterial serum concentration over time



The shaded area between minute 21 and minute 40 indicates the ibuprofen infusion period. Blue bars represent the median over the corresponding period or time point.

Supplemental material is neither peer-reviewed nor thoroughly edited by CJASN. The authors alone are responsible for the accuracy and presentation of the material.


Figure 4. Urea arterial serum concentration over time



The shaded area between minute 21 and minute 40 indicates the ibuprofen infusion period. Blue bars represent the median over the corresponding period or time point.

Supplemental material is neither peer-reviewed nor thoroughly edited by CJASN. The authors alone are responsible for the accuracy and presentation of the material.


Figure 5. Creatinine arterial serum concentration over time



The shaded area between minute 21 and minute 40 indicates the ibuprofen infusion period. Blue bars represent the median over the corresponding period or time point.

Supplemental material is neither peer-reviewed nor thoroughly edited by CJASN. The authors alone are responsible for the accuracy and presentation of the material.


Figure 6. Ibuprofen arterial serum concentration over time



The shaded area between minute 21 and minute 40 indicates the ibuprofen infusion period. Blue bars represent the median over the corresponding period or time point.

Supplemental material is neither peer-reviewed nor thoroughly edited by CJASN. The authors alone are responsible for the accuracy and presentation of the material.


Figure 7. Indoxyl sulfate venous serum concentration over time



The shaded area between minute 21 and minute 40 indicates the ibuprofen infusion period. Blue bars represent the median over the corresponding period or time point.

Supplemental material is neither peer-reviewed nor thoroughly edited by CJASN. The authors alone are responsible for the accuracy and presentation of the material.


Figure 8. p–Cresyl sulfate venous serum concentration over time



The shaded area between minute 21 and minute 40 indicates the ibuprofen infusion period. Blue bars represent the median over the corresponding period or time point.

Supplemental material is neither peer-reviewed nor thoroughly edited by CJASN. The authors alone are responsible for the accuracy and presentation of the material.


Figure 9. Trytophan venous serum concentration over time



The shaded area between minute 21 and minute 40 indicates the ibuprofen infusion period. Blue bars represent the median over the corresponding period or time point.

Supplemental material is neither peer-reviewed nor thoroughly edited by CJASN. The authors alone are responsible for the accuracy and presentation of the material.


Figure 10. Urea venous serum concentration over time



The shaded area between minute 21 and minute 40 indicates the ibuprofen infusion period. Blue bars represent the median over the corresponding period or time point.

Supplemental material is neither peer-reviewed nor thoroughly edited by CJASN. The authors alone are responsible for the accuracy and presentation of the material.


Figure 11. Creatinine venous serum concentration over time



The shaded area between minute 21 and minute 40 indicates the ibuprofen infusion period. Blue bars represent the median over the corresponding period or time point.

Supplemental material is neither peer-reviewed nor thoroughly edited by CJASN. The authors alone are responsible for the accuracy and presentation of the material.


Figure 12. Ibuprofen venous serum concentration over time



The shaded area between minute 21 and minute 40 indicates the ibuprofen infusion period. Blue bars represent the median over the corresponding period or time point.

Supplemental material is neither peer-reviewed nor thoroughly edited by CJASN. The authors alone are responsible for the accuracy and presentation of the material.


Figure 13. Indoxyl sulfate dialysate outlet concentration over time



The shaded area between minute 21 and minute 40 indicates the ibuprofen infusion period. Blue bars represent the median over the corresponding period or time point.

Supplemental material is neither peer-reviewed nor thoroughly edited by CJASN. The authors alone are responsible for the accuracy and presentation of the material.


Figure 14. p–Cresyl sulfate dialysate outlet concentration over time



The shaded area between minute 21 and minute 40 indicates the ibuprofen infusion period. Blue bars represent the median over the corresponding period or time point.

Supplemental material is neither peer-reviewed nor thoroughly edited by CJASN. The authors alone are responsible for the accuracy and presentation of the material.


Figure 15. Trytophan dialysate outlet concentration over time



The shaded area between minute 21 and minute 40 indicates the ibuprofen infusion period. Blue bars represent the median over the corresponding period or time point.

Supplemental material is neither peer-reviewed nor thoroughly edited by CJASN. The authors alone are responsible for the accuracy and presentation of the material.


Figure 16. Urea dialysate outlet concentration over time



The shaded area between minute 21 and minute 40 indicates the ibuprofen infusion period. Blue bars represent the median over the corresponding period or time point.

Supplemental material is neither peer-reviewed nor thoroughly edited by CJASN. The authors alone are responsible for the accuracy and presentation of the material.


Figure 17. Creatinine dialysate outlet concentration over time



The shaded area between minute 21 and minute 40 indicates the ibuprofen infusion period. Blue bars represent the median over the corresponding period or time point.

Supplemental material is neither peer-reviewed nor thoroughly edited by CJASN. The authors alone are responsible for the accuracy and presentation of the material.

Figure 18. Ibuprofen dialysate outlet concentration over time



The shaded area between minute 21 and minute 40 indicates the ibuprofen infusion period. Blue bars represent the median over the corresponding period or time point.