

Contributing Authors:

Jeffrey Perl¹, Laura M. Dember², Joanne M Bargman¹, Teri Browne³, David M Charytan⁴, Jennifer E. Flythe⁵, LaTonya J Hickson⁶, Adriana M Hung⁷, Michel Jadoul⁸, Timmy Chang Lee⁹, Klemens B. Meyer¹⁰, Hamid Moradi¹¹, Tariq Shafi¹², Isaac Teitelbaum¹³, Leslie Wong¹⁴, Christopher T Chan¹ On Behalf of the ASN – Dialysis Advisory Group

¹University of Toronto, Department of Medicine, Division of Nephrology, Ontario, Canada, ² Perelman School of Medicine, University of Pennsylvania Department of Medicine, Division of Nephrology, Philadelphia, PA, ³ University of South Carolina College of Social Work, Columbia, SC, ⁴ Harvard Medical School, Brigham & Women's Hospital, Boston, MA, ⁵ University of North Carolina, UNC Kidney Center, Chapel Hill, NC, ⁶Mayo Clinic, Rochester, MN, ⁷Vanderbilt University School of Medicine, Nashville TN, ⁸ Université Catholique de Louvain, Brussels, Belgium, ⁹University of Alabama, ¹⁰ Tufts Medical Center, Boston, MA, ¹¹ University of California, Irvine, ¹²John Hopkins School of Medicine, Baltimore, MD, ¹³University of Colorado, Aurora, CO, ¹⁴ Department of Nephrology and Hypertension Cleveland Clinic, Cleveland, OH