

Title: Transplant Center Patient Navigator to Increase Access to Kidney Transplantation among High-Risk Population

Authors: Mohua Basu^{1,2}; Lisa Petgrave-Nelson²; Kayla D Smith^{1,2}; Jennie P Perryman², Kevin Clark², Stephen O Pastan^{2,3}, Thomas C. Pearson^{1,2}, Christian P. Larsen^{1,2}, Sudeshna Paul⁴, Rachel E. Patzer^{1,2,3,5}

¹ Department of Surgery, Division of Transplantation, Emory University School of Medicine, Atlanta, GA

² Emory Transplant Center, Atlanta, GA

³ Renal Division, Emory University School of Medicine, Atlanta, GA

⁴ Office of Nursing Research, Nell Hodgson Woodruff School of Nursing, Emory University, Atlanta, GA

⁵ Department of Epidemiology, Rollins School of Public Health, Emory University, Atlanta, GA

Table S1: Risk Prediction Model for Waitlisting to Identify Eligible Study Patients (Risk of Waitlisting <40%)

| | Adjusted Odds Ratio (95% Confidence Intervals) |
|--|---|
| Age, years | 0.98 (0.97-0.99) |
| African American (vs. White) Race | 0.69 (0.52-0.91) |
| Body Mass Index ≥ 35 (vs. <35 kg/m²) | 0.47 (0.33-0.67) |
| Etiology of ESRD | |
| Diabetes | 0.64 (0.45-0.91) |
| Hypertension | 0.79 (0.55-1.14) |
| Glomerulonephritis | 1.02 (0.64-1.63) |
| Other | <i>Reference</i> |
| Education Level | |
| Less than High School | <i>Reference</i> |
| Completed High School | 1.85 (1.28-2.68) |
| Some College | 1.87 (1.26-2.77) |
| Completed College or Above | 3.09 (1.98-4.83) |
| Insurance Status | |
| Medicare | <i>Reference</i> |
| Medicaid | 1.43 (1.00-2.06) |
| Private | 2.65 (1.94-3.61) |
| Married (vs. Not Married) | 1.64 (1.25-2.16) |
| English-Speaking (vs. Non-English Speaking) | 5.12 (3.29-7.97) |
| No Dialysis (vs. On Dialysis) | 2.12 (1.55-2.89) |

* C-statistic: 0.90; 95% CI: 0.89-0.91

*Model adjusted for patient variables at time of referral: age, race, body mass index, ESRD etiology, education level, insurance status, marital status, primary language, and dialysis status