Nisha Bansal MD,¹ Dawei Xie PhD,² Kelvin Tao MS,² Jing Chen MD,³ Rajat Deo MD,² Ed Horwitz MD,⁴ Chi-yuan Hsu MD,^{5,12} Radha Krishna Kallem MD,² Martin G. Keane MD,⁶ Claudia M. Lora MD,⁷ Dominic Raj MD,⁸ Elsayed Z. Soliman MD,⁹ Louise Strauss RN,¹⁰ Myles Wolf MD,¹¹ Alan S. Go MD¹²⁻¹⁴ for the CRIC Study^{*}

1 University of Washington, Kidney Research Institute, Department of Medicine, Seattle, WA

2 Center for Clinical Epidemiology and Biostatistics, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA

3 Tulane Medical Center, Department of Epidemiology, New Orleans, LA

4 Metro Health, Department of Medicine, Cleveland, OH

5 University of California, San Francisco, Department of Medicine, San Francisco, CA

6 Temple University, Department of Medicine, Philadelphia, PA

7 University of Illinois, Chicago, Department of Medicine, Chicago, IL

8 George Washington University, Department of Medicine, Washington, DC

9 Epidemiological Cardiology Research Center (EPICARE), Department of Epidemiology and Prevention, and Department of Internal Medicine-Cardiology, Wake Forest School of Medicine, Winston Salem, NC.

10University Hospital, Department of Medicine, Cleveland, OH

11 Northwestern Memorial Hospital, Department of Medicine, Chicago, IL

12 Division of Research, Kaiser Permanente Northern California, Oakland, CA

13 Departments of Epidemiology, Biostatistics and Medicine, University of California, San Francisco, San Francisco, CA

14 Department of Health Research and Policy, Stanford University School of Medicine, Palo Alto, CA