

F

Supplementary Figure 1: Patients enrollment flowchart.

Supplementary Table 1: Clinical characteristics of COVID-19 infected children and 1: 1 matched parent.

Variables	Children $(N=203)$	Parents $(N = 203)$
Gender,n (%)		
Male	122 (60.1)	42 (20.7) <sup>†</sup>
Age (years)		
Mean (SD)	1.3 (0.8)	32.3 (5.2)
Symptom, <i>n</i> (%)		
Fever	198 (97.5)	158 (77.8)
Cough	128 (63.1)	164 (80.8)

Diarrhea	23 (11.3)	20 (9.9)
Vomiting	32 (15.8)	14 (6.9)*
Running nose	13 (6.4)	22 (10.8)
Febrile seizures	3 (1.5)	0 (0)
Severity, <i>n</i> (%)		
Mild	137 (67.5)	202 (99.5)†
Moderate	66 (32.5)	$0 (0)^{\dagger}$
Severe/critical	0 (0)	0 (0)
Asymptomatic	0 (0)	1 (0.5)
Vaccination, <i>n</i> (%) <sup>‡</sup>		
0	203 (100)	85 (41.9)
1	0	14 (6.9)
2	0	61 (30.0)
3	0	43 (21.2)

<sup>\*</sup>Compared to children P < 0.05. †Compared to children P < 0.001.‡Inactivated vaccines include CoronaVac (Sinovac), Covilo (Sinopharm, Beijing), and Inactivated (Vero cells) (Sinopharm, Wuhan). Two shots as full vaccination and three shots as a booster.IQR: Interquartile range; SD: Standard deviation.