

Supplementary Table 1: Characteristics of patients with well-differentiated G-NET at diagnosis.

Characteristic	No. of patients	
	(N = 94)	Percentage
Age at diagnosis, years (mean $\pm$ SD)	55.2 $\pm$ 10.4	
Gender ratio, male:female	36:58	38.3:61.7
Maximal tumor size (mean $\pm$ SD, mm)	8.4 $\pm$ 5.3	
Location of primary/-ies		
Body	67	71.3
Fundus	8	7.5
Antrum	6	6.4
Body + fundus	13	13.8
Number of gastric lesions		
<6	72	76.6
$\geq$ 6	22	23.4
Clinical classification		
Type 1	78	83.0
Type 2	4	4.3
Type 3	12	12.7
Depth of invasion		
Mucosa/submucosa	89	94.7
Muscularis propria	5	5.3
Grade		
Grade 1	66	70.2
Grade 2	26	27.7
Grade 3	2	2.1
Pre-treatment serum gastrin (mean $\pm$ SD, pg/mL)	1479 $\pm$ 532	

Ki 67 (median [IQR], %)	2 (1, 2.5)	
Presence of CAG	73	83.0
Treatment		
Endoscopic resection (EMR/ESD)	63 (14/49)	67.0
SSA	22	23.4
Surgical resection	9	9.6

CAG: Chronic atrophic gastritis; EMR: Endoscopic mucosal resection; ESD: Endoscopic submucosal dissection; G-NET: Gastric neuroendocrine tumors; SD: Standard deviation; SSA: Somatostatin analogue.