Supplemental Digital Content 2. Frequencies of correct answers.

	Physicians	Nurses	Nursing assistants	Other university educated staff	Administrative staff	Total	Missing
	487 (16)	1241 (41)	861 (28)	337 (11)	103 (3)	3029 (99)	15 (0.5)
What is the first thing you do when you see a person collapse?	192 (39)	684 (55)	353 (41)	159 (47)	33 (33)	1421 (47)	11 (0.4)
2. How long time should inspection of an unconscious patient take?	336 (69)	818 (66)	563 (66)	168 (50)	45 (45)	1930 (64)	17 (0.6)
3. Which first aid equipment should you prioritise?	443 (91)	1166 (94)	799 (94)	305 (91)	95 (92)	2808 (93)	12 (0.4)
4. What is the correct defibrillator electrode placement?	451 (93)	1168 (94)	748 (88)	246 (74)	65 (65)	2678 (89)	18 (0.6)
5. The patient is soaking wet of sweat, what should you do to be able to defibrillate?	397 (82)	1164 (94)	779 (91)	267 (80)	75 (74)	2682 (89)	13 (0.4)
6. How many shocks should be delivered in one sequence?	330 (68)	829 (68)	553 (66)	137 (41)	44 (44)	1893 (63)	37 (1.2)
7. Within which time frame should defibrillation be performed according to guidelines?	32 (7)	208 (17)	108 (13)	22 (7)	2 (2)	372 (12)	40 (1.3)
8. How can you tell if the ventilation is effective?	482 (99)	1239 (99)	853 (99)	333 (99)	102 (100)	3009 (99)	4 (0.1)
9. At which frequency should you perform compressions?	295 (61)	735 (60)	401 (48)	133 (40)	16 (17)	1580 (53)	43 (1.4)
Passed test ≥80% correct	195 (40)	626 (50)	331 (38)	80 (24)	7 (7)	1239 (41)	0 (0)
100% correct	3 (1)	72 (6)	19 (2)	2 (1)	0 (0)	96 (3)	0 (0)

Respondents are presented in frequency n and percent (%).