

Supplementary Table 1: Age and gender distribution of gastroenteritis cases of viral and unknown aetiology among children in Germany*

Description	ICD-10 code	Age [y]	Sex	N	CR	95% CI	
Norovirus	A08.1	< 1	m	7,541	361	353	369
		< 1	f	6,136	309	302	317
		1-3	m	12,034	188	185	191
		1-3	f	10,538	173	170	177
		4-10	m	4,201	26	25	27
		4-10	f	3,738	24	24	25
	AR total NV	0-10	m,f	44,188	89	88	89
Adenovirus	A08.2	< 1	m	3,102	149	143	154
		< 1	f	2,761	139	134	144
		1-3	m	4,002	62	61	64
		1-3	f	3,538	58	56	60
		4-10	m	1,271	8	7	8
		4-10	f	1,164	8	7	8
	AR total AV	0-10	m,f	15,838	32	32	33
Viral, other	A08.3-A08.5	< 1	m	4,321	207	201	213
		< 1	f	3,738	188	182	195
		1-3	m	6,609	103	101	106
		1-3	f	5,762	95	92	97
		4-10	m	5,058	31	30	32
		4-10	f	4,601	30	29	31
	AR total, other	0-10	m,f	30,089	62	61	62
AGE, nonspecific	A09*	< 1	m	38,514	1,844	1,826	1,863
		< 1	f	33,263	1,677	1,659	1,695
		1-3	m	61,565	961	953	969
		1-3	f	52,998	872	864	879
		4-10	m	48,69	301	298	303
		4-10	f	42,823	278	276	281
	AR total, nonspecific	0-10	m,f	277,853	569	567	571

*Number (N) of infections by the pathogen indicated, the crude rates (CR) by age and gender, and the age standardized rates (AR = total) per 100,000 person-years with 95% confidence limits (CI).