SDC 1. Characteristics of Pediatric Clostridium difficile Cases, 2010-2014

Characteristic	n/Total (%)
Sex, male	89/175 (50.9)
Race	
White	138/169 (81.7)
Black	10/169 (5.9)
Hispanic	12/169 (7.1)
Mixed	7/169 (4.1)
Asian	2/169 (1.2)
Inflammatory Bowel Disease (IBD) [^]	23/169 (13.6)
Immunosuppressed [^]	59/175 (33.7)
Oncology	26/59 (44.1)
Leukemia	14/59 (23.7)
Other Cancer	12/59 (20.3)
IBD	15/59 (25.4)
Crohn's Disease	11/59 (18.6)
Ulcerative Colitis	4/59 (6.8)
Transplant	8/59 (13.6)
Bone Marrow	7/59 (11.9)
Liver	1/59 (1.7)
Other [†]	10/59 (16.9)
Hospital-Acquired*	87/215 (40.5)
Recent Antibiotics*	103/209 (49.3)

[^]Data are shown by patient

[†]3 patients with Cystic Fibrosis, 2 unknown, and 1 each with familial hemophagocytic lymphohistiocytosis, lupus, panhypopituitarism, burn injury, and DiGeorge Syndrome.

^{*}Data are shown by case