

The following content was supplied by the authors as supporting material and has not been copy-edited or verified by JBJS.

Appendix Table 1. Medical comorbidities of 7979 patients who underwent hip fracture by cephalomedullary nail type (2014-2019).

Characteristic, n (%) unless specified	TFN (N=4007)	TFNA (N=3972)	Total (N=7979)
AIDS	13 (0.3)	4 (0.1)	17 (0.2)
Alcohol abuse	222 (5.5)	240 (6.0)	462 (5.8)
Chronic blood loss anemia	154 (3.8)	173 (4.4)	327 (4.1)
Chronic pulmonary disease	979 (24.4)	1093 (27.5)	2072 (26.0)
Coagulopathy	446 (11.1)	471 (11.9)	917 (11.5)
Congestive Heart Failure	679 (16.9)	735 (18.5)	1414 (17.7)
Diabetes, complicated	953 (23.8)	1080 (27.2)	2033 (25.5)
Diabetes, uncomplicated	288 (7.2)	507 (12.8)	795 (10.0)
Deficiency anemias	1263 (31.5)	1387 (34.9)	2650 (33.2)
Depression	425 (10.6)	681 (17.1)	1106 (13.9)
Drug abuse	74 (1.9)	77 (1.9)	151 (1.9)
Fluid and electrolyte disorders	967 (24.1)	937 (23.6)	1904 (23.9)
Hypertension	3002 (74.9)	3012 (75.8)	6014 (75.4)
Hypothyroidism	804 (20.1)	823 (20.7)	1627 (20.4)
Liver disease	176 (4.4)	206 (5.2)	382 (4.8)
Lymphoma	52 (1.3)	57 (1.4)	109 (1.4)
Obesity	466 (11.6)	576 (14.5)	1042 (13.1)
Other neurological disorders	862 (21.5)	916 (23.1)	1778 (22.3)
Paralysis	221 (5.52)	203 (5.1)	424 (5.3)
Peptic ulcer disease/bleeding	22 (0.6)	34 (0.9)	56 (0.7)
Peripheral vascular disease	2257 (56.3)	2185 (55.0)	4442 (55.7)
Psychoses	502 (12.5)	387 (9.7)	889 (11.1)
Pulmonary circulation disease	114 (2.9)	49 (1.2)	163 (2.0)
Renal failure	1293 (32.3)	1282 (32.3)	2575 (32.3)
RA/collagen vascular diseases	204 (5.1)	214 (5.4)	418 (5.2)
Tumor without metastasis	99 (2.5)	190 (4.8)	289 (3.6)
Valvular Disease	532 (13.3)	623 (15.7)	1155 (14.5)
Weight loss	518 (12.9)	375 (9.4)	893 (11.2)

AIDS=acquired immunodeficiency syndrome. RA=rheumatoid arthritis.

* Comorbidities were identified using the Elixhauser algorithm.