

Categorical variables	PROMIS Physical Function (p value)	PROMIS Pain Interference (p value)	FFI Pain (p value)	FFI Disability (p value)
Sex	.649	.714	.085	.677
Race	.111	.027*	.086	.282
Prior ankle instability operation	.927	.733	.660	.482
History of major trauma	.554	.515	.649	.680
Athlete	.001*	.004*	.011*	.001*
Comorbidities and substance use				
Smoking	.239	.164	.256	.152
Diabetes	.769	.492	.930	.854
RA	.033*	.115	.086	.775
Collagen pathologies	.080	.838	.332	.335
Generalized Hyperlaxity	.675	.928	.521	.515
Anterior Drawer Positive	.739	.712	.818	.545
Concomitant Pathologies				
Talar OCD	.143	.513	.269	.127
Tibial OCD	.425	.098	.228	.109
Peroneal Tendinopathy	.979	.376	.712	.550
Deltoid ligament injury	1.000	.944	.680	.320
Talar Tilt Varus	.353	.096	.102	.284
Preoperative ankle osteoarthritis	.727	.658	.552	.950
Anterior impingement	.126	.403	.235	.527
Posterior Impingement	.502	.209	.490	.662
Adjuvant Operations Performed				
Ankle arthroscopy	.469	.839	.936	.624
Calcaneal Osteotomy	.527	.120	.122	.376
Peroneal Debridement	.646	.623	.902	.946
Peroneal Groove Deepening	.681	.626	.979	.936
Peroneal Tenodesis	.709	.374	.736	.037*
Fibularis Longus to Brevis Transfer	.826	.584	.619	.994
Complications				
All complications	.084	.286	.097	.119
Wound complication	.484	.439	.415	.175
Sural Nerve Injury	.179	.873	.537	.597
Reflex Sympathetic Dystrophy	.831	.427	.090	.425
Reoperations	.207	.408	.239	.182
Recurrence of instability				
Failure	.163	.128	.266	.570
Continuous variables	PROMIS Physical Function R value (p value)	PROMIS Pain Interference R value (p value)	FAAM ADL R value (p value)	FAAM Sports R value (p value)
Age	-.224 (p=.035*)	-.178 (p=.096)	-.160 (p=.135)	-.134 (p=.237)
BMI	-.184 (p=.085)	.095 (p=.374)	-.210 (p=.048)*	-.103 (p=.364)
Time to Survey	.255 (p=.016)*	-.246 (p=.020)*	.337 (p=.001)*	.247 (p=.027)*
Resilience	.430 (p<.001)*	-.378 (p<.001)*	.424 (p<.001)*	.401 (p=.001)*