Appendix A: iKNOW PD Survey 1 Assessment of Facilitators and Barriers to Outcome Measurement in People with PD

Integrating Knowledge Translation Tools for Outcome Measurement in PD (iKNOW-PD)

Instructions: Clinical researchers from University of Michigan-Flint and Wayne State University are partnering with Genesys Integrated Physical Therapy in order to continue to improve the rehabilitation services provided to patients with Parkinson's Disease. We are interested in further understanding what your attitudes and behaviors are regarding using standardized assessments in people with Parkinson's Disease (PD). Please take 10 minutes to complete this anonymous survey.

Read each statement and rate from strongly agree to strongly disagree.

| | Strongly agree | Somewhat agree | Neither agree nor disagree | Somewhat disagree | Strongly disagree |
|--|-------------------|-------------------|----------------------------------|----------------------|----------------------|
| I have sufficient knowledge to use standardized assessments in patients with PD. | 0 | 0 | 0 | 0 | 0 |
| I would like to know more about the use of standardized assessments in patients with PD before I decide to use them. | 0 | 0 | 0 | 0 | 0 |
| I am free to make my own clinical decision when using standardized assessments in patients with PD. | 0 | 0 | 0 | 0 | 0 |
| In general, I resist using standardized assessments in patients with PD. | 0 | 0 | 0 | 0 | 0 |
| Using standardized assessments improves the quality of patient care in patients with PD. | 0 | 0 | 0 | 0 | 0 |
| The use of standardized assessments motivates my patients with PD. | 0 | 0 | 0 | 0 | 0 |

Read each statement and rate from strongly agree to strongly disagree.

| | Strongly agree | Somewhat agree | Neither agree nor disagree | Somewhat disagree | Strongly disagree |
|--|-------------------|-------------------|----------------------------------|----------------------|----------------------|
| I use standardized assessments to educate the patient/family. | 0 | 0 | 0 | 0 | 0 |
| I feel confident I choose the best standardized assessment for patient care. | 0 | 0 | 0 | 0 | 0 |
| There are so many different assessment; I do not know which ones to use. | 0 | 0 | 0 | 0 | 0 |
| I find using standardized assessment a problem because I have had no training in using them. | 0 | 0 | 0 | 0 | 0 |
| Patients value the use of standardized assessments to gain insight into their functioning. | 0 | 0 | 0 | 0 | 0 |
| The kinds of patients I treat are unsuitable for the use of standardized assessments. | 0 | 0 | 0 | 0 | 0 |

Read each statement and rate from strongly agree to strongly disagree.

| | Strongly agree | Somewhat agree | Neither agree nor disagree | Somewhat disagree | Strongly disagree |
|---|-------------------|-------------------|----------------------------------|----------------------|----------------------|
| Co-workers (PTs) support the use of standardized assessments. | 0 | 0 | 0 | 0 | 0 |
| My supervisor supports the use of standardized assessments. | 0 | 0 | 0 | 0 | 0 |

| 0 | 0 | 0 | 0 | 0 |
|---|---|---|---|---|
| 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 |
| | | | | |

What factors may increase the use of outcome measures in the patients you see with PD? What outcome measures in PD do you want to use more frequently? What barriers may exist to prevent you from using outcome measures in the patients you see with PD?