

**Supplementary Table S1:** Structural changes of kidney tissue in Wistar and MWF rats.

	<b>Wistar 50</b>	<b>MWF 50</b>	<b>MWF 60</b>	<b>MWF+LIS 60</b>	<b>MWF+LOS 60</b>
Tubular atrophy (score 0-3)	0.25±0.26 <sup>°°°</sup>	2.22±0.26	2.50±0.43	1.17±0.25 <sup>°°°</sup>	1.46±0.14 <sup>°°°</sup>
Interstitial fibrosis (score 0-3)	0.04±0.14 <sup>°°°</sup>	2.28±0.26	2.44±0.46	1.04±0.50 <sup>°°°</sup>	1.50±0.56 <sup>°°°</sup>
Interstitial Inflammation (score 0-3)	0.08±0.19 <sup>°°°</sup>	2.33±0.25	2.61±0.33	0.92±0.42 <sup>°°°</sup>	1.38±0.53 <sup>°°°</sup>
Luminal casts (score 0-3)	0.08±0.19 <sup>°°°</sup>	1.50±0.50	2.11±0.70	0.71±0.45 <sup>°°°</sup>	1.08±0.63 <sup>**</sup>

° P<0.05 vs MWF 50

°° P<0.01 vs MWF 50

\*\* P<0.01 vs MWF 60