

Supplemental Table S1. Adjusted hazards ratios overall and after initiation of maintenance renal replacement therapy in patients with end-stage renal disease due to multiple myeloma

Characteristic	Overall	P	AHR, year 1	P	AHR, year 2	P	AHR, year 3	P
All	2.05 (1.99-2.11)	0	2.64 (2.53-2.75)	0	1.87 (1.74-2)	0	1.59 (1.45-1.73)	0
Age, yrs.								
< 40	2.61 (1.83-3.72)	0	2.54 (1.48-4.35)	< 0.001	3.71 (1.46-9.4)	< 0.01	3.52 (1.24-9.99)	0.02
40-64	2.05 (2-2.12)	0	2.64 (2.54-2.75)	0	1.87 (1.75-2)	0	1.59 (1.45-1.74)	0
≥ 65	1.99 (1.92-2.06)	0	2.43 (2.31-2.54)	0	1.86 (1.71-2.02)	0	1.57 (1.4-1.75)	0
Sex								
Male	2.1 (2.02-2.19)	0	2.81 (2.66-2.97)	0	1.8 (1.64-1.97)	0	1.58 (1.4-1.79)	0
Female	1.98 (1.9-2.07)	0	2.43 (2.29-2.58)	0	1.96 (1.77-2.18)	0	1.59 (1.39-1.81)	0
Race/ethnicity								
White	1.92 (1.86-1.99)	0	2.48 (2.37-2.6)	0	1.74 (1.61-1.88)	0	1.41 (1.27-1.57)	0
Black	2.39 (2.23-2.55)	0	3.08 (2.8-3.4)	0	2.21 (1.89-2.58)	0	2.12 (1.75-2.58)	0
Hispanic	2.63 (2.32-2.99)	0	3.67 (3.04-4.42)	0	2.29 (1.73-3.02)	0	1.56 (1.07-2.28)	0.02
Other	2.32 (1.9-2.83)	0	2.61 (1.96-3.48)	0	2.5 (1.59-3.93)	< 0.001	4.92 (2.55-9.52)	0

Note: Hazards ratios are adjusted for age, sex, race, and ethnicity. Values are expressed as AHR (95% confidence interval).

AHR, adjusted hazard ratio.